



wwPDB EM Validation Summary Report ⓘ

Nov 3, 2024 – 07:07 AM EST

PDB ID : 487D
Title : SEVEN RIBOSOMAL PROTEINS FITTED TO A CRYO-ELECTRON MICROSCOPIC MAP OF THE LARGE 50S SUBUNIT AT 7.5 ANGSTROMS RESOLUTION
Authors : Brimacombe, R.; Mueller, F.
Deposited on : 2000-02-23
Resolution : 7.50 Å (reported)
Based on initial models : 1CSV, 1CSX, 1CSW

This is a wwPDB EM Validation Summary Report for a publicly released PDB/EMDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 7.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	H	224	89% 10% .
2	I	135	53% 36% 10%
3	J	164	77% 22% .
4	K	149	46% 44% 9% .
5	L	133	38% 44% 12% 5%
6	M	122	70% 27% .
7	N	94	81% 18% .

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8778 atoms, of which 953 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 50S ribosomal protein L1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	H	224	Total	C	N	O	S	0	0
			1712	1083	311	314	4		

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	179	CYS	SER	conflict	UNP P27150

- Molecule 2 is a protein called 50S ribosomal protein L2.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	I	135	Total	C	N	O	S	Se	0	0
			1024	645	187	188	1	3		

- Molecule 3 is a protein called 50S ribosomal protein L6.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	J	164	Total	C	N	O	S	0	0
			1251	787	225	237	2		

- Molecule 4 is a protein called 50S ribosomal protein L9.

Mol	Chain	Residues	Atoms					AltConf	Trace	
4	K	149	Total	C	H	N	O	S	0	0
			1435	729	287	206	212	1		

- Molecule 5 is a protein called 50S ribosomal protein L11.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	L	133	Total	C	N	O	S	0	0
			1000	642	169	183	6		

- Molecule 6 is a protein called 50S ribosomal protein L14.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	M	122	Total	C	N	O	S	0	0
			937	585	180	169	3		

- Molecule 7 is a protein called 50S ribosomal protein L25.

Mol	Chain	Residues	Atoms						AltConf	Trace
7	N	94	Total	C	H	N	O	S	0	0
			1419	479	666	137	134	3		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 50S ribosomal protein L1

Chain H: 



- Molecule 2: 50S ribosomal protein L2

Chain I: 




- Molecule 3: 50S ribosomal protein L6

Chain J: 




- Molecule 4: 50S ribosomal protein L9


Chain K: 




- Molecule 5: 50S ribosomal protein L11



- Molecule 6: 50S ribosomal protein L14



- Molecule 7: 50S ribosomal protein L25



4 Data and refinement statistics

Xtriage (Phenix) and EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	1.00Å 1.00Å 1.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 7.50	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-7.50)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	unknown	Depositor
R, R_{free}	(Not available) , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	8778	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	H	0.48	0/1743	0.74	0/2351
2	I	0.48	0/1034	0.65	0/1388
3	J	0.63	0/1270	0.76	0/1715
4	K	0.63	0/1160	0.79	0/1552
5	L	0.64	0/1016	0.89	6/1366 (0.4%)
6	M	0.86	2/946 (0.2%)	1.18	7/1269 (0.6%)
7	N	0.91	0/766	1.14	4/1025 (0.4%)
All	All	0.65	2/7935 (0.0%)	0.87	17/10666 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
6	M	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	M	86	ILE	C-N	-16.94	0.95	1.34
6	M	87	ILE	C-N	14.61	1.67	1.34

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	M	86	ILE	O-C-N	-15.53	97.85	122.70
6	M	87	ILE	CA-C-N	-14.87	84.49	117.20
6	M	87	ILE	C-N-CA	-14.70	84.95	121.70
6	M	86	ILE	C-N-CA	-13.35	88.31	121.70
6	M	87	ILE	O-C-N	9.88	138.50	122.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
6	M	86	ILE	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1712	0	1763	10	0
2	I	1024	0	1073	46	0
3	J	1251	0	1294	56	0
4	K	1148	287	1230	66	0
5	L	1000	0	1071	142	0
6	M	937	0	995	114	0
7	N	753	666	780	2	0
All	All	7825	953	8206	436	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 436 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:M:79:PHE:CZ	6:M:103:ALA:HB2	1.33	1.64
6:M:79:PHE:CZ	6:M:103:ALA:CB	1.87	1.54
6:M:63:VAL:CB	6:M:106:LEU:HD21	1.34	1.52
6:M:63:VAL:HG12	6:M:106:LEU:CD2	1.38	1.48
3:J:29:LYS:CG	3:J:81:GLY:N	1.78	1.46

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM

entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	222/224 (99%)	211 (95%)	9 (4%)	2 (1%)	14	52
2	I	133/135 (98%)	124 (93%)	7 (5%)	2 (2%)	8	40
3	J	159/164 (97%)	156 (98%)	3 (2%)	0	100	100
4	K	147/149 (99%)	130 (88%)	12 (8%)	5 (3%)	3	21
5	L	131/133 (98%)	101 (77%)	14 (11%)	16 (12%)	0	4
6	M	120/122 (98%)	114 (95%)	4 (3%)	2 (2%)	7	37
7	N	92/94 (98%)	74 (80%)	14 (15%)	4 (4%)	2	17
All	All	1004/1021 (98%)	910 (91%)	63 (6%)	31 (3%)	5	22

5 of 31 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	K	8	ASP
4	K	85	GLU
5	L	21	ALA
5	L	24	VAL
5	L	25	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	177/177 (100%)	169 (96%)	8 (4%)	23	45
2	I	107/104 (103%)	87 (81%)	20 (19%)	1	7
3	J	137/137 (100%)	132 (96%)	5 (4%)	30	50
4	K	119/119 (100%)	100 (84%)	19 (16%)	2	10
5	L	108/108 (100%)	94 (87%)	14 (13%)	3	14
6	M	101/101 (100%)	99 (98%)	2 (2%)	50	68

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
7	N	78/78 (100%)	71 (91%)	7 (9%)	8 24
All	All	827/824 (100%)	752 (91%)	75 (9%)	10 24

5 of 75 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	L	54	LEU
7	N	29	ILE
5	L	62	GLU
5	L	126	LYS
2	I	160	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
4	K	148	GLN
5	L	30	GLN
7	N	51	GLN
6	M	56	GLN
3	J	146	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
3	J	2
6	M	2

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	J	81:GLY	C	82:TYR	N	2.53
1	J	80:LYS	C	81:GLY	N	2.18
1	M	87:ILE	C	88:ARG	N	1.67
1	M	86:ILE	C	87:ILE	N	0.95