



wwPDB NMR Structure Validation Summary Report ⓘ

Mar 20, 2025 – 02:32 PM EDT

PDB ID : 193D
Title : SOLUTION STRUCTURE OF A QUINOMYCIN BISINTERCALATOR-DNA COMPLEX
Authors : Chen, H.; Patel, D.J.
Deposited on : 1994-09-30

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.41.4

i

SOLUTION NMR.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

12486

12%

2 Ensemble composition and analysis ⓘ

This entry contains 4 models.

Cyrange was unable to find well-defined residues.

Error message: No domains could be identified

NmrClust was unable to cluster the ensemble.

Error message: Wrapper check: not enough residues in core to run NmrClust

3 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 656 atoms, of which 252 are hydrogens and 0 are deuteriums.


- Molecule 1 is a DNA chain called DNA (5'-D(*AP*CP*AP*CP*GP*TP*GP*T)-3').

Mol	Chain	Residues	Atoms						Trace
1	A	8	Total	C	H	N	O	P	0
			253	78	92	30	46	7	
1	B	8	Total	C	H	N	O	P	0
			253	78	92	30	46	7	

- Molecule 2 is a protein called QUINOMYCIN.

Mol	Chain	Residues	Atoms						Trace
2	C	8	Total	C	H	N	O	S	0
			112	36	56	8	10	2	

- Molecule 3 is 3-HYDROXYQUINALDIC ACID (three-letter code: HQU) (formula: C₁₀H₇NO₃).




Mol	Chain	Residues	Atoms				
3	C	1	Total	C	H	N	O
			19	10	6	1	2
3	C	1	Total	C	H	N	O
			19	10	6	1	2

4 Residue-property plots [i](#)


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DNA (5'-D(*AP*CP*AP*CP*GP*TP*GP*T)-3')



- Molecule 1: DNA (5'-D(*AP*CP*AP*CP*GP*TP*GP*T)-3')




- Molecule 2: QUINOMYCIN



4.2 Residue scores for the first model from the NMR ensemble

No representative models were identified. Colouring as in section 4.1 above.

- Molecule 1: DNA (5'-D(*AP*CP*AP*CP*GP*TP*GP*T)-3')



- Molecule 1: DNA (5'-D(*AP*CP*AP*CP*GP*TP*GP*T)-3')





● Molecule 2: QUINOMYCIN



5 Refinement protocol and experimental data overview

The models were refined using the following method: *MOLECULAR DYNAMICS, MATRIX RELAXATION*.

Of the 4 calculated structures, 4 were deposited, based on the following criterion: *all calculated structures submitted*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	

No chemical shift data was provided.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HQU, NCY, NYB, CPC, DSN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.41±0.06	3±0/180 (1.5± 0.2%)	2.40±0.08	16±1/276 (6.0± 0.4%)
1	B	1.40±0.03	2±0/180 (1.4± 0.3%)	2.47±0.11	18±1/276 (6.3± 0.4%)
2	C	1.53±0.07	0±0/8 (0.0± 0.0%)	1.19±0.09	0±0/8 (0.0± 0.0%)
All	All	1.41	21/1472 (1.4%)	2.42	136/2240 (6.1%)

5 of 6 unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	B	6	DT	C5-C7	6.79	1.54	1.50	1	2
1	B	8	DT	C5-C7	6.79	1.54	1.50	3	4
1	A	8	DT	C5-C7	6.61	1.54	1.50	3	4
1	A	1	DA	N9-C8	-6.59	1.32	1.37	4	4
1	B	1	DA	N9-C8	-6.33	1.32	1.37	4	4

5 of 59 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	7	DG	O4'-C1'-N9	10.82	115.58	108.00	4	1
1	B	2	DC	O4'-C1'-N1	-10.74	100.48	108.00	4	4
1	B	7	DG	O4'-C1'-N9	10.63	115.44	108.00	4	1
1	A	2	DC	O4'-C1'-N1	-10.54	100.62	108.00	4	4
1	B	3	DA	O4'-C1'-N9	10.46	115.32	108.00	4	2

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	161	92	92	2±0
1	B	161	92	92	1±1
2	C	56	56	54	2±1
3	C	26	12	12	1±0
All	All	1616	1008	1000	18

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

5 of 13 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:C:3:NYB:O	2:C:3:NYB:SG	0.57	2.62	2	4
1:A:5:DG:H1'	3:C:9:HQU:O1	0.47	2.09	3	3
1:B:2:DC:H4'	1:B:3:DA:OP1	0.47	2.10	2	1
1:A:8:DT:OP2	1:A:8:DT:H6	0.46	1.94	4	1
1:B:8:DT:H6	1:B:8:DT:OP2	0.44	1.94	4	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	2/8 (25%)	2±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
All	All	8/32 (25%)	8 (100%)	0 (0%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	C	0	-	-	-
All	All	0	-	-	-

There are no protein residues with a non-rotameric sidechain to report.

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	NYB	C	3	2	8,10,11	1.77±0.07	2±0 (31±6%)
2	CPC	C	8	2	6,8,9	1.41±0.18	1±1 (20±13%)
2	CPC	C	4	2	6,8,9	1.54±0.30	2±1 (25±18%)
2	NCY	C	7	2	5,6,7	2.60±0.25	3±0 (60±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	NYB	C	3	2	7,11,13	1.24±0.07	1±1 (14±10%)
2	CPC	C	8	2	8,12,14	2.55±0.32	4±1 (46±16%)
2	CPC	C	4	2	8,12,14	2.84±0.57	4±1 (50±15%)
2	NCY	C	7	2	5,6,8	2.16±0.21	2±0 (45±8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NYB	C	3	2	-	0±0,8,11,13	-
2	CPC	C	8	2	-	0±0,1,14,17	0±0,1,1,1
2	CPC	C	4	2	-	0±0,1,14,17	0±0,1,1,1
2	NCY	C	7	2	-	0±0,3,6,8	-

5 of 12 unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	C	7	NCY	CB-CA	5.40	1.58	1.53	3	4
2	C	3	NYB	CB-CA	3.76	1.62	1.53	1	4
2	C	8	CPC	CG1-CA	2.89	1.55	1.51	3	2
2	C	3	NYB	CA-N	2.83	1.52	1.47	4	4
2	C	7	NCY	CA-N	2.77	1.52	1.47	3	4

5 of 17 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	C	4	CPC	CG2-CB-CG1	7.33	108.28	120.97	1	4
2	C	8	CPC	CG2-CB-CG1	5.26	111.86	120.97	4	4
2	C	7	NCY	CA-CB-SG	4.35	119.06	114.16	4	2
2	C	7	NCY	CB-CA-N	4.24	105.00	111.19	2	3
2	C	8	CPC	CG1-CB-CA	4.02	61.77	60.04	3	6

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
3	HQU	C	9	2	14,14,15	2.60±0.15	3±1 (21±8%)
3	HQU	C	0	2	14,14,15	2.51±0.05	2±0 (16±3%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
3	HQU	C	9	2	17,19,21	2.81±0.09	6±0 (33±2%)
3	HQU	C	0	2	17,19,21	2.87±0.11	6±0 (33±2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HQU	C	9	2	-	0±0,2,2,4	0±0,2,2,2
3	HQU	C	0	2	-	0±0,2,2,4	0±0,2,2,2

5 of 8 unique bond outliers are listed below. They are sorted according to the Z-score of the worst

occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
3	C	9	HQU	C2-C	9.10	1.38	1.48	4	4
3	C	0	HQU	C2-C	8.26	1.39	1.48	2	4
3	C	9	HQU	C9-N1	2.82	1.33	1.37	1	4
3	C	0	HQU	C9-N1	2.73	1.33	1.37	2	4
3	C	9	HQU	C4-C10	2.52	1.36	1.42	4	2

5 of 12 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
3	C	0	HQU	C3-C2-C	7.50	125.66	120.09	2	4
3	C	9	HQU	C3-C2-C	7.21	125.45	120.09	1	4
3	C	9	HQU	O1-C-C2	7.16	115.38	124.37	4	4
3	C	0	HQU	O1-C-C2	6.10	116.71	124.37	4	4
3	C	0	HQU	C8-C9-C10	5.57	124.61	119.04	2	4

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided