wwPDB EM Validation Summary Report

Sep 28, 2024 – 08:39 pm BST

PDB ID	:	8AIX
EMDB ID	:	EMD-15473
Title	:	Cryo-EM structure of crescentin filaments (wildtype, C2 symmetry and large box)
Authors	:	Liu, Y.; Lowe, J.
Deposited on	:	2022-07-27
Resolution	:	5.80 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references ⓘ) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev113
MolProbity	:	4.02b-467
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 5.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	457	8% 10% 82%
1	B	457	7% 11% 81%
1	E	457	8% 10% 82%
1	F	457	7% 11% 81%
1	G	457	14% 26% 60%
1	H	457	10% 29% 61%
1	K	457	7% 9% 84%
1	L	457	9% 9% 82%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	M	457	9% 23% 68%
1	N	457	13% 21% 66%
1	Q	457	14% 26% 60%
1	R	457	10% 29% 61%
1	U	457	6% 10% 84%
1	V	457	9% 10% 82%
1	W	457	9% 23% 68%
1	X	457	14% 20% 66%
2	C	907	7% 90%
2	D	907	8% 88%
2	I	907	9% 88%
2	J	907	8% 88%
2	O	907	7% 90%
2	P	907	5% 8% 88%
2	S	907	9% 88%
2	T	907	8% 88%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 21924 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Crescentin.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	84	Total	C	N	O	S	0	0
			655	390	134	130	1		
1	B	85	Total	C	N	O	S	0	0
			664	395	136	132	1		
1	G	183	Total	C	N	O	S	0	0
			1400	844	269	286	1		
1	H	179	Total	C	N	O	S	0	0
			1373	829	264	279	1		
1	K	72	Total	C	N	O	S	0	0
			556	341	104	110	1		
1	L	84	Total	C	N	O	S	0	0
			643	393	124	125	1		
1	M	147	Total	C	N	O	S	0	0
			1142	677	234	228	3		
1	N	156	Total	C	N	O	S	0	0
			1207	714	247	243	3		
1	E	84	Total	C	N	O	S	0	0
			655	390	134	130	1		
1	F	85	Total	C	N	O	S	0	0
			664	395	136	132	1		
1	Q	183	Total	C	N	O	S	0	0
			1400	844	269	286	1		
1	R	179	Total	C	N	O	S	0	0
			1373	829	264	279	1		
1	U	72	Total	C	N	O	S	0	0
			556	341	104	110	1		
1	V	84	Total	C	N	O	S	0	0
			643	393	124	125	1		
1	W	147	Total	C	N	O	S	0	0
			1142	677	234	228	3		
1	X	156	Total	C	N	O	S	0	0
			1207	714	247	243	3		

- Molecule 2 is a protein called Crescentin-specific megabody MB13.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	D	112	Total	C	N	O	S	0	0
			869	539	155	170	5		
2	J	111	Total	C	N	O	S	0	0
			860	534	154	167	5		
2	C	94	Total	C	N	O	S	0	0
			724	449	129	142	4		
2	I	112	Total	C	N	O	S	0	0
			869	539	155	170	5		
2	P	112	Total	C	N	O	S	0	0
			869	539	155	170	5		
2	T	111	Total	C	N	O	S	0	0
			860	534	154	167	5		
2	O	94	Total	C	N	O	S	0	0
			724	449	129	142	4		
2	S	112	Total	C	N	O	S	0	0
			869	539	155	170	5		

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Crescentin

Chain A: 8% 10% 82%



- Molecule 1: Crescentin

Chain B: 7% 11% 81%





• Molecule 1: Crescentin

Chain G:



• Molecule 1: Crescentin

Chain H:





• Molecule 1: Crescentin

Chain K:



• Molecule 1: Crescentin

Chain L:





• Molecule 1: Crescentin

Chain M:



• Molecule 1: Crescentin

Chain N:



• Molecule 1: Crescentin

Chain E: 8% 10% 82%



• Molecule 1: Crescentin

Chain F: 7% 11% 81%



• Molecule 1: Crescentin

Chain Q:



• Molecule 1: Crescentin

Chain R:



• Molecule 1: Crescentin

Chain U: 6% 10% 84%



• Molecule 1: Crescentin

Chain V:



• Molecule 1: Crescentin

Chain W:



• Molecule 1: Crescentin

Chain X:



• Molecule 2: Crescentin-specific megabody MB13

Chain D:





• Molecule 2: Crescentin-specific megabody MB13

Chain J: 8% 88%





• Molecule 2: Crescentin-specific megabody MB13

Chain C: 7% 90%



• Molecule 2: Crescentin-specific megabody MB13

Chain I: 9% 88%



• Molecule 2: Crescentin-specific megabody MB13

Chain P: 5% 8% 88%



• Molecule 2: Crescentin-specific megabody MB13

Chain T:



• Molecule 2: Crescentin-specific megabody MB13

Chain O:





• Molecule 2: Crescentin-specific megabody MB13

Chain S: 9% 88%




ILE ALA GLY GLU TYR PRO ASP TYR GLN ARG LYS ILE SER ALA THR ARG ASP GLY LEU LYS VAL THR PHE GLY LYS VAL ARG ALA THR TRP ASP LEU LEU THR SER GLY GLU SER GLU TYR GLN VAL HIS LYS SER LEU PRO VAL GLN THR GLU ILE ASN GLY ASN ARG PHE THR SER LYS
ALA HIS ILE ASN GLY SER THR THR LEU TYR THR THR TYR SER HIS LEU LEU THR ALA GLN GLU VAL SER LYS GLU GLN MET GLN ILE ARG ASP ILE LEU LEU ALA ARG PRO ALA PHE TYR LEU THR ALA SER GLN GLN ARG TRP GLU GLU TYR LEU LYS LYS GLY LEU THR ASN PRO ASP ALA
THR PRO GLU GLN THR ARG VAL ALA VAL LYS ALA ILE GLU THR LEU ASN GLY ASN TRP ARG SER PRO GLY GLY ALA VAL LYS PHE ASN THR VAL THR PRO SER VAL THR GLY ARG TRP PHE SER GLY ASN GLN THR TRP PRO TRP ASP THR TRP LYS GLN ALA PHE ALA MET ALA HIS PHE
ASN PRO ASP ILE ALA LYS GLU ASN ILE ARG ALA VAL PHE SER TRP GLN ILE GLN PRO GLY ASP SER VAL ARG PRO GLN ASP VAL GLY PHE VAL PRO ASP LEU ILE ALA TRP ASN LEU SER PRO GLU ARG GLY GLY ASP GLY GLY ASN TRP ASN THR LYS PRO SER LEU ALA
ALA TRP SER VAL MET GLU VAL TYR ASN VAL THR GLN ASP LYS THR TRP VAL ALA GLU MET TYR PRO LYS LEU VAL ALA TYR HIS ASP TRP TRP LEU ARG ASN ARG ASP HIS ASN GLY ASN GLY VAL PRO GLU GLY TYR GLY ALA THR ARG ASP LYS ALA HIS ASN THR GLU SER GLY MET
LEU PHE THR VAL LYS LYS D789 S790 L791 R792 L793 S794 C795 A796 S797 S798 R799 S800 I801 D802 G803 I804 N805 I806 M807 R808 W809 Y810 R811 Q812 A813 P814 G815 K816 Q817 R818 G819 M820 V821 A822 V823 V824 T825 G826 W827 G828 S829 T830 N831 Y832 V833 D834 S835 V836 R839 F840 I841 I842 S843
R844 D845 S846 A847 K848 D849 T850 V851 Y852 L853 Q854 M855 N856 N857 L858 K859 P860 E861 D862 T863 A864 V865 Y866 N869 A870 I871 Y872 R873 E876 Y877 W878 T882 Q883 V884 T885 V886 S887 SER GLY GLU ASN LEU TYR PHE GLN GLY SER HIS HIS HIS HIS HIS HIS HIS HIS HIS

4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C2	Depositor
Number of particles used	550574	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	34	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	3500	Depositor
Magnification	75000	Depositor
Image detector	FEI FALCON IV (4k x 4k)	Depositor
Maximum map value	1.942	Depositor
Minimum map value	-0.003	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.007	Depositor
Recommended contour level	0.017	Depositor
Map size (Å)	972.0, 972.0, 972.0	wwPDB
Map dimensions	500, 500, 500	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.944, 1.944, 1.944	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	0.36	0/655	0.63	0/875
1	B	0.42	0/664	0.71	0/887
1	E	0.36	0/655	0.63	0/875
1	F	0.42	0/664	0.71	0/887
1	G	0.30	0/1407	0.58	0/1899
1	H	0.32	0/1380	0.56	1/1862 (0.1%)
1	K	0.27	0/560	0.44	0/755
1	L	0.32	0/648	0.48	0/872
1	M	0.32	0/1143	0.56	0/1528
1	N	0.38	0/1208	0.65	1/1615 (0.1%)
1	Q	0.30	0/1407	0.58	0/1899
1	R	0.32	0/1380	0.56	0/1862
1	U	0.27	0/560	0.44	0/755
1	V	0.32	0/648	0.48	0/872
1	W	0.32	0/1143	0.56	0/1528
1	X	0.38	0/1208	0.65	1/1615 (0.1%)
2	C	0.45	0/734	0.63	1/988 (0.1%)
2	D	0.44	0/883	0.61	0/1192
2	I	0.46	0/883	0.63	0/1192
2	J	0.45	0/874	0.64	0/1180
2	O	0.45	0/734	0.63	1/988 (0.1%)
2	P	0.44	0/883	0.61	0/1192
2	S	0.47	0/883	0.63	0/1192
2	T	0.45	0/874	0.64	0/1180
All	All	0.37	0/22078	0.60	5/29690 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	N	403	ARG	CB-CG-CD	5.41	125.66	111.60
1	X	403	ARG	CB-CG-CD	5.38	125.59	111.60
2	O	812	GLN	C-N-CA	-5.04	109.10	121.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
2	C	812	GLN	C-N-CA	-5.03	109.14	121.70
1	H	164	LEU	CA-CB-CG	5.01	126.83	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	655	0	667	70	0
1	B	664	0	674	67	0
1	E	655	0	667	68	0
1	F	664	0	674	68	0
1	G	1400	0	1389	136	0
1	H	1373	0	1364	178	0
1	K	556	0	557	43	0
1	L	643	0	646	48	0
1	M	1142	0	1151	142	0
1	N	1207	0	1214	117	0
1	Q	1400	0	1389	137	0
1	R	1373	0	1364	176	0
1	U	556	0	557	45	0
1	V	643	0	646	50	0
1	W	1142	0	1151	138	0
1	X	1207	0	1214	114	0
2	C	724	0	697	103	0
2	D	869	0	846	92	0
2	I	869	0	846	123	0
2	J	860	0	837	97	0
2	O	724	0	697	100	0
2	P	869	0	846	89	0
2	S	869	0	846	122	0
2	T	860	0	837	95	0
All	All	21924	0	21776	2130	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 49.

The worst 5 of 2130 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:O:802:ASP:OD1	2:O:844:ARG:HD3	1.52	1.09
2:C:802:ASP:OD1	2:C:844:ARG:HD3	1.52	1.08
2:J:832:TYR:HB2	2:J:836:VAL:HG21	1.42	1.01
1:A:404:ARG:NH2	1:X:320:ARG:HD2	1.76	1.01
2:I:805:ASN:HD21	2:I:872:TYR:HD1	1.01	1.01

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	82/457 (18%)	76 (93%)	6 (7%)	0	100	100
1	B	83/457 (18%)	72 (87%)	11 (13%)	0	100	100
1	E	82/457 (18%)	76 (93%)	6 (7%)	0	100	100
1	F	83/457 (18%)	72 (87%)	11 (13%)	0	100	100
1	G	181/457 (40%)	166 (92%)	15 (8%)	0	100	100
1	H	177/457 (39%)	166 (94%)	11 (6%)	0	100	100
1	K	70/457 (15%)	67 (96%)	3 (4%)	0	100	100
1	L	82/457 (18%)	77 (94%)	5 (6%)	0	100	100
1	M	145/457 (32%)	139 (96%)	6 (4%)	0	100	100
1	N	154/457 (34%)	151 (98%)	3 (2%)	0	100	100
1	Q	181/457 (40%)	166 (92%)	15 (8%)	0	100	100
1	R	177/457 (39%)	166 (94%)	11 (6%)	0	100	100
1	U	70/457 (15%)	67 (96%)	3 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	V	82/457 (18%)	77 (94%)	5 (6%)	0	100	100
1	W	145/457 (32%)	139 (96%)	6 (4%)	0	100	100
1	X	154/457 (34%)	151 (98%)	3 (2%)	0	100	100
2	C	86/907 (10%)	70 (81%)	16 (19%)	0	100	100
2	D	108/907 (12%)	88 (82%)	20 (18%)	0	100	100
2	I	108/907 (12%)	81 (75%)	27 (25%)	0	100	100
2	J	107/907 (12%)	80 (75%)	27 (25%)	0	100	100
2	O	86/907 (10%)	70 (81%)	16 (19%)	0	100	100
2	P	108/907 (12%)	88 (82%)	20 (18%)	0	100	100
2	S	108/907 (12%)	81 (75%)	27 (25%)	0	100	100
2	T	107/907 (12%)	80 (75%)	27 (25%)	0	100	100
All	All	2766/14568 (19%)	2466 (89%)	300 (11%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	62/345 (18%)	62 (100%)	0	100	100
1	B	63/345 (18%)	63 (100%)	0	100	100
1	E	62/345 (18%)	62 (100%)	0	100	100
1	F	63/345 (18%)	63 (100%)	0	100	100
1	G	140/345 (41%)	137 (98%)	3 (2%)	48	66
1	H	137/345 (40%)	136 (99%)	1 (1%)	81	87
1	K	55/345 (16%)	54 (98%)	1 (2%)	54	71
1	L	63/345 (18%)	63 (100%)	0	100	100
1	M	110/345 (32%)	107 (97%)	3 (3%)	40	58
1	N	116/345 (34%)	115 (99%)	1 (1%)	75	83

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	Q	140/345 (41%)	137 (98%)	3 (2%)	48 66
1	R	137/345 (40%)	136 (99%)	1 (1%)	81 87
1	U	55/345 (16%)	54 (98%)	1 (2%)	54 71
1	V	63/345 (18%)	63 (100%)	0	100 100
1	W	110/345 (32%)	107 (97%)	3 (3%)	40 58
1	X	116/345 (34%)	115 (99%)	1 (1%)	75 83
2	C	78/749 (10%)	77 (99%)	1 (1%)	65 77
2	D	94/749 (13%)	94 (100%)	0	100 100
2	I	94/749 (13%)	93 (99%)	1 (1%)	70 80
2	J	93/749 (12%)	93 (100%)	0	100 100
2	O	78/749 (10%)	77 (99%)	1 (1%)	65 77
2	P	94/749 (13%)	94 (100%)	0	100 100
2	S	94/749 (13%)	93 (99%)	1 (1%)	70 80
2	T	93/749 (12%)	93 (100%)	0	100 100
All	All	2210/11512 (19%)	2188 (99%)	22 (1%)	71 82

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	R	99	ARG
1	W	380	ARG
1	W	337	ARG
1	W	403	ARG
1	M	380	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
2	P	856	ASN
1	R	100	GLN
1	R	40	GLN
1	V	75	GLN
1	L	75	GLN

5.3.3 RNA

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Map visualisation

This section contains visualisations of the EMDB entry EMD-15473. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections

6.1.1 Primary map



X



Y



Z

6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map



X Index: 250



Y Index: 250

Z Index: 250

6.2.2 Raw map



X Index: 250



Y Index: 250



Z Index: 250

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 208



Y Index: 274

Z Index: 254

6.3.2 Raw map



X Index: 211



Y Index: 226



Z Index: 250

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map

X

Y

Z

6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map

X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.017. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map

X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency

A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

6.6.1 emd_15473_msk_1.map (i)



X



Y



Z

7 Map analysis

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate



The volume at the recommended contour level is 465 nm^3; this corresponds to an approximate mass of 420 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.172 Å^{-1}

8 Fourier-Shell correlation

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC



*Reported resolution corresponds to spatial frequency of 0.172 Å^{-1}

8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	5.80	-	-
Author-provided FSC curve	5.74	7.44	5.84
Unmasked-calculated*	9.24	15.36	9.80

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 9.24 differs from the reported value 5.8 by more than 10 %

9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-15473 and PDB model 8AIX. Per-residue inclusion information can be found in section 3 on page 6.

9.1 Map-model overlay

X

Y

Z

The images above show the 3D surface view of the map at the recommended contour level 0.017 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model (i)

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.017).

9.4 Atom inclusion (i)



At the recommended contour level, 98% of all backbone atoms, 90% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.017) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8950	0.1390
A	0.8860	0.1330
B	0.9190	0.1380
C	0.8810	0.1620
D	0.8810	0.1690
E	0.8730	0.1210
F	0.9280	0.1550
G	0.9100	0.1400
H	0.9140	0.1320
I	0.8840	0.1320
J	0.8960	0.1600
K	0.7510	0.0560
L	0.8670	0.1160
M	0.9340	0.1560
N	0.9200	0.1390
O	0.8620	0.1580
P	0.9060	0.1630
Q	0.9050	0.1390
R	0.9170	0.1240
S	0.9090	0.1630
T	0.8670	0.1560
U	0.7730	0.0570
V	0.8930	0.1130
W	0.9200	0.1670
X	0.9070	0.1250