



## wwPDB EM Validation Summary Report ⓘ

Apr 1, 2025 – 10:48 pm BST

PDB ID : 7AK6 / pdb\_00007ak6  
EMDB ID : EMD-11811  
Title : Cryo-EM structure of ND6-P25L mutant respiratory complex I from Mus musculus at 3.8 Å  
Authors : Yin, Z.; Bridges, H.R.; Grba, D.; Hirst, J.  
Deposited on : 2020-09-29  
Resolution : 3.82 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev117  
Mogul : 1.8.4, CSD as541be (2020)  
MolProbity : 4.02b-467  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.42

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	115	<div> <div>52%</div> <div>79%</div> <div>19%</div> </div>
2	B	224	<div> <div>16%</div> <div>69%</div> <div>31%</div> </div>
3	C	263	<div> <div>19%</div> <div>78%</div> <div>21%</div> </div>
4	D	463	<div> <div>25%</div> <div>90%</div> <div>10%</div> </div>
5	E	248	<div> <div>55%</div> <div>85%</div> <div>15%</div> </div>
6	F	464	<div> <div>56%</div> <div>92%</div> <div>8%</div> </div>
7	G	727	<div> <div>41%</div> <div>94%</div> <div>6%</div> </div>
8	H	318	<div> <div>45%</div> <div>96%</div> <div>.</div> </div>
9	I	212	<div> <div>16%</div> <div>84%</div> <div>16%</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
10	J	172	68% 99% .
11	K	98	50% 99% .
12	L	607	61% 100%
13	M	459	36% 100%
14	N	345	32% 99%
15	O	355	50% 90% 10%
16	P	377	47% 76% 24%
17	Q	175	26% 71% 29%
18	R	116	34% 80% 19%
19	S	99	73% 83% 17%
20	T	156	44% 46% 54%
20	U	156	48% 56% 44%
21	V	116	62% 97% .
22	W	131	53% 86% 13%
23	X	172	59% 99% ..
24	Y	143	85% 98% .
25	Z	144	43% 97% .
26	a	68	46% 100%
27	b	84	60% 98% .
28	c	76	43% 61% 39%
29	d	120	49% 99% .
30	e	106	42% 97% ..
31	f	57	63% 89% 11%
32	g	151	40% 68% 32%
33	h	189	30% 72% 28%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
34	i	128	
35	j	105	
36	k	104	
37	l	186	
38	m	129	
39	n	179	
40	o	137	
41	p	176	
42	q	145	
43	r	112	
44	s	104	

## 2 Entry composition

There are 54 unique types of molecules in this entry. The entry contains 65323 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADH-ubiquinone oxidoreductase chain 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	93	Total	C	N	O	S	0	0
			761	527	105	124	5		

- Molecule 2 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 7, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	155	Total	C	N	O	S	0	0
			1241	793	222	212	14		

- Molecule 3 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 3, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	207	Total	C	N	O	S	0	0
			1721	1111	296	311	3		

- Molecule 4 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 2, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	417	Total	C	N	O	S	0	0
			3369	2155	576	614	24		

- Molecule 5 is a protein called NADH dehydrogenase [ubiquinone] flavoprotein 2, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	210	Total	C	N	O	S	0	0
			1639	1043	275	310	11		

- Molecule 6 is a protein called NADH dehydrogenase [ubiquinone] flavoprotein 1, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	428	Total	C	N	O	S	0	0
			3301	2080	590	609	22		

- Molecule 7 is a protein called NADH-ubiquinone oxidoreductase 75 kDa subunit, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	G	686	Total	C	N	O	S	0	0
			5281	3313	917	1010	41		

- Molecule 8 is a protein called NADH-ubiquinone oxidoreductase chain 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	H	305	Total	C	N	O	S	0	0
			2441	1642	367	411	21		

- Molecule 9 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 8, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	I	178	Total	C	N	O	S	0	0
			1408	885	243	268	12		

- Molecule 10 is a protein called NADH-ubiquinone oxidoreductase chain 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	J	171	Total	C	N	O	S	0	0
			1278	858	185	220	15		

- Molecule 11 is a protein called NADH-ubiquinone oxidoreductase chain 4L.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	K	98	Total	C	N	O	S	0	0
			737	477	112	137	11		

- Molecule 12 is a protein called NADH-ubiquinone oxidoreductase chain 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	L	606	Total	C	N	O	S	0	0
			4800	3182	746	827	45		

- Molecule 13 is a protein called NADH-ubiquinone oxidoreductase chain 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	M	459	Total	C	N	O	S	0	0
			3632	2408	567	617	40		

- Molecule 14 is a protein called NADH-ubiquinone oxidoreductase chain 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	N	344	Total	C	N	O	S	0	0
			2696	1791	416	452	37		

- Molecule 15 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 10, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	O	319	Total	C	N	O	S	0	0
			2599	1668	430	491	10		

- Molecule 16 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 9, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	P	288	Total	C	N	O	S	0	0
			2281	1453	412	409	7		

- Molecule 17 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 4, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	Q	125	Total	C	N	O	S	0	0
			1015	642	179	190	4		

- Molecule 18 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 6, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	R	94	Total	C	N	O	S	0	0
			738	458	135	142	3		

- Molecule 19 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	S	82	Total	C	N	O	S	0	0
			659	413	125	118	3		

- Molecule 20 is a protein called Acyl carrier protein, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	T	72	Total	C	N	O	S	0	0
			584	376	86	117	5		
20	U	88	Total	C	N	O	S	0	0
			706	453	104	144	5		

- Molecule 21 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	V	112	Total	C	N	O	S	0	0
			915	596	152	164	3		

- Molecule 22 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	W	114	Total	C	N	O	S	0	0
			970	619	180	165	6		

- Molecule 23 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 8.

Mol	Chain	Residues	Atoms					AltConf	Trace
23	X	171	Total	C	N	O	S	0	0
			1396	889	250	247	10		

- Molecule 24 is a protein called MCG5603.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	Y	140	Total	C	N	O	S	0	0
			1037	662	175	192	8		

- Molecule 25 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 13.

Mol	Chain	Residues	Atoms					AltConf	Trace
25	Z	139	Total	C	N	O	S	0	0
			1152	741	204	199	8		

- Molecule 26 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 1.



Mol	Chain	Residues	Atoms					AltConf	Trace
26	a	68	Total	C	N	O	S	0	0
			556	360	99	93	4		

- Molecule 27 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
27	b	82	Total	C	N	O	S	0	0
			643	422	104	113	4		

- Molecule 28 is a protein called NADH dehydrogenase [ubiquinone] 1 subunit C1, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
28	c	46	Total	C	N	O	S	0	0
			381	249	66	65	1		

- Molecule 29 is a protein called NADH dehydrogenase [ubiquinone] 1 subunit C2.

Mol	Chain	Residues	Atoms					AltConf	Trace
29	d	119	Total	C	N	O	S	0	0
			985	645	167	164	9		

- Molecule 30 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
30	e	104	Total	C	N	O	S	0	0
			870	550	161	151	8		

- Molecule 31 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
31	f	51	Total	C	N	O	S	0	0
			433	278	78	75	2		

- Molecule 32 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 11, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
32	g	102	Total	C	N	O	S	0	0
			858	555	137	162	4		

- Molecule 33 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 5, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
33	h	137	Total	C	N	O	S	0	0
			1153	756	192	202	3		

- Molecule 34 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
34	i	95	Total	C	N	O	S	0	0
			802	524	138	137	3		

- Molecule 35 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 2, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
35	j	62	Total	C	N	O	S	0	0
			537	355	88	93	1		

- Molecule 36 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
36	k	77	Total	C	N	O	S	0	0
			626	414	106	104	2		

- Molecule 37 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 8, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
37	l	156	Total	C	N	O	S	0	0
			1312	846	219	236	11		

- Molecule 38 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
38	m	125	Total	C	N	O		0	0
			1044	673	188	183			

- Molecule 39 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 9.

Mol	Chain	Residues	Atoms					AltConf	Trace
39	n	176	Total	C	N	O	S	0	0
			1527	976	274	266	11		

- Molecule 40 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 7.

Mol	Chain	Residues	Atoms					AltConf	Trace
40	o	114	Total	C	N	O	S	0	0
			984	620	185	171	8		

- Molecule 41 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 10.

Mol	Chain	Residues	Atoms					AltConf	Trace
41	p	168	Total	C	N	O	S	0	0
			1424	896	256	264	8		

- Molecule 42 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 12.

Mol	Chain	Residues	Atoms					AltConf	Trace
42	q	143	Total	C	N	O	S	0	0
			1192	766	212	210	4		

- Molecule 43 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 7.

Mol	Chain	Residues	Atoms					AltConf	Trace
43	r	96	Total	C	N	O	S	0	0
			776	490	145	138	3		

- Molecule 44 is a protein called NADH dehydrogenase [ubiquinone] flavoprotein 3, mitochondrial.


Mol	Chain	Residues	Atoms				AltConf	Trace
44	s	39	Total	C	N	O	0	0
			329	207	59	63		

- Molecule 45 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe<sub>4</sub>S<sub>4</sub>).




Mol	Chain	Residues	Atoms			AltConf
45	B	1	Total	Fe	S	0
			8	4	4	
45	F	1	Total	Fe	S	0
			8	4	4	
45	G	1	Total	Fe	S	0
			8	4	4	
45	G	1	Total	Fe	S	0
			8	4	4	
45	I	1	Total	Fe	S	0
			8	4	4	
45	I	1	Total	Fe	S	0
			8	4	4	

- Molecule 46 is 1,2-DIACYL-SN-GLYCERO-3-PHOSPHOCHOLINE (CCD ID: PC1) (formula: C<sub>44</sub>H<sub>88</sub>NO<sub>8</sub>P).




Mol	Chain	Residues	Atoms					AltConf
			Total	C	N	O	P	
46	B	1	26	16	1	8	1	0

- Molecule 47 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula:  $\text{Fe}_2\text{S}_2$ ).




Mol	Chain	Residues	Atoms			AltConf
			Total	Fe	S	
47	E	1	4	2	2	0
47	G	1	4	2	2	0

- Molecule 48 is FLAVIN MONONUCLEOTIDE (CCD ID: FMN) (formula:  $\text{C}_{17}\text{H}_{21}\text{N}_4\text{O}_9\text{P}$ ).



Mol	Chain	Residues	Atoms					AltConf
48	F	1	Total 31	C 17	N 4	O 9	P 1	0

- Molecule 49 is 1,2-Distearoyl-sn-glycerophosphoethanolamine (CCD ID: 3PE) (formula:  $C_{41}H_{82}NO_8P$ ).




Mol	Chain	Residues	Atoms					AltConf
49	H	1	Total 42	C 32	N 1	O 8	P 1	0
49	J	1	Total 38	C 28	N 1	O 8	P 1	0

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms					AltConf
49	L	1	Total	C	N	O	P	0
			45	35	1	8	1	
49	M	1	Total	C	N	O	P	0
			38	28	1	8	1	

- Molecule 50 is CARDIOLIPIN (CCD ID: CDL) (formula:  $C_{81}H_{156}O_{17}P_2$ ).



Mol	Chain	Residues	Atoms				AltConf
50	L	1	Total	C	O	P	0
			54	35	17	2	
50	d	1	Total	C	O	P	0
			50	31	17	2	


- Molecule 51 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula:  $C_{10}H_{16}N_5O_{13}P_3$ ).






Mol	Chain	Residues	Atoms		AltConf
53	R	1	Total	Zn	0
			1	1	


- Molecule 54 is {S}-[2-[3-[[2 {R}]-3,3-dimethyl-2-oxidanyl-4-phosphonooxy-butanoyl]amino]propanoylamino]ethyl] (3 {S})-3-oxidanyltetradecanethioate (CCD ID: EHZ) (formula: C<sub>25</sub>H<sub>49</sub>N<sub>2</sub>O<sub>9</sub>PS).





Mol	Chain	Residues	Atoms						AltConf
54	T	1	Total	C	N	O	P	S	0
			37	25	2	8	1	1	
54	U	1	Total	C	N	O	P	S	0
			37	25	2	8	1	1	









• Molecule 7: NADH-ubiquinone oxidoreductase 75 kDa subunit, mitochondrial




- Molecule 8: NADH-ubiquinone oxidoreductase chain 1



- Molecule 9: NADH dehydrogenase [ubiquinone] iron-sulfur protein 8, mitochondrial




- Molecule 10: NADH-ubiquinone oxidoreductase chain 6




- Molecule 11: NADH-ubiquinone oxidoreductase chain 4L







• Molecule 12: NADH-ubiquinone oxidoreductase chain 5




• Molecule 13: NADH-ubiquinone oxidoreductase chain 4








• Molecule 14: NADH-ubiquinone oxidoreductase chain 2





• Molecule 15: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 10, mitochondrial





















- Molecule 22: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 6




- Molecule 23: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 8




- Molecule 24: MCG5603




- Molecule 25: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 13








- Molecule 26: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 1




- Molecule 27: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 3





- Molecule 28: NADH dehydrogenase [ubiquinone] 1 subunit C1, mitochondrial





- Molecule 29: NADH dehydrogenase [ubiquinone] 1 subunit C2





- Molecule 30: NADH dehydrogenase [ubiquinone] iron-sulfur protein 5





- Molecule 31: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 1




- Molecule 32: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 11, mitochondrial




- Molecule 33: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 5, mitochondrial




- Molecule 34: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 6




- Molecule 35: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 2, mitochondrial




- Molecule 36: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 3




- Molecule 37: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 8, mitochondrial





- Molecule 38: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 4





- Molecule 39: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 9




- Molecule 40: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 7




- Molecule 41: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 10








- Molecule 42: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 12



- Molecule 43: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 7



- Molecule 44: NADH dehydrogenase [ubiquinone] flavoprotein 3, mitochondrial



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	25629	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	50.015	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	130000	Depositor
Image detector	GATAN K2 QUANTUM (4k x 4k)	Depositor
Maximum map value	0.182	Depositor
Minimum map value	-0.046	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.0268	Depositor
Map size (Å)	474.30002, 474.30002, 474.30002	wwPDB
Map dimensions	450, 450, 450	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.054, 1.054, 1.054	Depositor



## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, SF4, ATP, EHZ, CDL, PC1, 2MR, FES, ZN, FME, AYA, 3PE, NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.31	0/771	0.52	2/1054 (0.2%)
2	B	0.43	0/1272	0.48	0/1722
3	C	0.38	0/1771	0.47	0/2412
4	D	0.37	0/3440	0.45	0/4656
5	E	0.31	0/1679	0.45	0/2288
6	F	0.31	0/3376	0.46	0/4561
7	G	0.34	0/5368	0.47	0/7273
8	H	0.33	0/2504	0.43	0/3420
9	I	0.41	0/1438	0.48	0/1946
10	J	0.32	0/1297	0.44	0/1764
11	K	0.31	0/738	0.43	0/1002
12	L	0.31	0/4913	0.45	0/6686
13	M	0.31	0/3709	0.46	0/5052
14	N	0.33	0/2748	0.44	0/3741
15	O	0.32	0/2666	0.44	0/3615
16	P	0.32	0/2329	0.47	0/3145
17	Q	0.32	0/1038	0.47	0/1401
18	R	0.34	0/751	0.46	0/1011
19	S	0.27	0/670	0.47	0/904
20	T	0.28	0/593	0.44	0/799
20	U	0.29	0/718	0.41	0/970
21	V	0.31	0/937	0.42	0/1270
22	W	0.31	0/993	0.45	0/1335
23	X	0.30	0/1434	0.44	0/1937
24	Y	0.29	0/1061	0.45	0/1439
25	Z	0.32	0/1183	0.44	0/1597
26	a	0.34	0/569	0.43	0/766
27	b	0.29	0/666	0.41	0/914
28	c	0.30	0/392	0.40	0/533
29	d	0.34	0/1017	0.44	0/1373
30	e	0.30	0/892	0.45	0/1187
31	f	0.28	0/443	0.44	0/596

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
32	g	0.33	0/887	0.42	0/1209
33	h	0.33	0/1188	0.42	0/1610
34	i	0.30	0/828	0.45	0/1125
35	j	0.27	0/561	0.41	0/768
36	k	0.28	0/646	0.43	0/873
37	l	0.31	0/1367	0.41	0/1866
38	m	0.31	0/1073	0.45	0/1455
39	n	0.30	0/1581	0.41	0/2140
40	o	0.27	0/1009	0.42	0/1355
41	p	0.33	0/1457	0.42	0/1969
42	q	0.35	0/1234	0.45	0/1681
43	r	0.31	0/786	0.43	0/1062
44	s	0.27	0/338	0.45	0/458
All	All	0.33	0/66331	0.45	2/89940 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	49	LEU	CB-CG-CD2	5.81	120.88	111.00
1	A	49	LEU	CB-CG-CD1	5.60	120.51	111.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	89/115 (77%)	83 (93%)	6 (7%)	0	100	100
2	B	153/224 (68%)	136 (89%)	17 (11%)	0	100	100
3	C	205/263 (78%)	181 (88%)	24 (12%)	0	100	100
4	D	410/463 (89%)	379 (92%)	30 (7%)	1 (0%)	44	74
5	E	208/248 (84%)	180 (86%)	28 (14%)	0	100	100
6	F	426/464 (92%)	385 (90%)	41 (10%)	0	100	100
7	G	684/727 (94%)	617 (90%)	67 (10%)	0	100	100
8	H	299/318 (94%)	267 (89%)	32 (11%)	0	100	100
9	I	176/212 (83%)	157 (89%)	19 (11%)	0	100	100
10	J	169/172 (98%)	154 (91%)	15 (9%)	0	100	100
11	K	96/98 (98%)	90 (94%)	6 (6%)	0	100	100
12	L	604/607 (100%)	544 (90%)	60 (10%)	0	100	100
13	M	457/459 (100%)	411 (90%)	46 (10%)	0	100	100
14	N	342/345 (99%)	316 (92%)	26 (8%)	0	100	100
15	O	317/355 (89%)	283 (89%)	34 (11%)	0	100	100
16	P	280/377 (74%)	244 (87%)	36 (13%)	0	100	100
17	Q	123/175 (70%)	111 (90%)	12 (10%)	0	100	100
18	R	92/116 (79%)	84 (91%)	8 (9%)	0	100	100
19	S	80/99 (81%)	76 (95%)	4 (5%)	0	100	100
20	T	70/156 (45%)	62 (89%)	8 (11%)	0	100	100
20	U	86/156 (55%)	75 (87%)	11 (13%)	0	100	100
21	V	110/116 (95%)	99 (90%)	11 (10%)	0	100	100
22	W	112/131 (86%)	103 (92%)	9 (8%)	0	100	100
23	X	169/172 (98%)	153 (90%)	15 (9%)	1 (1%)	22	57
24	Y	138/143 (96%)	129 (94%)	9 (6%)	0	100	100
25	Z	137/144 (95%)	126 (92%)	11 (8%)	0	100	100
26	a	66/68 (97%)	62 (94%)	4 (6%)	0	100	100
27	b	80/84 (95%)	69 (86%)	11 (14%)	0	100	100
28	c	44/76 (58%)	41 (93%)	3 (7%)	0	100	100
29	d	117/120 (98%)	109 (93%)	8 (7%)	0	100	100
30	e	102/106 (96%)	87 (85%)	15 (15%)	0	100	100
31	f	49/57 (86%)	41 (84%)	8 (16%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
32	g	100/151 (66%)	88 (88%)	12 (12%)	0	100	100
33	h	135/189 (71%)	121 (90%)	14 (10%)	0	100	100
34	i	91/128 (71%)	71 (78%)	20 (22%)	0	100	100
35	j	60/105 (57%)	55 (92%)	5 (8%)	0	100	100
36	k	75/104 (72%)	69 (92%)	6 (8%)	0	100	100
37	l	154/186 (83%)	132 (86%)	22 (14%)	0	100	100
38	m	123/129 (95%)	114 (93%)	9 (7%)	0	100	100
39	n	174/179 (97%)	162 (93%)	12 (7%)	0	100	100
40	o	112/137 (82%)	95 (85%)	17 (15%)	0	100	100
41	p	166/176 (94%)	151 (91%)	15 (9%)	0	100	100
42	q	141/145 (97%)	122 (86%)	19 (14%)	0	100	100
43	r	92/112 (82%)	80 (87%)	12 (13%)	0	100	100
44	s	37/104 (36%)	34 (92%)	3 (8%)	0	100	100
All	All	7950/9211 (86%)	7148 (90%)	800 (10%)	2 (0%)	100	100

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
23	X	168	PHE
4	D	86	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	84/103 (82%)	84 (100%)	0	100	100
2	B	131/185 (71%)	131 (100%)	0	100	100
3	C	189/227 (83%)	188 (100%)	1 (0%)	86	90
4	D	361/394 (92%)	361 (100%)	0	100	100
5	E	183/206 (89%)	183 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
6	F	343/370 (93%)	343 (100%)	0	100	100
7	G	578/610 (95%)	575 (100%)	3 (0%)	86	90
8	H	267/279 (96%)	267 (100%)	0	100	100
9	I	145/178 (82%)	145 (100%)	0	100	100
10	J	131/137 (96%)	131 (100%)	0	100	100
11	K	87/87 (100%)	87 (100%)	0	100	100
12	L	548/549 (100%)	548 (100%)	0	100	100
13	M	414/414 (100%)	413 (100%)	1 (0%)	92	94
14	N	306/307 (100%)	306 (100%)	0	100	100
15	O	283/309 (92%)	283 (100%)	0	100	100
16	P	249/325 (77%)	247 (99%)	2 (1%)	79	85
17	Q	112/153 (73%)	112 (100%)	0	100	100
18	R	79/96 (82%)	78 (99%)	1 (1%)	65	76
19	S	73/80 (91%)	73 (100%)	0	100	100
20	T	67/135 (50%)	66 (98%)	1 (2%)	60	75
20	U	81/135 (60%)	81 (100%)	0	100	100
21	V	100/102 (98%)	100 (100%)	0	100	100
22	W	108/114 (95%)	107 (99%)	1 (1%)	75	83
23	X	153/154 (99%)	153 (100%)	0	100	100
24	Y	105/107 (98%)	105 (100%)	0	100	100
25	Z	120/123 (98%)	120 (100%)	0	100	100
26	a	58/58 (100%)	58 (100%)	0	100	100
27	b	72/73 (99%)	72 (100%)	0	100	100
28	c	40/67 (60%)	40 (100%)	0	100	100
29	d	106/107 (99%)	106 (100%)	0	100	100
30	e	92/94 (98%)	91 (99%)	1 (1%)	70	79
31	f	47/53 (89%)	47 (100%)	0	100	100
32	g	93/129 (72%)	93 (100%)	0	100	100
33	h	122/162 (75%)	122 (100%)	0	100	100
34	i	90/120 (75%)	89 (99%)	1 (1%)	70	79
35	j	58/87 (67%)	58 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
36	k	60/78 (77%)	59 (98%)	1 (2%)	56	72
37	l	141/161 (88%)	141 (100%)	0	100	100
38	m	111/114 (97%)	111 (100%)	0	100	100
39	n	161/164 (98%)	160 (99%)	1 (1%)	84	88
40	o	106/121 (88%)	106 (100%)	0	100	100
41	p	153/158 (97%)	153 (100%)	0	100	100
42	q	129/131 (98%)	128 (99%)	1 (1%)	79	85
43	r	86/95 (90%)	86 (100%)	0	100	100
44	s	38/95 (40%)	38 (100%)	0	100	100
All	All	7060/7946 (89%)	7045 (100%)	15 (0%)	91	94

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
18	R	80	LYS
39	n	174	ARG
20	T	12	LYS
42	q	60	ARG
34	i	120	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 89 such sidechains are listed below:

Mol	Chain	Res	Type
23	X	34	GLN
33	h	44	ASN
23	X	102	GLN
25	Z	89	ASN
37	l	71	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

9 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
8	FME	H	1	8	8,9,10	0.93	0	7,9,11	1.13	1 (14%)
13	FME	M	1	13	8,9,10	0.95	0	7,9,11	0.67	0
43	AYA	r	1	43	6,7,8	1.29	1 (16%)	5,8,10	1.26	1 (20%)
1	FME	A	1	1	8,9,10	0.89	0	7,9,11	1.46	1 (14%)
12	FME	L	1	12	8,9,10	0.95	0	7,9,11	0.93	0
10	FME	J	1	10	8,9,10	0.92	0	7,9,11	0.85	0
4	2MR	D	85	4	10,12,13	2.46	2 (20%)	5,13,15	1.72	2 (40%)
11	FME	K	1	11	8,9,10	0.94	0	7,9,11	1.35	1 (14%)
14	FME	N	1	14	8,9,10	0.89	0	7,9,11	1.22	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	FME	H	1	8	-	2/7/9/11	-
13	FME	M	1	13	-	1/7/9/11	-
43	AYA	r	1	43	-	1/4/6/8	-
1	FME	A	1	1	-	4/7/9/11	-
12	FME	L	1	12	-	3/7/9/11	-
10	FME	J	1	10	-	1/7/9/11	-
4	2MR	D	85	4	-	3/10/13/15	-
11	FME	K	1	11	-	3/7/9/11	-
14	FME	N	1	14	-	4/7/9/11	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	85	2MR	CZ-NE	5.31	1.45	1.34
4	D	85	2MR	CZ-NH2	4.94	1.44	1.33
43	r	1	AYA	CA-N	-2.54	1.43	1.46

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1	FME	C-CA-N	3.12	115.36	109.73
11	K	1	FME	C-CA-N	2.90	114.97	109.73
4	D	85	2MR	CD-NE-CZ	2.87	128.79	123.41
43	r	1	AYA	CB-CA-N	2.50	112.39	109.61
4	D	85	2MR	NE-CZ-NH2	-2.46	117.23	119.48

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	1	FME	O-C-CA-CB
8	H	1	FME	CB-CA-N-CN
8	H	1	FME	CA-CB-CG-SD
11	K	1	FME	N-CA-CB-CG
12	L	1	FME	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 1 is monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
47	FES	G	803	7	0,4,4	-	-	-		
45	SF4	B	201	2	0,12,12	-	-	-		
47	FES	E	301	5	0,4,4	-	-	-		



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
50	CDL	d	201	-	49,49,99	1.22	7 (14%)	55,61,111	1.35	5 (9%)
54	EHZ	U	201	20	29,36,37	1.72	5 (17%)	35,44,47	1.55	6 (17%)
50	CDL	L	702	-	53,53,99	1.18	8 (15%)	59,65,111	1.13	4 (6%)
45	SF4	I	201	9	0,12,12	-	-	-	-	-
45	SF4	I	202	9	0,12,12	-	-	-	-	-
49	3PE	H	401	-	41,41,50	0.95	4 (9%)	44,46,55	1.26	4 (9%)
45	SF4	G	801	7	0,12,12	-	-	-	-	-
45	SF4	G	802	7	0,12,12	-	-	-	-	-
51	ATP	O	401	-	26,33,33	0.92	1 (3%)	31,52,52	1.73	5 (16%)
52	NDP	P	501	-	45,52,52	2.23	5 (11%)	53,80,80	1.76	8 (15%)
46	PC1	B	202	-	25,25,53	1.36	4 (16%)	31,33,61	1.10	2 (6%)
49	3PE	J	401	-	37,37,50	0.99	4 (10%)	40,42,55	1.11	2 (5%)
45	SF4	F	502	6	0,12,12	-	-	-	-	-
49	3PE	L	701	-	44,44,50	0.90	4 (9%)	47,49,55	1.12	2 (4%)
48	FMN	F	501	-	33,33,33	1.09	2 (6%)	48,50,50	1.25	9 (18%)
49	3PE	M	501	-	37,37,50	0.99	4 (10%)	40,42,55	1.10	2 (5%)
54	EHZ	T	201	20	29,36,37	1.69	5 (17%)	35,44,47	1.53	4 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
47	FES	G	803	7	-	-	0/1/1/1
45	SF4	B	201	2	-	-	0/6/5/5
47	FES	E	301	5	-	-	0/1/1/1
50	CDL	d	201	-	-	22/59/59/110	-
54	EHZ	U	201	20	-	24/42/44/45	-
50	CDL	L	702	-	-	25/64/64/110	-
45	SF4	I	201	9	-	-	0/6/5/5
45	SF4	I	202	9	-	-	0/6/5/5
49	3PE	H	401	-	-	13/45/45/54	-
45	SF4	G	801	7	-	-	0/6/5/5
45	SF4	G	802	7	-	-	0/6/5/5
51	ATP	O	401	-	-	5/18/38/38	0/3/3/3
52	NDP	P	501	-	-	9/30/77/77	0/5/5/5
46	PC1	B	202	-	-	15/29/29/57	-

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
49	3PE	J	401	-	-	18/41/41/54	-
45	SF4	F	502	6	-	-	0/6/5/5
49	3PE	L	701	-	-	25/48/48/54	-
48	FMN	F	501	-	-	10/18/18/18	0/3/3/3
49	3PE	M	501	-	-	23/41/41/54	-
54	EHZ	T	201	20	-	15/42/44/45	-

The worst 5 of 53 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
52	P	501	NDP	P2B-O2B	12.32	1.82	1.59
54	U	201	EHZ	C15-N2	5.60	1.45	1.33
54	T	201	EHZ	C15-N2	5.51	1.45	1.33
54	U	201	EHZ	C12-N1	5.38	1.45	1.33
54	T	201	EHZ	C12-N1	5.28	1.45	1.33

The worst 5 of 53 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
52	P	501	NDP	PN-O3-PA	-7.51	107.07	132.83
54	T	201	EHZ	C8-C9-S1	5.46	120.39	113.63
54	U	201	EHZ	C8-C9-S1	5.36	120.26	113.63
50	d	201	CDL	OA6-CA5-C11	4.75	119.83	111.09
49	L	701	3PE	O21-C21-C22	4.64	121.51	111.50

There are no chirality outliers.

5 of 204 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
46	B	202	PC1	C1-O11-P-O12
46	B	202	PC1	C1-O11-P-O14
46	B	202	PC1	O13-C11-C12-N
48	F	501	FMN	N10-C1'-C2'-O2'
48	F	501	FMN	N10-C1'-C2'-C3'



There are no ring outliers.


No monomer is involved in short contacts.



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will

also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.












## Ligand PC1 B 202





## Ligand 3PE J 401











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-11811. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 225




Y Index: 225




Z Index: 225


### 6.2.2 Raw map



X Index: 225



Y Index: 225




Z Index: 225


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 225




Y Index: 216




Z Index: 279


### 6.3.2 Raw map



X Index: 225



Y Index: 216




Z Index: 245

The images above show the largest variance slices of the map in three orthogonal directions.




## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0268. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.




## 6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


### 6.6.1 emd\_11811\_msk\_1.map [i](#)



X



Y




Z

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 146 nm<sup>3</sup>; this corresponds to an approximate mass of 132 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of 0.262 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of  $0.262 \text{ \AA}^{-1}$

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.82	-	-
Author-provided FSC curve	3.80	4.37	3.91
Unmasked-calculated*	4.66	8.31	6.30

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.66 differs from the reported value 3.82 by more than 10 %

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-11811 and PDB model 7AK6. Per-residue inclusion information can be found in [section 3](#) on [page 18](#).

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.0268 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0268).









































































## 9.4 Atom inclusion ⓘ



At the recommended contour level, 50% of all backbone atoms, 38% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary ⓘ























The table lists the average atom inclusion at the recommended contour level (0.0268) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.3810	 0.4470
A	 0.3090	 0.4300
B	 0.5680	 0.4850
C	 0.5340	 0.4930
D	 0.5190	 0.4770
E	 0.3040	 0.4230
F	 0.3490	 0.4430
G	 0.4310	 0.4570
H	 0.4020	 0.4510
I	 0.5890	 0.4930
J	 0.2760	 0.4190
K	 0.3850	 0.4600
L	 0.3490	 0.4400
M	 0.4610	 0.4670
N	 0.4800	 0.4640
O	 0.3610	 0.4480
P	 0.3410	 0.4350
Q	 0.4420	 0.4760
R	 0.4400	 0.4750
S	 0.1920	 0.3990
T	 0.0760	 0.3060
U	 0.1930	 0.3950
V	 0.3270	 0.4380
W	 0.3420	 0.4330
X	 0.3480	 0.4540
Y	 0.2180	 0.4140
Z	 0.4400	 0.4540
a	 0.4260	 0.4560
b	 0.3340	 0.4240
c	 0.2810	 0.4230
d	 0.3780	 0.4630
e	 0.3950	 0.4520
f	 0.2800	 0.4130
g	 0.3510	 0.4440
h	 0.4250	 0.4620



*Continued on next page...*

*Continued from previous page...*

Chain	Atom inclusion	Q-score
i	 0.2660	 0.4070
j	 0.1650	 0.3950
k	 0.1680	 0.4170
l	 0.3220	 0.4450
m	 0.2950	 0.4260
n	 0.3130	 0.4290
o	 0.2060	 0.3850
p	 0.3540	 0.4300
q	 0.4260	 0.4660
r	 0.3950	 0.4560
s	 0.2250	 0.4390