



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 12, 2024 – 02:56 PM EDT

PDB ID : 1ARL  
Title : CARBOXYPEPTIDASE A WITH ZN REMOVED  
Authors : Greenblatt, H.M.; Feinberg, H.; Tucker, P.A.; Shoham, G.  
Deposited on : 1994-11-22  
Resolution : 1.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.36.2
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	10282 (1.90-1.86)
Ramachandran outliers	138981	10152 (1.90-1.86)
Sidechain outliers	138945	10152 (1.90-1.86)
RSRZ outliers	127900	9303 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	307	<div> <div></div> <div>59%</div> <div>33%</div> <div>7%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2616 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called APO-CARBOXYPEPTIDASE A=ALPHA= (COX).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	307	Total	C	N	O	S	0	0	0
			2437	1561	403	468	5			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	179	Total	O	0	0
			179	179		

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: APO-CARBOXYPEPTIDASE A=ALPHA= (COX)



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.70Å 60.32Å 47.20Å 90.00° 97.39° 90.00°	Depositor
Resolution (Å)	13.00 – 1.88 13.01 – 1.88	Depositor EDS
% Data completeness (in resolution range)	88.0 (13.00-1.88) 87.9 (13.01-1.88)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	14.93 (at 1.88Å)	Xtriage
Refinement program	TNT, PROLSQ	Depositor
R, $R_{free}$	0.143 , (Not available) 0.137 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.2	Xtriage
Anisotropy	0.120	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 83.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2616	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.06	9/2503 (0.4%)	1.62	42/3402 (1.2%)

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	254	SER	CB-OG	-7.24	1.32	1.42
1	A	43	GLU	CD-OE1	7.01	1.33	1.25
1	A	108	GLU	CD-OE2	6.96	1.33	1.25
1	A	88	GLU	CD-OE1	6.34	1.32	1.25
1	A	218	GLU	CD-OE2	6.01	1.32	1.25
1	A	108	GLU	CD-OE1	-5.74	1.19	1.25
1	A	270	GLU	CD-OE2	5.70	1.31	1.25
1	A	292	GLU	CD-OE1	5.55	1.31	1.25
1	A	72	GLU	CD-OE2	-5.09	1.20	1.25

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	71	ARG	NE-CZ-NH1	14.58	127.59	120.30
1	A	130	ARG	NE-CZ-NH2	-14.26	113.17	120.30
1	A	130	ARG	NE-CZ-NH1	12.74	126.67	120.30
1	A	71	ARG	NE-CZ-NH2	-11.10	114.75	120.30
1	A	20	ASP	CB-CG-OD1	10.15	127.44	118.30
1	A	142	ASP	CB-CG-OD2	-10.08	109.23	118.30
1	A	65	ASP	CB-CG-OD2	-9.53	109.72	118.30
1	A	65	ASP	CB-CG-OD1	8.56	126.00	118.30
1	A	20	ASP	CB-CG-OD2	-7.31	111.72	118.30
1	A	48	TYR	CB-CG-CD2	-7.21	116.67	121.00
1	A	101	ASP	CB-CG-OD2	-7.20	111.82	118.30
1	A	142	ASP	CB-CG-OD1	7.09	124.68	118.30
1	A	185	ASP	CB-CG-OD1	6.82	124.44	118.30
1	A	256	ASP	CB-CG-OD1	6.67	124.31	118.30
1	A	240	TYR	CB-CG-CD1	-6.67	117.00	121.00

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	104	ASP	CB-CG-OD1	6.64	124.27	118.30
1	A	148	ASP	CB-CG-OD1	6.56	124.21	118.30
1	A	135	SER	N-CA-CB	6.56	120.33	110.50
1	A	23	ASP	CB-CG-OD2	6.46	124.11	118.30
1	A	148	ASP	CB-CG-OD2	-6.44	112.50	118.30
1	A	185	ASP	CB-CG-OD2	-6.40	112.54	118.30
1	A	23	ASP	CB-CG-OD1	-6.33	112.60	118.30
1	A	101	ASP	CB-CG-OD1	6.19	123.87	118.30
1	A	89	ASP	CB-CG-OD2	-6.17	112.75	118.30
1	A	104	ASP	CB-CG-OD2	-6.16	112.75	118.30
1	A	93	ASP	CB-CG-OD2	-6.14	112.77	118.30
1	A	281	LEU	CB-CG-CD2	6.08	121.34	111.00
1	A	114	ASP	CB-CG-OD2	6.05	123.75	118.30
1	A	45	ARG	NE-CZ-NH2	-5.94	117.33	120.30
1	A	145	ARG	NE-CZ-NH2	5.84	123.22	120.30
1	A	216	LYS	CA-CB-CG	-5.56	101.17	113.40
1	A	80	VAL	CA-CB-CG2	-5.51	102.64	110.90
1	A	90	TYR	CB-CG-CD2	-5.42	117.75	121.00
1	A	93	ASP	CB-CG-OD1	5.39	123.15	118.30
1	A	194	SER	N-CA-CB	5.32	118.48	110.50
1	A	221	GLN	CB-CG-CD	5.31	125.41	111.60
1	A	100	LEU	N-CA-CB	5.27	120.94	110.40
1	A	215	ASP	CB-CG-OD2	5.18	122.96	118.30
1	A	11	THR	CA-CB-OG1	-5.13	98.23	109.00
1	A	268	THR	CA-CB-CG2	-5.09	105.27	112.40
1	A	254	SER	CB-CA-C	-5.07	100.46	110.10
1	A	139	VAL	CA-CB-CG2	5.05	118.48	110.90

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2437	0	2346	74	0
2	A	179	0	0	10	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	2616	0	2346	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:137:LEU:HD22	1:A:137:LEU:H	1.29	0.98
1:A:186:HIS:HD2	1:A:188:ASN:H	1.14	0.93
1:A:293:THR:HB	2:A:477:HOH:O	1.70	0.91
1:A:71:ARG:HD2	1:A:127:ARG:HB3	1.55	0.89
1:A:242:SER:O	1:A:246:THR:HG23	1.76	0.85
1:A:289:THR:O	1:A:293:THR:HG22	1.82	0.79
1:A:91:GLY:H	1:A:97:THR:CG2	2.00	0.75
1:A:186:HIS:HD2	1:A:188:ASN:N	1.87	0.73
1:A:186:HIS:CD2	1:A:188:ASN:H	2.04	0.73
1:A:175:GLU:O	1:A:179:ILE:HD13	1.95	0.67
1:A:31:GLN:HB2	2:A:493:HOH:O	1.94	0.66
1:A:137:LEU:HD22	1:A:137:LEU:N	2.08	0.66
1:A:71:ARG:HD3	2:A:421:HOH:O	1.97	0.65
1:A:295:LEU:HD22	2:A:580:HOH:O	1.96	0.63
1:A:247:ILE:HG22	1:A:248:TYR:N	2.14	0.63
1:A:91:GLY:H	1:A:97:THR:HG21	1.66	0.60
1:A:93:ASP:O	1:A:97:THR:HG23	2.01	0.59
1:A:62:ILE:HD12	1:A:191:ALA:HB3	1.85	0.59
1:A:204:TYR:HB2	1:A:205:PRO:CD	2.33	0.58
1:A:137:LEU:H	1:A:137:LEU:CD2	2.08	0.58
1:A:29:HIS:N	1:A:30:PRO:HD3	2.19	0.57
1:A:32:LEU:HD11	1:A:100:LEU:HD13	1.85	0.56
1:A:132:VAL:O	1:A:133:THR:HG23	2.05	0.56
1:A:168:LYS:HG2	2:A:546:HOH:O	2.06	0.55
1:A:257:TRP:O	1:A:261:GLN:HG2	2.08	0.53
1:A:71:ARG:HD2	1:A:127:ARG:CB	2.33	0.53
1:A:42:TYR:OH	1:A:132:VAL:HG22	2.09	0.53
1:A:216:LYS:NZ	2:A:494:HOH:O	2.40	0.53
1:A:132:VAL:HG23	1:A:133:THR:N	2.25	0.52
1:A:168:LYS:HD2	1:A:169:TYR:CE1	2.44	0.52
1:A:272:ARG:HH11	1:A:285:GLN:HE21	1.58	0.51
1:A:204:TYR:HB2	1:A:205:PRO:HD2	1.94	0.49
1:A:287:ILE:N	1:A:288:PRO:HD2	2.28	0.49

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:154:ALA:O	1:A:249:GLN:OE1	2.30	0.49
1:A:281:LEU:HD13	1:A:285:GLN:HB2	1.95	0.49
1:A:24:LEU:HD23	2:A:553:HOH:O	2.13	0.48
1:A:91:GLY:N	1:A:97:THR:HG21	2.29	0.48
1:A:112:ASN:ND2	1:A:115:GLY:H	2.11	0.47
1:A:233:LEU:HD13	1:A:234:TYR:CE2	2.50	0.47
1:A:196:HIS:ND1	1:A:268:THR:HG21	2.30	0.47
1:A:216:LYS:HG2	1:A:216:LYS:O	2.12	0.47
1:A:179:ILE:N	1:A:179:ILE:CD1	2.77	0.47
1:A:248:TYR:CD2	1:A:249:GLN:N	2.83	0.47
1:A:86:PHE:HE1	1:A:294:TRP:CZ3	2.34	0.46
1:A:239:LYS:HE3	2:A:468:HOH:O	2.15	0.45
1:A:248:TYR:CG	1:A:249:GLN:N	2.84	0.45
1:A:272:ARG:HH11	1:A:285:GLN:NE2	2.15	0.45
1:A:287:ILE:N	1:A:288:PRO:CD	2.80	0.45
1:A:85:LYS:HE3	1:A:89:ASP:OD2	2.18	0.44
1:A:179:ILE:HD11	2:A:403:HOH:O	2.16	0.44
1:A:268:THR:HG22	1:A:269:PHE:N	2.33	0.44
1:A:137:LEU:N	1:A:137:LEU:HD13	2.33	0.43
1:A:192:PHE:O	1:A:193:LEU:HD12	2.19	0.43
1:A:177:LYS:HA	1:A:177:LYS:HD3	1.81	0.43
1:A:156:ALA:HB1	1:A:166:HIS:HB3	2.01	0.43
1:A:71:ARG:CD	1:A:127:ARG:HB3	2.36	0.42
1:A:160:PRO:HA	1:A:165:TYR:CG	2.54	0.42
1:A:5:ASN:HD21	1:A:84:LYS:HZ1	1.66	0.42
1:A:55:GLY:O	1:A:59:ARG:HD2	2.20	0.42
1:A:40:ARG:HH11	1:A:40:ARG:HD3	1.75	0.41
1:A:196:HIS:CE1	1:A:268:THR:HG21	2.55	0.41
1:A:247:ILE:CG2	1:A:248:TYR:N	2.76	0.41
1:A:163:GLU:HG3	2:A:568:HOH:O	2.21	0.41
1:A:28:GLU:C	1:A:30:PRO:HD3	2.41	0.41
1:A:60:PRO:HB2	1:A:103:MET:HE3	2.03	0.41
1:A:50:LEU:HD23	1:A:50:LEU:HA	1.89	0.41
1:A:192:PHE:C	1:A:193:LEU:HD12	2.41	0.41
1:A:219:LEU:HG	1:A:267:PHE:CZ	2.55	0.41
1:A:255:ILE:HB	1:A:266:SER:HB3	2.03	0.41
1:A:11:THR:OG1	1:A:12:TYR:N	2.54	0.40
1:A:119:THR:HA	1:A:123:ASN:O	2.21	0.40
1:A:32:LEU:O	1:A:32:LEU:HD12	2.21	0.40
1:A:133:THR:HB	1:A:134:SER:H	1.28	0.40
1:A:3:SER:HB3	1:A:6:THR:HB	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	305/307 (99%)	293 (96%)	9 (3%)	3 (1%)	15 6

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	199	SER
1	A	135	SER
1	A	133	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	263/263 (100%)	235 (89%)	28 (11%)	6 2

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	THR
1	A	33	VAL
1	A	34	SER
1	A	57	SER
1	A	58	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	71	ARG
1	A	97	THR
1	A	112	ASN
1	A	137	LEU
1	A	139	VAL
1	A	166	HIS
1	A	168	LYS
1	A	179	ILE
1	A	184	LYS
1	A	202	LEU
1	A	203	LEU
1	A	212	SER
1	A	216	LYS
1	A	233	LEU
1	A	239	LYS
1	A	246	THR
1	A	254	SER
1	A	271	LEU
1	A	276	ARG
1	A	281	LEU
1	A	295	LEU
1	A	298	LEU
1	A	305	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	ASN
1	A	29	HIS
1	A	37	GLN
1	A	112	ASN
1	A	122	GLN
1	A	171	ASN
1	A	186	HIS
1	A	220	ASN
1	A	285	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	307/307 (100%)	-0.78	4 (1%) 77 79	10, 18, 47, 97	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	133	THR	3.5
1	A	137	LEU	2.7
1	A	135	SER	2.6
1	A	2	ARG	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.