



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2024 – 09:59 PM EDT

PDB ID : 6AXJ  
Title : Crystal structure of the Yaf9 YEATS domain bound to H3K27ac  
Authors : Klein, B.J.; Kutateladze, T.G.  
Deposited on : 2017-09-06  
Resolution : 2.38 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.38 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5509 (2.40-2.36)
Clashscore	141614	6082 (2.40-2.36)
Ramachandran outliers	138981	5973 (2.40-2.36)
Sidechain outliers	138945	5975 (2.40-2.36)
RSRZ outliers	127900	5397 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	169	
1	B	169	
1	C	169	
1	D	169	
2	E	11	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	11	 36% 9% 55%
2	G	11	 36% 9% 55%
2	H	11	 36% 9% 55%

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5624 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein AF-9 homolog.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	143	Total	C	N	O	S	0	0	0
			1180	770	196	212	2			
1	C	142	Total	C	N	O	S	0	0	0
			1175	767	195	211	2			
1	B	142	Total	C	N	O	S	0	2	0
			1193	777	197	217	2			
1	D	142	Total	C	N	O	S	0	0	0
			1175	767	195	211	2			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	3	GLY	-	expression tag	UNP P53930
A	4	PRO	-	expression tag	UNP P53930
A	5	LEU	-	expression tag	UNP P53930
A	6	GLY	-	expression tag	UNP P53930
A	7	SER	-	expression tag	UNP P53930
C	3	GLY	-	expression tag	UNP P53930
C	4	PRO	-	expression tag	UNP P53930
C	5	LEU	-	expression tag	UNP P53930
C	6	GLY	-	expression tag	UNP P53930
C	7	SER	-	expression tag	UNP P53930
B	3	GLY	-	expression tag	UNP P53930
B	4	PRO	-	expression tag	UNP P53930
B	5	LEU	-	expression tag	UNP P53930
B	6	GLY	-	expression tag	UNP P53930
B	7	SER	-	expression tag	UNP P53930
D	3	GLY	-	expression tag	UNP P53930
D	4	PRO	-	expression tag	UNP P53930
D	5	LEU	-	expression tag	UNP P53930
D	6	GLY	-	expression tag	UNP P53930
D	7	SER	-	expression tag	UNP P53930

- Molecule 2 is a protein called ALY-SER-ALA-PRO-ALA.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	G	5	Total	C	N	O	0	0	0
			35	22	6	7			
2	E	5	Total	C	N	O	0	0	0
			35	22	6	7			
2	H	5	Total	C	N	O	0	0	0
			35	22	6	7			
2	F	5	Total	C	N	O	0	0	0
			35	22	6	7			

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	179	Total	O	0	0
			179	179		
5	C	162	Total	O	0	0
			162	162		

*Continued on next page...*


*Continued from previous page...*


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	147	Total 147	O 147	0	0
5	D	177	Total 177	O 177	0	0
5	G	4	Total 4	O 4	0	0
5	E	6	Total 6	O 6	0	0
5	H	3	Total 3	O 3	0	0
5	F	3	Total 3	O 3	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Protein AF-9 homolog

Chain A: 





- Molecule 1: Protein AF-9 homolog

Chain C: 




- Molecule 1: Protein AF-9 homolog

Chain B: 





- Molecule 1: Protein AF-9 homolog

Chain D: 




- Molecule 2: ALY-SER-ALA-PRO-ALA

Chain G: 






- Molecule 2: ALY-SER-ALA-PRO-ALA

Chain E:  36% 9% 55%




- Molecule 2: ALY-SER-ALA-PRO-ALA

Chain H:  36% 9% 55%



- Molecule 2: ALY-SER-ALA-PRO-ALA

Chain F:  36% 9% 55%



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	140.32Å 140.32Å 129.05Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	35.08 – 2.38 35.08 – 2.38	Depositor EDS
% Data completeness (in resolution range)	98.2 (35.08-2.38) 81.3 (35.08-2.38)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.37 (at 2.39Å)	Xtriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, $R_{free}$	0.179 , 0.227 0.182 , 0.229	Depositor DCC
$R_{free}$ test set	2015 reflections (5.38%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.3	Xtriage
Anisotropy	0.330	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 56.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.56$ , $\langle L^2 \rangle = 0.40$	Xtriage
Estimated twinning fraction	0.468 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	5624	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 55.52 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.1238e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ALY, SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/1216	0.56	0/1650
1	B	0.42	0/1229	0.58	0/1667
1	C	0.43	0/1211	0.59	0/1643
1	D	0.42	0/1211	0.58	0/1643
2	E	0.38	0/23	0.44	0/31
2	F	0.33	0/23	0.39	0/31
2	G	0.37	0/23	0.50	0/31
2	H	0.37	0/23	0.29	0/31
All	All	0.42	0/4959	0.58	0/6727

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1180	0	1157	11	0
1	B	1193	0	1162	5	0
1	C	1175	0	1152	9	0
1	D	1175	0	1152	19	0
2	E	35	0	37	3	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	35	0	37	2	0
2	G	35	0	37	0	0
2	H	35	0	37	1	0
3	A	10	0	0	0	0
3	C	10	0	0	0	0
4	A	12	0	16	0	0
4	B	24	0	32	0	0
4	C	6	0	8	0	0
4	D	18	0	24	2	0
5	A	179	0	0	1	2
5	B	147	0	0	0	1
5	C	162	0	0	1	0
5	D	177	0	0	4	0
5	E	6	0	0	0	0
5	F	3	0	0	0	0
5	G	4	0	0	0	0
5	H	3	0	0	0	0
All	All	5624	0	4851	40	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:91:GLU:O	2:E:27:ALY:HCA	1.90	0.71
1:D:116:ARG:HH22	1:D:121:ALA:H	1.42	0.68
1:D:27:MET:HE3	1:D:42:LEU:HB2	1.76	0.67
1:A:34:ASN:OD1	5:A:301:HOH:O	2.14	0.66
1:D:91:GLU:O	2:H:27:ALY:HCA	1.99	0.63
1:A:89:TRP:N	2:E:27:ALY:OH	2.30	0.60
1:D:16:ARG:NH2	1:D:166:LEU:O	2.29	0.60
1:D:110:ASN:ND2	5:D:302:HOH:O	2.28	0.59
1:D:106:GLU:OE2	5:D:301:HOH:O	2.17	0.58
1:D:169:ARG:NH2	5:D:305:HOH:O	2.40	0.55
1:C:119:PRO:HA	1:C:145:VAL:HG11	1.89	0.54
1:D:116:ARG:HH22	1:D:121:ALA:N	2.05	0.54
1:A:89:TRP:CZ2	1:C:72:ASN:HB2	2.44	0.53
1:A:89:TRP:CE2	1:C:72:ASN:HB2	2.44	0.52
1:B:116:ARG:HH22	1:B:121:ALA:H	1.56	0.52
1:B:72:ASN:HB2	1:D:89:TRP:CZ2	2.46	0.51

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:16:ARG:NH2	1:A:166:LEU:O	2.30	0.51
1:C:70:TYR:CZ	2:F:27:ALY:HH32	2.49	0.48
1:C:91:GLU:O	2:F:27:ALY:HCA	2.12	0.48
1:B:72:ASN:HB2	1:D:89:TRP:CE2	2.49	0.48
1:D:149:TYR:CD2	4:D:202:GOL:H12	2.48	0.48
1:A:55:ILE:HD12	1:A:59:ILE:HD12	1.95	0.48
1:D:27:MET:CE	1:D:42:LEU:HB2	2.43	0.48
1:C:144:GLU:N	5:C:305:HOH:O	2.47	0.47
1:D:161:GLU:HG3	5:D:409:HOH:O	2.14	0.47
1:D:16:ARG:HD3	1:D:16:ARG:HA	1.77	0.47
1:A:119:PRO:HA	1:A:145:VAL:HG11	1.96	0.46
1:C:45:ILE:HD13	1:C:94:ILE:HD12	1.98	0.46
1:A:16:ARG:HD3	1:A:16:ARG:HA	1.78	0.44
1:B:10:LYS:HB2	1:B:10:LYS:HE3	1.81	0.44
1:D:116:ARG:NH2	1:D:120:TYR:HA	2.32	0.44
1:D:67:HIS:HB2	1:D:92:PHE:CD2	2.52	0.44
1:C:10:LYS:HE3	1:C:10:LYS:HB2	1.77	0.43
1:B:15:SER:HA	1:B:153:ILE:O	2.19	0.43
1:D:149:TYR:HD2	4:D:202:GOL:H12	1.84	0.43
1:D:15:SER:HA	1:D:153:ILE:O	2.19	0.42
1:C:22:ASN:HA	1:C:44:THR:O	2.20	0.42
1:A:89:TRP:CH2	2:E:27:ALY:HG3	2.55	0.42
1:D:10:LYS:HB2	1:D:10:LYS:HE3	1.67	0.42
1:A:22:ASN:HA	1:A:44:THR:O	2.20	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:304:HOH:O	5:A:381:HOH:O[3_655]	2.06	0.14
5:A:438:HOH:O	5:B:371:HOH:O[3_655]	2.14	0.06

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	139/169 (82%)	136 (98%)	3 (2%)	0	100	100
1	B	140/169 (83%)	137 (98%)	3 (2%)	0	100	100
1	C	138/169 (82%)	135 (98%)	3 (2%)	0	100	100
1	D	138/169 (82%)	135 (98%)	3 (2%)	0	100	100
2	E	3/11 (27%)	3 (100%)	0	0	100	100
2	F	3/11 (27%)	3 (100%)	0	0	100	100
2	G	3/11 (27%)	3 (100%)	0	0	100	100
2	H	3/11 (27%)	3 (100%)	0	0	100	100
All	All	567/720 (79%)	555 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	130/153 (85%)	129 (99%)	1 (1%)	81	91
1	B	132/153 (86%)	131 (99%)	1 (1%)	81	91
1	C	130/153 (85%)	129 (99%)	1 (1%)	81	91
1	D	130/153 (85%)	130 (100%)	0	100	100
2	E	2/5 (40%)	2 (100%)	0	100	100
2	F	2/5 (40%)	2 (100%)	0	100	100
2	G	2/5 (40%)	2 (100%)	0	100	100
2	H	2/5 (40%)	2 (100%)	0	100	100
All	All	530/632 (84%)	527 (99%)	3 (1%)	86	93

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	ASN
1	C	22	ASN
1	B	22	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	ALY	F	27	2	10,11,12	3.43	3 (30%)	7,12,14	2.77	3 (42%)
2	ALY	H	27	2	10,11,12	3.18	2 (20%)	7,12,14	2.26	2 (28%)
2	ALY	G	27	2	10,11,12	2.91	2 (20%)	7,12,14	2.01	1 (14%)
2	ALY	E	27	2	10,11,12	3.20	2 (20%)	7,12,14	2.87	3 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ALY	F	27	2	-	0/9/10/12	-
2	ALY	H	27	2	-	0/9/10/12	-
2	ALY	G	27	2	-	1/9/10/12	-
2	ALY	E	27	2	-	1/9/10/12	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	27	ALY	OH-CH	8.67	1.42	1.23
2	E	27	ALY	OH-CH	8.11	1.41	1.23
2	H	27	ALY	OH-CH	7.70	1.40	1.23
2	G	27	ALY	OH-CH	6.92	1.38	1.23
2	H	27	ALY	CH-NZ	6.06	1.51	1.34
2	G	27	ALY	CH-NZ	5.86	1.51	1.34
2	F	27	ALY	CH-NZ	5.74	1.50	1.34
2	E	27	ALY	CH-NZ	5.52	1.50	1.34
2	F	27	ALY	CE-NZ	2.37	1.51	1.46

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	27	ALY	CE-NZ-CH	-5.18	114.59	122.56
2	F	27	ALY	CE-NZ-CH	-4.99	114.88	122.56
2	E	27	ALY	OH-CH-CH3	-4.64	113.44	122.06
2	G	27	ALY	CE-NZ-CH	-4.59	115.50	122.56
2	H	27	ALY	CE-NZ-CH	-4.40	115.79	122.56
2	F	27	ALY	CH3-CH-NZ	-4.00	109.01	116.09
2	H	27	ALY	OH-CH-CH3	-3.19	116.13	122.06
2	E	27	ALY	OH-CH-NZ	2.76	129.52	121.74
2	F	27	ALY	CD-CE-NZ	-2.72	104.44	112.21

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	G	27	ALY	O-C-CA-CB
2	E	27	ALY	CG-CD-CE-NZ

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	27	ALY	2	0
2	H	27	ALY	1	0
2	E	27	ALY	3	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.



## 5.6 Ligand geometry

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	SO4	A	202	-	4,4,4	0.12	0	6,6,6	0.13	0
4	GOL	B	202	-	5,5,5	0.39	0	5,5,5	0.41	0
4	GOL	D	203	-	5,5,5	0.38	0	5,5,5	0.40	0
4	GOL	B	201	-	5,5,5	0.37	0	5,5,5	0.36	0
4	GOL	D	201	-	5,5,5	0.40	0	5,5,5	0.35	0
3	SO4	C	202	-	4,4,4	0.32	0	6,6,6	0.05	0
3	SO4	C	201	-	4,4,4	0.18	0	6,6,6	0.30	0
4	GOL	B	203	-	5,5,5	0.35	0	5,5,5	0.57	0
4	GOL	D	202	-	5,5,5	0.34	0	5,5,5	0.33	0
4	GOL	C	203	-	5,5,5	0.30	0	5,5,5	0.63	0
4	GOL	B	204	-	5,5,5	0.33	0	5,5,5	0.58	0
4	GOL	A	203	-	5,5,5	0.34	0	5,5,5	0.42	0
4	GOL	A	204	-	5,5,5	0.34	0	5,5,5	0.21	0
3	SO4	A	201	-	4,4,4	0.31	0	6,6,6	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	D	203	-	-	2/4/4/4	-
4	GOL	B	201	-	-	4/4/4/4	-
4	GOL	D	201	-	-	3/4/4/4	-
4	GOL	B	203	-	-	2/4/4/4	-
4	GOL	D	202	-	-	4/4/4/4	-
4	GOL	C	203	-	-	2/4/4/4	-
4	GOL	B	204	-	-	2/4/4/4	-
4	GOL	A	203	-	-	2/4/4/4	-
4	GOL	A	204	-	-	2/4/4/4	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	202	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (27) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	203	GOL	O1-C1-C2-C3
4	A	204	GOL	O1-C1-C2-C3
4	B	201	GOL	O1-C1-C2-C3
4	B	202	GOL	O1-C1-C2-C3
4	B	202	GOL	C1-C2-C3-O3
4	B	203	GOL	O1-C1-C2-C3
4	D	202	GOL	O1-C1-C2-O2
4	D	202	GOL	O1-C1-C2-C3
4	D	203	GOL	O1-C1-C2-O2
4	D	203	GOL	O1-C1-C2-C3
4	A	203	GOL	O1-C1-C2-O2
4	A	204	GOL	O1-C1-C2-O2
4	D	202	GOL	O2-C2-C3-O3
4	B	201	GOL	C1-C2-C3-O3
4	B	204	GOL	C1-C2-C3-O3
4	D	201	GOL	C1-C2-C3-O3
4	D	202	GOL	C1-C2-C3-O3
4	B	201	GOL	O1-C1-C2-O2
4	B	202	GOL	O1-C1-C2-O2
4	B	202	GOL	O2-C2-C3-O3
4	B	203	GOL	O1-C1-C2-O2
4	B	204	GOL	O2-C2-C3-O3
4	C	203	GOL	O1-C1-C2-O2
4	B	201	GOL	O2-C2-C3-O3
4	D	201	GOL	O2-C2-C3-O3
4	C	203	GOL	O1-C1-C2-C3
4	D	201	GOL	O1-C1-C2-C3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	202	GOL	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	143/169 (84%)	-0.51	0 100 100	28, 40, 78, 94	0
1	B	142/169 (84%)	-0.56	0 100 100	28, 40, 76, 94	0
1	C	142/169 (84%)	-0.54	0 100 100	28, 40, 76, 106	0
1	D	142/169 (84%)	-0.55	0 100 100	29, 41, 73, 125	0
2	E	4/11 (36%)	-0.32	0 100 100	40, 41, 46, 70	0
2	F	4/11 (36%)	0.08	0 100 100	44, 47, 51, 75	0
2	G	4/11 (36%)	-0.03	0 100 100	41, 44, 47, 66	0
2	H	4/11 (36%)	0.08	0 100 100	39, 45, 45, 78	0
All	All	585/720 (81%)	-0.52	0 100 100	28, 40, 77, 125	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ALY	E	27	12/13	0.95	0.14	24,32,39,42	0
2	ALY	H	27	12/13	0.96	0.12	31,36,39,39	0
2	ALY	F	27	12/13	0.96	0.13	31,37,41,46	0
2	ALY	G	27	12/13	0.98	0.13	32,37,40,41	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	D	202	6/6	0.86	0.11	82,87,89,89	0
4	GOL	D	203	6/6	0.87	0.15	86,89,92,94	0
4	GOL	D	201	6/6	0.92	0.12	44,57,59,63	0
4	GOL	A	203	6/6	0.92	0.10	78,81,86,86	0
4	GOL	B	202	6/6	0.92	0.14	64,71,75,75	0
4	GOL	B	204	6/6	0.95	0.10	67,74,80,83	0
4	GOL	B	203	6/6	0.95	0.11	63,68,70,71	0
4	GOL	B	201	6/6	0.96	0.15	50,68,71,75	0
4	GOL	C	203	6/6	0.97	0.12	47,60,69,69	0
4	GOL	A	204	6/6	0.97	0.13	51,65,67,70	0
3	SO4	A	201	5/5	0.99	0.12	33,34,38,39	0
3	SO4	C	202	5/5	0.99	0.12	31,36,38,40	0
3	SO4	C	201	5/5	1.00	0.12	32,34,38,42	0
3	SO4	A	202	5/5	1.00	0.12	26,32,40,40	0

## 6.5 Other polymers

There are no such residues in this entry.