



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 22, 2024 – 10:13 AM EDT

PDB ID : 6B05  
Title : The Crystal Structure of the Ferredoxin Protease FusC E83A mutant in complex with Arabidopsis Ferredoxin  
Authors : Grinter, R.  
Deposited on : 2017-09-13  
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	924	<div> <div>2%</div> <div>85%</div> <div>12%</div> <div>.</div> </div>
2	B	104	<div> <div>28%</div> <div>28%</div> <div>11%</div> <div>.</div> <div>61%</div> </div>
2	C	104	<div> <div>5%</div> <div>7%</div> <div>93%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 8533 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative zinc protease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	896	7203	4522	1291	1367	23	0	12	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	83	ALA	GLU	conflict	UNP Q6D8U3

- Molecule 2 is a protein called Ferredoxin-2, chloroplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	41	324	205	44	74	1	0	1	0
2	C	7	48	30	7	11		0	0	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	95	ALA	-	expression tag	UNP P16972
B	96	ILE	-	expression tag	UNP P16972
B	97	MET	-	expression tag	UNP P16972
B	98	LEU	-	expression tag	UNP P16972
B	99	GLU	-	expression tag	UNP P16972
B	100	HIS	-	expression tag	UNP P16972
B	101	HIS	-	expression tag	UNP P16972
B	102	HIS	-	expression tag	UNP P16972
B	103	HIS	-	expression tag	UNP P16972
B	104	HIS	-	expression tag	UNP P16972
B	105	HIS	-	expression tag	UNP P16972
C	95	ALA	-	expression tag	UNP P16972
C	96	ILE	-	expression tag	UNP P16972

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	97	MET	-	expression tag	UNP P16972
C	98	LEU	-	expression tag	UNP P16972
C	99	GLU	-	expression tag	UNP P16972
C	100	HIS	-	expression tag	UNP P16972
C	101	HIS	-	expression tag	UNP P16972
C	102	HIS	-	expression tag	UNP P16972
C	103	HIS	-	expression tag	UNP P16972
C	104	HIS	-	expression tag	UNP P16972
C	105	HIS	-	expression tag	UNP P16972

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	938	Total	O	0	0
			938	938		
4	B	16	Total	O	0	0
			16	16		
4	C	3	Total	O	0	0
			3	3		



GLN	SER	PHE	LEU	ASP	GLU	GLN	ILE	GLY	GLU	GLY	PHE	VAL	LEU	THR	CYS	ALA	ALA	TYR	PRO	THR	SER	D85	V86	E89	T90	H91	LYS	GLU	GLU	ALA	ILE	MET	LEU	GLU	HIS	HIS	HIS	HIS
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.34Å 126.50Å 133.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.99 – 1.90 39.99 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.0 (39.99-1.90) 99.6 (39.99-1.90)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.43 (at 1.89Å)	Xtriage
Refinement program	PHENIX 1.11.1 _2575	Depositor
R, $R_{free}$	0.195 , 0.239 0.196 , 0.238	Depositor DCC
$R_{free}$ test set	5472 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.6	Xtriage
Anisotropy	0.222	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 51.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8533	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.85% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.40	0/7351	0.58	1/9967 (0.0%)
2	B	0.31	0/327	0.50	0/445
2	C	0.27	0/47	0.44	0/64
All	All	0.39	0/7725	0.58	1/10476 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	529	ARG	NE-CZ-NH2	-6.05	117.27	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7203	0	7192	73	0
2	B	324	0	302	10	0
2	C	48	0	43	0	0
3	A	1	0	0	0	0
4	A	938	0	0	9	1
4	B	16	0	0	0	0
4	C	3	0	0	0	0

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8533	0	7537	80	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 80 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:512:LYS:NZ	4:A:1101:HOH:O	2.02	0.93
1:A:275:GLN:HE22	1:A:652:VAL:H	1.18	0.89
1:A:318:GLN:HE22	1:A:489:ARG:H	1.19	0.89
1:A:314:GLN:HE21	1:A:318:GLN:HE21	1.28	0.79
1:A:409:THR:HG22	1:A:413:GLN:HE21	1.47	0.78

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1126:HOH:O	4:A:1771:HOH:O[2_565]	2.04	0.16

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	906/924 (98%)	896 (99%)	10 (1%)	0	100	100
2	B	38/104 (36%)	35 (92%)	2 (5%)	1 (3%)	5	1
2	C	5/104 (5%)	5 (100%)	0	0	100	100
All	All	949/1132 (84%)	936 (99%)	12 (1%)	1 (0%)	51	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	36	LEU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	776/788 (98%)	769 (99%)	7 (1%)	78	79
2	B	36/88 (41%)	35 (97%)	1 (3%)	43	36
2	C	5/88 (6%)	5 (100%)	0	100	100
All	All	817/964 (85%)	809 (99%)	8 (1%)	78	76

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	21	ASP
1	A	781	LYS
1	A	364	GLU
1	A	257	TYR
1	A	776	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	477	GLN
1	A	523	GLN
1	A	710	GLN
1	A	601	GLN
1	A	318	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	896/924 (96%)	0.11	22 (2%) 57 60	18, 32, 59, 89	1 (0%)
2	B	41/104 (39%)	3.26	29 (70%) 0 0	29, 92, 136, 140	0
2	C	7/104 (6%)	2.90	5 (71%) 0 0	51, 74, 93, 95	0
All	All	944/1132 (83%)	0.26	56 (5%) 22 25	18, 33, 68, 140	1 (0%)

The worst 5 of 56 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	88	ILE	8.6
2	B	8	PHE	8.2
2	B	86	VAL	8.0
2	B	87	THR	7.6
2	B	89	GLU	7.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ZN	A	1001	1/1	0.99	0.06	37,37,37,37	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.