



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 26, 2024 – 01:04 AM EDT

PDB ID : 7B38
Title : Torpedo californica acetylcholinesterase complexed with Mg+2
Authors : Silman, I.; Shnyrov, V.L.; Ashani, Y.; Roth, E.; Nicolas, A.; Sussman, J.L.; Weiner, L.
Deposited on : 2020-11-29
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	532	<div> <div>8%</div> <div>89%</div> <div>10%</div> </div>
2	B	2	<div> <div>100%</div> </div>

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 4742 atoms, of which 12 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetylcholinesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	532	Total	C	N	O	S	0	3	0
			4206	2704	711	768	23			

- Molecule 2 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			24	14	1	9			

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Mg	0	0
			4	4		

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C O 4 2 2	0	0
8	A	1	Total C H O 10 2 6 2	0	0
8	A	1	Total C O 4 2 2	0	0
8	A	1	Total C O 4 2 2	0	0
8	A	1	Total C H O 10 2 6 2	0	0
8	A	1	Total C O 4 2 2	0	0

- Molecule 9 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	1	Total C O 7 4 3	0	0
9	A	1	Total C O 7 4 3	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	397	Total O 397 397	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Acetylcholinesterase



• Molecule 2: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	138.22Å 138.22Å 71.38Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	34.83 – 1.85 34.83 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.6 (34.83-1.85) 99.6 (34.83-1.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.13 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.8.0267, PHENIX 1.15.2_3472	Depositor
R, R_{free}	0.202 , 0.241 0.206 , 0.244	Depositor DCC
R_{free} test set	3292 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	30.0	Xtriage
Anisotropy	0.410	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 50.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.024 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4742	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, FUC, GOL, PEG, ZN, EDO, NAG, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/4344	0.55	0/5903

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4206	0	4015	51	0
2	B	24	0	22	0	0
3	A	4	0	0	0	0
4	A	1	0	0	0	0
5	A	28	0	26	0	0
6	A	10	0	14	4	0
7	A	18	0	24	3	0
8	A	28	12	42	4	0
9	A	14	0	20	1	0
10	A	397	0	0	4	0
All	All	4730	12	4163	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:330:PHE:CD1	6:A:605:PGE:H12	1.96	1.01
1:A:330:PHE:CE1	6:A:605:PGE:H12	2.04	0.93
1:A:384:GLY:H	8:A:609:EDO:H21	1.41	0.83
1:A:238:VAL:HG12	7:A:611:GOL:H11	1.66	0.76
1:A:498:LYS:HG2	1:A:499:GLU:HG3	1.68	0.76
1:A:498:LYS:HD2	10:A:1009:HOH:O	1.87	0.73
1:A:384:GLY:N	8:A:609:EDO:H21	2.04	0.72
1:A:255:ASN:HB3	1:A:257:ASN:HD21	1.55	0.72
1:A:135:LEU:HA	1:A:470:MET:CE	2.21	0.70
1:A:27:ILE:HD11	1:A:100:TRP:HD1	1.56	0.69
1:A:271:PRO:HD3	8:A:615:EDO:H22	1.76	0.68
1:A:330:PHE:CE1	6:A:605:PGE:C1	2.77	0.68
1:A:255:ASN:HB3	1:A:257:ASN:ND2	2.12	0.65
7:A:616:GOL:H11	10:A:761:HOH:O	1.98	0.62
1:A:135:LEU:HD13	1:A:470:MET:HE3	1.81	0.62
1:A:405:MET:HA	1:A:408:VAL:HG12	1.83	0.61
1:A:496:THR:HB	1:A:498:LYS:HD3	1.86	0.58
1:A:238:VAL:HG12	7:A:611:GOL:C1	2.34	0.58
1:A:264:HIS:NE2	1:A:268:GLU:OE2	2.36	0.58
1:A:142:VAL:HG11	1:A:184:ILE:HD11	1.86	0.57
1:A:515:ARG:HB3	1:A:518:VAL:HB	1.89	0.54
1:A:220:ARG:HG3	1:A:221:ARG:HG3	1.89	0.53
1:A:502:PHE:CZ	1:A:513:HIS:HB2	2.43	0.53
1:A:135:LEU:HD13	1:A:470:MET:CE	2.38	0.53
1:A:211:LEU:HD23	1:A:314:PHE:HB3	1.91	0.52
1:A:228:SER:HB2	1:A:229:PRO:HD2	1.92	0.51
1:A:197:PHE:HB2	1:A:223:ILE:HB	1.93	0.50
1:A:257:ASN:OD1	1:A:261:GLU:HG3	2.12	0.49
1:A:496:THR:CB	1:A:498:LYS:HD3	2.43	0.49
6:A:605:PGE:H52	10:A:1041:HOH:O	2.13	0.48
1:A:197:PHE:CB	1:A:223:ILE:HB	2.45	0.47
1:A:135:LEU:HA	1:A:470:MET:HE2	1.95	0.47
1:A:498:LYS:HG2	1:A:499:GLU:N	2.30	0.47
1:A:498:LYS:HE2	1:A:499:GLU:OE2	2.16	0.46
1:A:531:LEU:C	1:A:531:LEU:HD23	2.36	0.45
1:A:146:LEU:C	1:A:146:LEU:HD12	2.36	0.45
1:A:451:PRO:HA	1:A:458:TYR:CD1	2.52	0.45
1:A:408:VAL:CG2	1:A:418:THR:HG21	2.47	0.45
1:A:212:SER:HB2	1:A:300:PHE:CE1	2.51	0.45
1:A:330:PHE:CE1	1:A:334:TYR:CE1	3.05	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:330:PHE:HE1	1:A:334:TYR:CE1	2.36	0.44
1:A:346:LYS:NZ	9:A:612:PEG:H12	2.33	0.43
1:A:487:SER:OG	1:A:489:GLU:HG2	2.18	0.43
1:A:199:GLU:HA	1:A:225:GLN:O	2.19	0.43
1:A:223:ILE:HA	1:A:320:LEU:O	2.18	0.43
1:A:344:GLU:OE2	1:A:346:LYS:HE3	2.18	0.43
1:A:44:ARG:O	1:A:45:PHE:HB2	2.19	0.42
1:A:384:GLY:H	8:A:609:EDO:C2	2.23	0.42
1:A:257:ASN:ND2	1:A:257:ASN:H	2.18	0.41
1:A:135:LEU:HA	1:A:470:MET:HE1	2.01	0.41
1:A:420:LEU:HD23	1:A:502:PHE:HB3	2.01	0.41
1:A:212:SER:O	1:A:216:ARG:HG3	2.21	0.41
1:A:34:PRO:HD3	1:A:58:TRP:CE2	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	533/532 (100%)	512 (96%)	21 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	449/465 (97%)	442 (98%)	7 (2%)	62 49

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	103	SER
1	A	197	PHE
1	A	200	SER
1	A	288	PHE
1	A	330	PHE
1	A	473	TRP
1	A	498	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	253	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	1,2	14,14,15	0.39	0	17,19,21	0.53	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FUC	B	2	2	10,10,11	0.77	0	14,14,16	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	2/6/23/26	0/1/1/1
2	FUC	B	2	2	-	-	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	NAG	C4-C5-C6-O6
2	B	1	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

Of 20 ligands modelled in this entry, 5 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	PEG	A	612	-	6,6,6	0.47	0	5,5,5	0.27	0
7	GOL	A	616	-	5,5,5	0.75	0	5,5,5	1.13	0
8	EDO	A	617	-	3,3,3	0.50	0	2,2,2	0.24	0
8	EDO	A	615	-	3,3,3	0.51	0	2,2,2	0.31	0
8	EDO	A	610	-	3,3,3	0.56	0	2,2,2	0.27	0
5	NAG	A	603	1	14,14,15	0.58	0	17,19,21	0.59	0
8	EDO	A	607	-	3,3,3	0.60	0	2,2,2	0.09	0
9	PEG	A	608	-	6,6,6	0.49	0	5,5,5	0.17	0
8	EDO	A	609	-	3,3,3	0.41	0	2,2,2	0.12	0
7	GOL	A	611	-	5,5,5	1.03	0	5,5,5	0.93	0
5	NAG	A	604	1	14,14,15	0.25	0	17,19,21	0.61	0
6	PGE	A	605	-	9,9,9	0.37	0	8,8,8	0.28	0
7	GOL	A	606	-	5,5,5	0.70	0	5,5,5	1.13	1 (20%)
8	EDO	A	614	-	3,3,3	0.48	0	2,2,2	0.37	0
8	EDO	A	613	-	3,3,3	0.51	0	2,2,2	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	PEG	A	612	-	-	2/4/4/4	-
7	GOL	A	616	-	-	2/4/4/4	-
8	EDO	A	617	-	-	1/1/1/1	-
8	EDO	A	615	-	-	1/1/1/1	-
8	EDO	A	610	-	-	0/1/1/1	-
5	NAG	A	603	1	-	0/6/23/26	0/1/1/1
8	EDO	A	607	-	-	1/1/1/1	-
9	PEG	A	608	-	-	2/4/4/4	-
8	EDO	A	609	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	A	611	-	-	2/4/4/4	-
5	NAG	A	604	1	-	2/6/23/26	0/1/1/1
6	PGE	A	605	-	-	4/7/7/7	-
7	GOL	A	606	-	-	0/4/4/4	-
8	EDO	A	614	-	-	1/1/1/1	-
8	EDO	A	613	-	-	0/1/1/1	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	606	GOL	C3-C2-C1	-2.13	103.44	111.70

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	616	GOL	O1-C1-C2-C3
5	A	604	NAG	C8-C7-N2-C2
5	A	604	NAG	O7-C7-N2-C2
9	A	612	PEG	O1-C1-C2-O2
7	A	611	GOL	O1-C1-C2-C3
9	A	608	PEG	O2-C3-C4-O4
7	A	611	GOL	O1-C1-C2-O2
7	A	616	GOL	O1-C1-C2-O2
6	A	605	PGE	O2-C3-C4-O3
8	A	614	EDO	O1-C1-C2-O2
6	A	605	PGE	C3-C4-O3-C5
6	A	605	PGE	C6-C5-O3-C4
9	A	608	PEG	C4-C3-O2-C2
8	A	607	EDO	O1-C1-C2-O2
8	A	615	EDO	O1-C1-C2-O2
9	A	612	PEG	C1-C2-O2-C3
6	A	605	PGE	O1-C1-C2-O2
8	A	617	EDO	O1-C1-C2-O2

There are no ring outliers.

6 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	A	612	PEG	1	0
7	A	616	GOL	1	0
8	A	615	EDO	1	0
8	A	609	EDO	3	0
7	A	611	GOL	2	0
6	A	605	PGE	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	532/532 (100%)	0.27	45 (8%)	10 10	20, 34, 53, 68	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	4	SER	5.0
1	A	114	TRP	4.0
1	A	115	ILE	4.0
1	A	206	VAL	3.9
1	A	197	PHE	3.5
1	A	106	PRO	3.5
1	A	55	SER	3.5
1	A	485	PRO	3.4
1	A	57	VAL	3.4
1	A	224	LEU	3.4
1	A	200	SER	3.3
1	A	535	THR	3.2
1	A	116	TYR	3.0
1	A	113	VAL	2.9
1	A	226	SER	2.9
1	A	146	LEU	2.8
1	A	16[A]	MET	2.7
1	A	223	ILE	2.7
1	A	7	LEU	2.7
1	A	198	GLY	2.6
1	A	6	LEU	2.5
1	A	119	GLY	2.5
1	A	201	ALA	2.4
1	A	54	TRP	2.4
1	A	120	PHE	2.4
1	A	401	ILE	2.4
1	A	186	PHE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	160	GLY	2.3
1	A	202	GLY	2.3
1	A	231	CYS	2.3
1	A	444	ILE	2.2
1	A	199	GLU	2.2
1	A	56	GLY	2.2
1	A	323	VAL	2.1
1	A	24	SER	2.1
1	A	15	VAL	2.1
1	A	196	ILE	2.1
1	A	204	ALA	2.1
1	A	12	SER	2.1
1	A	130	TYR	2.1
1	A	117	GLY	2.1
1	A	148	TYR	2.1
1	A	404	LEU	2.0
1	A	203	GLY	2.0
1	A	207	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	B	1	14/15	0.79	0.24	52,58,63,67	0
2	FUC	B	2	10/11	0.85	0.41	69,72,77,78	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
9	PEG	A	612	7/7	0.60	0.30	62,68,70,70	0
5	NAG	A	604	14/15	0.77	0.32	60,63,70,73	0
9	PEG	A	608	7/7	0.78	0.22	55,59,63,68	0
7	GOL	A	611	6/6	0.82	0.34	47,51,55,56	0
3	MG	A	619	1/1	0.83	0.07	59,59,59,59	0
8	EDO	A	607	4/4	0.83	0.40	40,41,45,48	0
3	MG	A	618	1/1	0.86	0.12	51,51,51,51	0
8	EDO	A	610	4/4	0.86	0.24	40,48,51,54	0
7	GOL	A	616	6/6	0.87	0.39	39,49,51,51	0
6	PGE	A	605	10/10	0.88	0.14	41,41,46,47	0
8	EDO	A	615	4/4	0.89	0.20	31,51,61,62	0
8	EDO	A	617	4/4	0.89	0.10	49,49,51,51	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	NAG	A	603	14/15	0.91	0.19	43,50,57,59	0
8	EDO	A	614	4/4	0.91	0.28	44,44,48,59	0
7	GOL	A	606	6/6	0.93	0.09	36,42,50,52	0
4	ZN	A	602	1/1	0.93	0.09	45,45,45,45	1
8	EDO	A	613	4/4	0.94	0.30	42,46,54,55	0
8	EDO	A	609	4/4	0.95	0.28	44,44,47,53	0
3	MG	A	620	1/1	0.95	0.07	32,32,32,32	0
3	MG	A	601	1/1	0.99	0.06	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around MG A 619:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG A 618:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MG A 620:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MG A 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.