



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 3, 2024 – 10:37 AM EST

PDB ID : 2BBH
Title : X-ray structure of T.maritima CorA soluble domain
Authors : Lunin, V.V.; Dobrovetsky, E.; Khutoreskaya, G.; Bochkarev, A.; Maguire, M.E.; Edwards, A.M.; Koth, C.M.; Structural Genomics Consortium (SGC)
Deposited on : 2005-10-17
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3097 (1.86-1.86)
Clashscore	180529	3359 (1.86-1.86)
Ramachandran outliers	177936	3335 (1.86-1.86)
Sidechain outliers	177891	3335 (1.86-1.86)
RSRZ outliers	164620	3097 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	269	<div> <div>7%</div> <div>66%</div> <div>15%</div> <div>•</div> <div>17%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	DMU	A	401	X	-	-	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	DMU	A	402	X	-	-	-
2	DMU	A	403	X	-	-	-
2	DMU	A	404	X	-	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2285 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called divalent cation transport-related protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	223	1915	1233	321	357	1	3	0	4	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	cloning artifact	UNP Q9WZ31
A	-1	SER	-	cloning artifact	UNP Q9WZ31
A	0	HIS	-	cloning artifact	UNP Q9WZ31
A	1	MSE	MET	modified residue	UNP Q9WZ31
A	32	MSE	MET	modified residue	UNP Q9WZ31
A	82	THR	ILE	cloning artifact	UNP Q9WZ31
A	112	MSE	MET	modified residue	UNP Q9WZ31
A	138	MSE	MET	modified residue	UNP Q9WZ31

- Molecule 2 is DECYL-BETA-D-MALTOPYRANOSIDE (three-letter code: DMU) (formula: $C_{22}H_{42}O_{11}$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			33	22	11		
2	A	1	Total	C	O	0	0
			33	22	11		
2	A	1	Total	C	O	0	0
			33	22	11		
2	A	1	Total	C	O	0	0
			33	22	11		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	236	Total	O	0	0
			236	236		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: divalent cation transport-related protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	78.00Å 78.00Å 101.61Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.60 – 1.85 42.60 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.8 (42.60-1.85) 99.8 (42.60-1.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.47 (at 1.86Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.196 , 0.233 0.195 , 0.233	Depositor DCC
R_{free} test set	1378 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	28.4	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 45.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2285	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, DMU, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	0/1951	0.82	1/2634 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	242	LEU	CA-CB-CG	5.10	127.03	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1915	0	1910	40	0
2	A	132	0	148	12	1
3	A	1	0	0	0	0
4	A	1	0	0	0	0
5	A	236	0	0	5	1
All	All	2285	0	2058	44	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:198:GLU:CB	1:A:199:VAL:HA	1.68	1.16
1:A:198:GLU:HB3	1:A:199:VAL:HA	1.23	1.10
1:A:40[B]:ARG:HG3	1:A:40[B]:ARG:HH21	1.19	1.07
1:A:198:GLU:HB2	1:A:199:VAL:HA	1.56	0.86
1:A:198:GLU:CB	1:A:199:VAL:CA	2.55	0.84
1:A:229[B]:ARG:NH1	5:A:603:HOH:O	2.12	0.83
2:A:402:DMU:H30	5:A:530:HOH:O	1.82	0.78
1:A:224:THR:HG21	2:A:402:DMU:H1	1.68	0.76
1:A:241:PRO:HG2	5:A:526:HOH:O	1.93	0.69
1:A:224:THR:CG2	2:A:402:DMU:H1	2.23	0.67
1:A:198:GLU:HB3	1:A:199:VAL:CA	2.14	0.67
1:A:165:ARG:HE	2:A:401:DMU:H30	1.61	0.65
1:A:40[B]:ARG:HG3	1:A:40[B]:ARG:NH2	1.97	0.65
1:A:240:PRO:HB2	1:A:241:PRO:HD3	1.83	0.61
1:A:198:GLU:HB2	1:A:199:VAL:CA	2.30	0.60
1:A:40[B]:ARG:HD2	1:A:41:GLU:N	2.21	0.56
1:A:96:ARG:HA	2:A:403:DMU:H1	1.87	0.55
1:A:124:SER:OG	2:A:402:DMU:O61	2.23	0.53
1:A:115:TYR:N	1:A:115:TYR:CD2	2.76	0.53
1:A:124:SER:HG	2:A:402:DMU:H31	1.49	0.51
1:A:151:ARG:O	1:A:155:ARG:HG3	2.12	0.50
2:A:404:DMU:H28	2:A:404:DMU:C10	2.26	0.47
2:A:404:DMU:O1	2:A:404:DMU:O55	2.22	0.47
1:A:40[B]:ARG:NH2	1:A:40[B]:ARG:CG	2.73	0.45
1:A:61:TRP:HB2	1:A:169:LEU:HD21	1.98	0.45
1:A:152:GLU:OE1	1:A:156:TYR:HE2	2.00	0.45
1:A:51:LEU:N	1:A:52:PRO:CD	2.80	0.45
1:A:124:SER:HB3	2:A:402:DMU:H32	1.98	0.44
1:A:229[B]:ARG:HD3	1:A:229[B]:ARG:C	2.37	0.44
1:A:175:ASP:OD1	2:A:404:DMU:H11	2.16	0.44
1:A:13:PRO:N	5:A:628:HOH:O	2.50	0.43
1:A:92:ASN:O	1:A:95:GLN:HG2	2.19	0.43
1:A:120:HIS:HA	1:A:121:GLU:HA	1.38	0.43
1:A:212:HIS:O	1:A:216:ARG:HG2	2.19	0.43
2:A:401:DMU:H8	5:A:528:HOH:O	2.18	0.43
1:A:196:GLU:HG3	1:A:215:LYS:HD2	2.01	0.43
1:A:210:ARG:HH21	1:A:214:LEU:HD11	1.84	0.42
1:A:31:VAL:O	1:A:41:GLU:HA	2.20	0.42
1:A:30:GLU:OE2	1:A:43:LYS:HE2	2.19	0.42
1:A:24:ARG:HA	1:A:68:HIS:CD2	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:32:MSE:HE3	1:A:39:PHE:CE2	2.55	0.41
1:A:240:PRO:HB2	1:A:241:PRO:CD	2.51	0.41
1:A:192:ILE:HD13	1:A:219:VAL:HG12	2.03	0.40
1:A:194:VAL:O	1:A:198:GLU:HB2	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:403:DMU:O4	5:A:540:HOH:O[7_555]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	221/269 (82%)	216 (98%)	5 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	218/252 (86%)	210 (96%)	8 (4%)	29	14

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	84	PRO
1	A	96	ARG
1	A	115	TYR
1	A	120	HIS
1	A	125	GLU
1	A	210	ARG
1	A	219	VAL
1	A	220	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	209	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	DMU	A	403	-	34,34,34	0.87	1 (2%)	45,45,45	2.89	18 (40%)
2	DMU	A	402	-	34,34,34	1.14	1 (2%)	45,45,45	3.00	17 (37%)
2	DMU	A	401	-	34,34,34	1.08	4 (11%)	45,45,45	2.93	19 (42%)
2	DMU	A	404	-	34,34,34	1.01	2 (5%)	45,45,45	2.92	18 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DMU	A	403	-	7/7/10/10	7/19/59/59	0/2/2/2
2	DMU	A	402	-	7/7/10/10	12/19/59/59	0/2/2/2
2	DMU	A	401	-	7/7/10/10	10/19/59/59	0/2/2/2
2	DMU	A	404	-	7/7/10/10	9/19/59/59	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	402	DMU	O16-C6	4.60	1.47	1.40
2	A	401	DMU	O16-C6	3.45	1.45	1.40
2	A	404	DMU	O16-C6	3.24	1.45	1.40
2	A	401	DMU	O5-C6	2.49	1.48	1.41
2	A	403	DMU	O16-C6	2.43	1.44	1.40
2	A	404	DMU	C3-C4	-2.27	1.46	1.52
2	A	401	DMU	O1-C10	2.25	1.47	1.41
2	A	401	DMU	O7-C10	2.22	1.48	1.41

All (72) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	402	DMU	O16-C6-C1	9.17	122.19	108.27
2	A	403	DMU	O5-C6-C1	7.66	126.11	110.37
2	A	404	DMU	O5-C6-C1	7.01	124.76	110.37
2	A	401	DMU	O16-C6-C1	6.34	117.90	108.27
2	A	401	DMU	O7-C3-C4	6.24	125.85	109.48
2	A	402	DMU	O7-C3-C2	6.16	122.88	107.23
2	A	402	DMU	O1-C9-C8	6.07	120.64	109.70
2	A	401	DMU	O1-C9-C8	5.85	120.24	109.70
2	A	404	DMU	O1-C9-C8	5.73	120.03	109.70

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	DMU	O5-C4-C57	5.52	120.13	106.44
2	A	401	DMU	O1-C9-C11	5.52	120.11	106.44
2	A	404	DMU	O1-C9-C11	5.47	120.01	106.44
2	A	404	DMU	O16-C6-C1	5.46	116.57	108.27
2	A	403	DMU	O1-C9-C8	5.46	119.54	109.70
2	A	404	DMU	O5-C4-C57	5.41	119.84	106.44
2	A	402	DMU	O1-C10-C5	5.39	121.44	110.37
2	A	404	DMU	O5-C4-C3	5.25	120.57	109.72
2	A	403	DMU	C2-C3-C4	5.24	122.55	110.93
2	A	402	DMU	O5-C4-C3	5.23	120.53	109.72
2	A	404	DMU	O1-C10-C5	5.20	121.06	110.37
2	A	401	DMU	O5-C6-O16	5.11	122.13	110.04
2	A	403	DMU	O5-C4-C57	5.08	119.03	106.44
2	A	402	DMU	O5-C4-C57	5.07	119.00	106.44
2	A	401	DMU	O7-C10-C5	5.00	120.40	108.09
2	A	402	DMU	O1-C9-C11	4.98	118.79	106.44
2	A	404	DMU	C2-C3-C4	4.98	121.96	110.93
2	A	403	DMU	O5-C4-C3	4.96	119.98	109.72
2	A	402	DMU	O5-C6-O16	4.93	121.70	110.04
2	A	403	DMU	O7-C10-C5	4.77	119.81	108.09
2	A	401	DMU	O1-C10-C5	4.76	120.14	110.37
2	A	401	DMU	O5-C6-C1	4.66	119.95	110.37
2	A	404	DMU	O7-C10-C5	4.61	119.42	108.09
2	A	403	DMU	O7-C10-O1	4.52	122.59	110.69
2	A	404	DMU	O7-C3-C2	4.46	118.57	107.23
2	A	403	DMU	O1-C9-C11	4.35	117.23	106.44
2	A	402	DMU	O7-C10-C5	4.33	118.75	108.09
2	A	403	DMU	O16-C6-C1	4.21	114.66	108.27
2	A	403	DMU	O7-C3-C4	4.20	120.48	109.48
2	A	403	DMU	C11-C9-C8	4.13	123.17	113.02
2	A	402	DMU	O7-C3-C4	3.95	119.83	109.48
2	A	403	DMU	O5-C6-O16	3.89	119.23	110.04
2	A	404	DMU	O7-C3-C4	3.80	119.44	109.48
2	A	402	DMU	O16-C18-C19	3.73	122.01	109.37
2	A	401	DMU	C10-O7-C3	-3.71	109.19	117.98
2	A	401	DMU	O7-C3-C2	3.69	116.60	107.23
2	A	403	DMU	O7-C3-C2	3.67	116.55	107.23
2	A	404	DMU	O5-C6-O16	3.64	118.66	110.04
2	A	403	DMU	O1-C10-C5	3.52	117.59	110.37
2	A	402	DMU	O7-C10-O1	3.46	119.81	110.69
2	A	401	DMU	O5-C4-C3	3.44	116.84	109.72
2	A	401	DMU	C57-C4-C3	3.43	123.02	113.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	404	DMU	O7-C10-O1	3.35	119.52	110.69
2	A	403	DMU	C8-C7-C5	3.34	116.70	110.83
2	A	401	DMU	O7-C10-O1	3.31	119.39	110.69
2	A	404	DMU	C1-C2-C3	3.18	116.89	109.68
2	A	401	DMU	C6-C1-C2	-3.13	103.44	110.01
2	A	402	DMU	C11-C9-C8	3.07	120.56	113.02
2	A	404	DMU	C6-C1-C2	3.01	116.33	110.01
2	A	401	DMU	C2-C3-C4	2.98	117.53	110.93
2	A	404	DMU	O16-C18-C19	2.88	119.13	109.37
2	A	402	DMU	C2-C3-C4	2.86	117.27	110.93
2	A	404	DMU	C11-C9-C8	2.83	119.97	113.02
2	A	403	DMU	C10-O1-C9	-2.77	108.31	113.72
2	A	402	DMU	O5-C6-C1	2.75	116.03	110.37
2	A	401	DMU	C11-C9-C8	2.70	119.65	113.02
2	A	403	DMU	C57-C4-C3	2.68	120.91	113.38
2	A	402	DMU	C57-C4-C3	2.52	120.46	113.38
2	A	403	DMU	C6-C1-C2	2.40	115.05	110.01
2	A	402	DMU	C18-O16-C6	2.35	117.70	113.68
2	A	401	DMU	C22-C19-C18	-2.24	103.73	113.47
2	A	404	DMU	C57-C4-C3	2.21	119.59	113.38
2	A	401	DMU	O16-C18-C19	2.06	116.37	109.37

All (28) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	401	DMU	C4
2	A	401	DMU	C2
2	A	401	DMU	C3
2	A	401	DMU	C5
2	A	401	DMU	C9
2	A	401	DMU	C6
2	A	401	DMU	C10
2	A	402	DMU	C4
2	A	402	DMU	C2
2	A	402	DMU	C3
2	A	402	DMU	C5
2	A	402	DMU	C9
2	A	402	DMU	C6
2	A	402	DMU	C10
2	A	403	DMU	C4
2	A	403	DMU	C2
2	A	403	DMU	C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atom
2	A	403	DMU	C5
2	A	403	DMU	C9
2	A	403	DMU	C6
2	A	403	DMU	C10
2	A	404	DMU	C4
2	A	404	DMU	C2
2	A	404	DMU	C3
2	A	404	DMU	C5
2	A	404	DMU	C9
2	A	404	DMU	C6
2	A	404	DMU	C10

All (38) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	403	DMU	O5-C6-O16-C18
2	A	401	DMU	C2-C3-O7-C10
2	A	402	DMU	C2-C3-O7-C10
2	A	402	DMU	O6-C11-C9-O1
2	A	403	DMU	O6-C11-C9-O1
2	A	403	DMU	O5-C4-C57-O61
2	A	401	DMU	C3-C4-C57-O61
2	A	401	DMU	O6-C11-C9-C8
2	A	401	DMU	O5-C6-O16-C18
2	A	402	DMU	C3-C4-C57-O61
2	A	404	DMU	C3-C4-C57-O61
2	A	402	DMU	O16-C18-C19-C22
2	A	403	DMU	O16-C18-C19-C22
2	A	401	DMU	O16-C18-C19-C22
2	A	402	DMU	C1-C6-O16-C18
2	A	404	DMU	C1-C6-O16-C18
2	A	401	DMU	C4-C3-O7-C10
2	A	404	DMU	O6-C11-C9-O1
2	A	402	DMU	O5-C6-O16-C18
2	A	401	DMU	C18-C19-C22-C25
2	A	402	DMU	C19-C22-C25-C28
2	A	401	DMU	C19-C18-O16-C6
2	A	403	DMU	O1-C10-O7-C3
2	A	404	DMU	O6-C11-C9-C8
2	A	402	DMU	C5-C10-O7-C3
2	A	404	DMU	C25-C28-C31-C34
2	A	404	DMU	O16-C18-C19-C22

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	403	DMU	C34-C37-C40-C43
2	A	402	DMU	C25-C28-C31-C34
2	A	404	DMU	O1-C10-O7-C3
2	A	402	DMU	C19-C18-O16-C6
2	A	404	DMU	C2-C3-O7-C10
2	A	401	DMU	C5-C10-O7-C3
2	A	402	DMU	C28-C31-C34-C37
2	A	402	DMU	C18-C19-C22-C25
2	A	401	DMU	C22-C25-C28-C31
2	A	404	DMU	C19-C18-O16-C6
2	A	403	DMU	C25-C28-C31-C34


There are no ring outliers.


4 monomers are involved in 13 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	403	DMU	1	1
2	A	402	DMU	6	0
2	A	401	DMU	2	0
2	A	404	DMU	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	220/269 (81%)	0.29	18 (8%)	19 20	15, 30, 81, 93	4 (1%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	115	TYR	5.4
1	A	199	VAL	5.1
1	A	207	THR	3.8
1	A	211	THR	3.0
1	A	214	LEU	2.9
1	A	208	VAL	2.9
1	A	122	LEU	2.8
1	A	120	HIS	2.6
1	A	114	THR	2.6
1	A	13	PRO	2.6
1	A	210	ARG	2.6
1	A	217	ASN	2.4
1	A	195	LEU	2.4
1	A	113	PHE	2.4
1	A	226	TRP	2.3
1	A	212	HIS	2.3
1	A	182	PHE	2.2
1	A	117	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	DMU	A	401	33/33	0.69	0.19	28,74,86,87	0
2	DMU	A	402	33/33	0.71	0.20	49,78,85,86	0
2	DMU	A	404	33/33	0.78	0.15	47,81,89,90	0
2	DMU	A	403	33/33	0.80	0.18	24,51,70,71	0
4	MG	A	302	1/1	0.98	0.10	34,34,34,34	0
3	NA	A	301	1/1	0.99	0.03	25,25,25,25	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around DMU A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around DMU A 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around DMU A 404:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around DMU A 403:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.