



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 18, 2025 – 05:06 PM EDT

PDB ID : 9BBD / pdb\_00009bbd  
Title : Structure of S1\_8B, a lambda-carrageenan specific sulfatase  
Authors : Hettle, J.A.; Vickers, C.; Boraston, A.B.  
Deposited on : 2024-04-05  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.006 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.45.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	5963 (2.30-2.30)
Clashscore	180529	6698 (2.30-2.30)
Ramachandran outliers	177936	6640 (2.30-2.30)
Sidechain outliers	177891	6640 (2.30-2.30)
RSRZ outliers	164620	5963 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	541	<div> <div>%</div> <div> <div></div> <div>79%</div> <div>5%</div> <div>15%</div> </div> </div>
1	B	541	<div> <div>2%</div> <div> <div></div> <div>75%</div> <div>10%</div> <div>15%</div> </div> </div>

## 2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 7747 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called S1\_8B sulfatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	458	Total	C	N	O	S	0	0	0
			3649	2339	619	683	8			
1	B	459	Total	C	N	O	S	0	1	0
			3669	2351	623	686	9			

- Molecule 2 is ACETATE ION (CCD ID: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	4	Total Cl 4 4	0	0
3	B	3	Total Cl 3 3	0	0

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula:  $O_4S$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		
6	B	1	Total	Ca	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	209	Total 209	O 209	0	0
7	B	147	Total 147	O 147	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: S1\_8B sulfatase



#### • Molecule 1: S1\_8B sulfatase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	93.84Å 103.09Å 148.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.93 – 2.30 29.93 – 2.30	Depositor EDS
% Data completeness (in resolution range)	93.8 (29.93-2.30) 94.0 (29.93-2.30)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.19 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.15.2 _3472	Depositor
R, $R_{free}$	0.203 , 0.235 0.204 , 0.235	Depositor DCC
$R_{free}$ test set	3117 reflections (4.81%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	34.9	Xtriage
Anisotropy	0.091	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 32.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	7747	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, SO4, CA, EDO, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.37	0/3754	0.55	0/5110
1	B	0.34	0/3774	0.51	0/5135
All	All	0.35	0/7528	0.53	0/10245

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3649	0	3443	19	0
1	B	3669	0	3464	38	0
2	A	4	0	3	0	0
2	B	4	0	3	1	0
3	A	4	0	0	0	0
3	B	3	0	0	0	0
4	A	12	0	18	4	0
4	B	4	0	6	0	0
5	A	20	0	0	0	0
5	B	20	0	0	1	0
6	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	1	0	0	0	0
7	A	209	0	0	1	0
7	B	147	0	0	1	0
All	All	7747	0	6937	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:354:LYS:NZ	1:A:362:ASP:OD2	2.00	0.94
1:A:218:THR:H	4:A:607:EDO:H12	1.37	0.89
1:B:274:LEU:HD11	1:B:357[A]:MET:HE1	1.63	0.81
1:B:354:LYS:HE3	1:B:357[A]:MET:HE3	1.78	0.66
1:B:354:LYS:CE	1:B:357[A]:MET:HE3	2.28	0.63
1:B:259:ILE:HG22	1:B:259:ILE:O	1.99	0.61
1:A:91:VAL:HG11	1:A:357:MET:SD	2.42	0.59
1:A:218:THR:H	4:A:607:EDO:C1	2.12	0.58
1:A:86:HIS:HB2	1:A:350:VAL:HG11	1.89	0.55
1:B:47:ILE:HG21	1:B:322:SER:HB3	1.90	0.54
1:B:118:ALA:HB2	1:B:533:ILE:HD11	1.92	0.52
1:A:144:PHE:CZ	1:A:192:GLY:HA3	2.44	0.52
1:B:132:VAL:HG13	1:B:133:PRO:HD3	1.90	0.52
1:B:274:LEU:CD1	1:B:357[A]:MET:HE1	2.37	0.51
1:B:354:LYS:NZ	1:B:362:ASP:OD2	2.41	0.51
1:B:64:ASP:OD1	1:B:66:THR:OG1	2.28	0.50
1:B:295:TRP:CE3	1:B:298:GLY:HA3	2.46	0.50
1:B:143:THR:OG1	1:B:161:TYR:HA	2.12	0.49
1:B:52:MET:HE1	1:B:267:TRP:CE2	2.47	0.48
1:A:447:GLU:OE1	1:A:450:ARG:NH1	2.46	0.48
1:B:144:PHE:CZ	1:B:192:GLY:HA3	2.48	0.48
1:B:526:TYR:OH	5:B:608:SO4:O2	2.23	0.48
1:A:171:GLN:HG3	7:A:848:HOH:O	2.12	0.48
1:A:115:SER:O	1:A:117:ARG:HD3	2.13	0.47
1:B:108:LEU:HD12	7:B:810:HOH:O	2.14	0.47
1:B:305:LYS:NZ	1:B:344:PHE:O	2.47	0.47
1:B:314:SER:HB2	1:B:350:VAL:HG13	1.96	0.47
1:A:52:MET:HE1	1:A:267:TRP:CE2	2.51	0.46
1:B:113:HIS:HA	1:B:114:ARG:HA	1.71	0.46
1:A:44:PRO:HD2	1:A:325:LEU:O	2.16	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:57:PRO:HB3	1:B:290:PRO:HB3	1.97	0.46
1:B:186:LYS:HA	1:B:259:ILE:HD11	1.97	0.46
1:B:217:MET:O	1:B:230:ARG:NH1	2.48	0.46
1:B:371:LYS:HE2	1:B:429:GLN:OE1	2.16	0.45
1:B:296:PRO:HA	1:B:299:LYS:HG3	1.97	0.45
1:B:78:GLN:HE22	1:B:294:SER:CB	2.30	0.45
1:B:354:LYS:HE2	1:B:357[A]:MET:HE3	2.00	0.44
1:B:501:SER:OG	2:B:601:ACT:H3	2.17	0.44
1:B:254:LEU:HD23	1:B:259:ILE:CG2	2.47	0.44
1:B:87:GLN:OE1	1:B:315:GLY:HA3	2.18	0.44
1:A:52:MET:O	1:A:269:SER:HA	2.18	0.44
1:B:259:ILE:HG22	1:B:263:THR:OG1	2.17	0.43
1:B:91:VAL:HG11	1:B:357[A]:MET:SD	2.58	0.43
1:A:254:LEU:HD22	1:A:259:ILE:HG22	2.00	0.43
1:A:306:GLY:HA2	4:A:606:EDO:H12	2.00	0.43
1:A:172:GLY:H	4:A:608:EDO:H22	1.83	0.42
1:A:357:MET:HB3	1:A:357:MET:HE2	1.52	0.42
1:B:378:TYR:C	1:B:380:PRO:HD3	2.44	0.42
1:A:87:GLN:OE1	1:A:315:GLY:HA3	2.20	0.42
1:A:113:HIS:HA	1:A:114:ARG:HA	1.76	0.41
1:B:132:VAL:CG1	1:B:133:PRO:HD3	2.49	0.41
1:B:259:ILE:HD13	1:B:259:ILE:HG21	1.84	0.41
1:B:311:ASP:OD1	1:B:311:ASP:N	2.47	0.41
1:B:477:GLU:HG3	1:B:533:ILE:CD1	2.50	0.41
1:B:376:ARG:HD2	1:B:455:LEU:HD23	2.03	0.40
1:B:145:ASN:O	1:B:176:GLY:HA3	2.21	0.40
1:A:501:SER:O	1:A:502:VAL:C	2.65	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	454/541 (84%)	442 (97%)	11 (2%)	1 (0%)	44	55
1	B	456/541 (84%)	441 (97%)	15 (3%)	0	100	100
All	All	910/1082 (84%)	883 (97%)	26 (3%)	1 (0%)	48	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	464	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	374/471 (79%)	373 (100%)	1 (0%)	91	96
1	B	376/471 (80%)	376 (100%)	0	100	100
All	All	750/942 (80%)	749 (100%)	1 (0%)	92	97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	354	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	71	ASN
1	A	193	GLN
1	A	503	ASN
1	B	71	ASN
1	B	78	GLN
1	B	203	ASN
1	B	310	ASN
1	B	503	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 9 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	EDO	A	606	-	3,3,3	0.46	0	2,2,2	0.35	0
4	EDO	B	605	-	3,3,3	0.51	0	2,2,2	0.20	0
2	ACT	A	601	-	3,3,3	1.18	0	3,3,3	1.77	1 (33%)
5	SO4	A	611	-	4,4,4	0.35	0	6,6,6	0.30	0
4	EDO	A	607	-	3,3,3	0.48	0	2,2,2	0.22	0
2	ACT	B	601	-	3,3,3	1.52	1 (33%)	3,3,3	1.36	0
5	SO4	A	612	-	4,4,4	0.28	0	6,6,6	0.25	0
5	SO4	A	610	-	4,4,4	0.26	0	6,6,6	0.14	0
5	SO4	B	606	-	4,4,4	0.25	0	6,6,6	0.16	0
4	EDO	A	608	-	3,3,3	0.51	0	2,2,2	0.32	0
5	SO4	B	609	-	4,4,4	0.29	0	6,6,6	0.25	0
5	SO4	B	607	-	4,4,4	0.25	0	6,6,6	0.24	0
5	SO4	A	609	-	4,4,4	0.25	0	6,6,6	0.17	0
5	SO4	B	608	-	4,4,4	0.21	0	6,6,6	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	606	-	-	0/1/1/1	-
4	EDO	B	605	-	-	0/1/1/1	-
4	EDO	A	607	-	-	0/1/1/1	-
4	EDO	A	608	-	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	ACT	CH3-C	2.25	1.57	1.49

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	ACT	OXT-C-O	2.58	131.61	122.03

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	608	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	606	EDO	1	0
4	A	607	EDO	2	0
2	B	601	ACT	1	0
4	A	608	EDO	1	0
5	B	608	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	458/541 (84%)	-0.33	4 (0%) 81 81	20, 32, 46, 67	0
1	B	459/541 (84%)	-0.04	10 (2%) 62 63	21, 38, 57, 77	2 (0%)
All	All	917/1082 (84%)	-0.19	14 (1%) 71 72	20, 35, 54, 77	2 (0%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	421	PRO	5.9
1	A	539	GLN	3.9
1	B	259	ILE	3.2
1	B	539	GLN	3.0
1	B	42	LYS	2.8
1	B	496	LEU	2.8
1	A	205	ASN	2.5
1	B	206	GLY	2.3
1	B	528	SER	2.2
1	B	380	PRO	2.2
1	B	170	PHE	2.2
1	B	204	PHE	2.2
1	A	357	MET	2.1
1	B	502	VAL	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ACT	B	601	4/4	0.57	0.27	53,59,63,63	0
3	CL	B	604	1/1	0.82	0.19	74,74,74,74	0
4	EDO	B	605	4/4	0.82	0.16	43,47,52,52	0
3	CL	B	603	1/1	0.83	0.14	56,56,56,56	0
4	EDO	A	606	4/4	0.84	0.15	43,44,49,50	0
4	EDO	A	607	4/4	0.85	0.14	34,35,46,48	0
3	CL	A	602	1/1	0.85	0.17	72,72,72,72	0
5	SO4	B	609	5/5	0.88	0.12	55,61,66,74	0
5	SO4	A	609	5/5	0.89	0.17	45,50,69,70	0
2	ACT	A	601	4/4	0.89	0.17	41,42,43,50	0
3	CL	A	605	1/1	0.91	0.16	62,62,62,62	0
5	SO4	B	606	5/5	0.91	0.14	58,65,72,75	0
3	CL	A	604	1/1	0.91	0.12	59,59,59,59	0
5	SO4	A	612	5/5	0.94	0.09	48,49,57,62	0
5	SO4	B	607	5/5	0.95	0.08	48,56,60,71	0
5	SO4	B	608	5/5	0.95	0.11	48,50,56,59	0
4	EDO	A	608	4/4	0.95	0.08	36,37,39,43	0
3	CL	A	603	1/1	0.96	0.14	52,52,52,52	0
6	CA	A	613	1/1	0.96	0.08	19,19,19,19	0
3	CL	B	602	1/1	0.97	0.18	66,66,66,66	0
5	SO4	A	610	5/5	0.97	0.06	48,49,51,61	0
5	SO4	A	611	5/5	0.97	0.09	35,38,45,46	0
6	CA	B	610	1/1	0.98	0.10	26,26,26,26	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.