



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 17, 2024 – 06:08 pm BST

PDB ID : 8BGL  
Title : Structure of the dimeric rsCherryRev1.4  
Authors : Bui, T.Y.H.; Van Meervelt, L.  
Deposited on : 2022-10-28  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	269	 74% 7% 19%
1	B	269	 74% 7% 19%

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3792 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PAmCherry1 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	216	Total	C	N	O	S	0	2	0
			1745	1117	286	330	12			
1	B	216	Total	C	N	O	S	0	3	0
			1743	1116	285	330	12			

There are 106 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-37	MET	-	initiating methionine	UNP D1MPT3
A	-36	ARG	-	expression tag	UNP D1MPT3
A	-35	GLY	-	expression tag	UNP D1MPT3
A	-34	SER	-	expression tag	UNP D1MPT3
A	-33	HIS	-	expression tag	UNP D1MPT3
A	-32	HIS	-	expression tag	UNP D1MPT3
A	-31	HIS	-	expression tag	UNP D1MPT3
A	-30	HIS	-	expression tag	UNP D1MPT3
A	-29	HIS	-	expression tag	UNP D1MPT3
A	-28	HIS	-	expression tag	UNP D1MPT3
A	-27	GLY	-	expression tag	UNP D1MPT3
A	-26	MET	-	expression tag	UNP D1MPT3
A	-25	ALA	-	expression tag	UNP D1MPT3
A	-24	SER	-	expression tag	UNP D1MPT3
A	-23	MET	-	expression tag	UNP D1MPT3
A	-22	THR	-	expression tag	UNP D1MPT3
A	-21	GLY	-	expression tag	UNP D1MPT3
A	-20	GLY	-	expression tag	UNP D1MPT3
A	-19	GLN	-	expression tag	UNP D1MPT3
A	-18	GLN	-	expression tag	UNP D1MPT3
A	-17	MET	-	expression tag	UNP D1MPT3
A	-16	GLY	-	expression tag	UNP D1MPT3
A	-15	ARG	-	expression tag	UNP D1MPT3
A	-14	ASP	-	expression tag	UNP D1MPT3
A	-13	LEU	-	expression tag	UNP D1MPT3

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	TYR	-	expression tag	UNP D1MPT3
A	-11	ASP	-	expression tag	UNP D1MPT3
A	-10	ASP	-	expression tag	UNP D1MPT3
A	-9	ASP	-	expression tag	UNP D1MPT3
A	-8	ASP	-	expression tag	UNP D1MPT3
A	-7	LYS	-	expression tag	UNP D1MPT3
A	-6	ASP	-	expression tag	UNP D1MPT3
A	-5	PRO	-	expression tag	UNP D1MPT3
A	24	CYS	GLY	conflict	UNP D1MPT3
A	26	GLU	VAL	conflict	UNP D1MPT3
A	36	HIS	ARG	conflict	UNP D1MPT3
A	57	ALA	THR	conflict	UNP D1MPT3
A	66	QYX	MET	chromophore	UNP D1MPT3
A	66	Q2K	TYR	chromophore	UNP D1MPT3
A	66	QYX	GLY	chromophore	UNP D1MPT3
A	70	LYS	ASN	conflict	UNP D1MPT3
A	83	LEU	PHE	conflict	UNP D1MPT3
A	98	ASN	LYS	conflict	UNP D1MPT3
A	125	CYS	ARG	conflict	UNP D1MPT3
A	144	PHE	GLU	conflict	UNP D1MPT3
A	146	CYS	LEU	conflict	UNP D1MPT3
A	149	GLN	ARG	conflict	UNP D1MPT3
A	160	LEU	GLU	conflict	UNP D1MPT3
A	161	SER	VAL	conflict	UNP D1MPT3
A	163	MET	PRO	conflict	UNP D1MPT3
A	165	LEU	VAL	conflict	UNP D1MPT3
A	177	PHE	VAL	conflict	UNP D1MPT3
A	197	ILE	ARG	conflict	UNP D1MPT3
B	-37	MET	-	initiating methionine	UNP D1MPT3
B	-36	ARG	-	expression tag	UNP D1MPT3
B	-35	GLY	-	expression tag	UNP D1MPT3
B	-34	SER	-	expression tag	UNP D1MPT3
B	-33	HIS	-	expression tag	UNP D1MPT3
B	-32	HIS	-	expression tag	UNP D1MPT3
B	-31	HIS	-	expression tag	UNP D1MPT3
B	-30	HIS	-	expression tag	UNP D1MPT3
B	-29	HIS	-	expression tag	UNP D1MPT3
B	-28	HIS	-	expression tag	UNP D1MPT3
B	-27	GLY	-	expression tag	UNP D1MPT3
B	-26	MET	-	expression tag	UNP D1MPT3
B	-25	ALA	-	expression tag	UNP D1MPT3
B	-24	SER	-	expression tag	UNP D1MPT3

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	-23	MET	-	expression tag	UNP D1MPT3
B	-22	THR	-	expression tag	UNP D1MPT3
B	-21	GLY	-	expression tag	UNP D1MPT3
B	-20	GLY	-	expression tag	UNP D1MPT3
B	-19	GLN	-	expression tag	UNP D1MPT3
B	-18	GLN	-	expression tag	UNP D1MPT3
B	-17	MET	-	expression tag	UNP D1MPT3
B	-16	GLY	-	expression tag	UNP D1MPT3
B	-15	ARG	-	expression tag	UNP D1MPT3
B	-14	ASP	-	expression tag	UNP D1MPT3
B	-13	LEU	-	expression tag	UNP D1MPT3
B	-12	TYR	-	expression tag	UNP D1MPT3
B	-11	ASP	-	expression tag	UNP D1MPT3
B	-10	ASP	-	expression tag	UNP D1MPT3
B	-9	ASP	-	expression tag	UNP D1MPT3
B	-8	ASP	-	expression tag	UNP D1MPT3
B	-7	LYS	-	expression tag	UNP D1MPT3
B	-6	ASP	-	expression tag	UNP D1MPT3
B	-5	PRO	-	expression tag	UNP D1MPT3
B	24	CYS	GLY	conflict	UNP D1MPT3
B	26	GLU	VAL	conflict	UNP D1MPT3
B	36	HIS	ARG	conflict	UNP D1MPT3
B	57	ALA	THR	conflict	UNP D1MPT3
B	66	QYX	MET	chromophore	UNP D1MPT3
B	66	Q2K	TYR	chromophore	UNP D1MPT3
B	66	QYX	GLY	chromophore	UNP D1MPT3
B	70	LYS	ASN	conflict	UNP D1MPT3
B	83	LEU	PHE	conflict	UNP D1MPT3
B	98	ASN	LYS	conflict	UNP D1MPT3
B	125	CYS	ARG	conflict	UNP D1MPT3
B	144	PHE	GLU	conflict	UNP D1MPT3
B	146	CYS	LEU	conflict	UNP D1MPT3
B	149	GLN	ARG	conflict	UNP D1MPT3
B	160	LEU	GLU	conflict	UNP D1MPT3
B	161	SER	VAL	conflict	UNP D1MPT3
B	163	MET	PRO	conflict	UNP D1MPT3
B	165	LEU	VAL	conflict	UNP D1MPT3
B	177	PHE	VAL	conflict	UNP D1MPT3
B	197	ILE	ARG	conflict	UNP D1MPT3

- Molecule 2 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula:  $C_{10}H_{22}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			16	10	6		
3	B	1	Total	C	O	0	0
			16	10	6		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	133	Total	O	0	3
			133	133		
5	B	128	Total	O	0	2
			128	128		

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PAmCherry1 protein

Chain A: 



- Molecule 1: PAmCherry1 protein

Chain B: 





## 4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.10Å 48.82Å 99.12Å 90.00° 101.16° 90.00°	Depositor
Resolution (Å)	62.62 – 2.00 76.18 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (62.62-2.00) 97.1 (76.18-2.00)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.72 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, $R_{free}$	0.188 , 0.231 0.188 , 0.231	Depositor DCC
$R_{free}$ test set	1543 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.6	Xtriage
Anisotropy	0.561	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 32.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.209 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3792	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.81% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 1PE, Q2K, QYX, PEG, EDO, ABA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.28	0/1744	0.53	0/2351
1	B	0.28	0/1742	0.52	0/2349
All	All	0.28	0/3486	0.53	0/4700

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	215[B]	ABA	Mainchain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1745	0	1626	10	0
1	B	1743	0	1621	10	0
2	A	7	0	10	0	0
3	A	16	0	22	0	0
3	B	16	0	22	1	0
4	B	4	0	6	0	0
5	A	133	0	0	2	0
5	B	128	0	0	1	0
All	All	3792	0	3307	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:125:CYS:SG	5:B:502:HOH:O	2.45	0.74
1:B:178:LYS:HB3	3:B:301:1PE:H161	1.70	0.73
1:B:28:GLU:HB2	1:B:50:LYS:HB2	1.92	0.51
1:B:14:PHE:CZ	1:B:33:GLY:HA3	2.48	0.49
1:A:198:LYS:NZ	1:A:200:ASP:OD1	2.45	0.49
1:A:14:PHE:CZ	1:A:33:GLY:HA3	2.48	0.49
1:A:59:ASP:HB3	1:A:165:LEU:HD21	1.95	0.48
1:B:14:PHE:HB3	1:B:118:PHE:HB2	1.98	0.46
1:B:59:ASP:HB3	1:B:165:LEU:HD21	1.99	0.44
1:A:200:ASP:OD2	5:A:601:HOH:O	2.21	0.43
1:A:164:ARG:HG2	1:A:174:ASP:OD1	2.20	0.42
1:A:28:GLU:HB2	1:A:50:LYS:HB2	2.01	0.42
1:A:141:MET:SD	1:A:168:LYS:HA	2.59	0.42
1:B:99:PHE:CD2	1:B:175:ALA:HB2	2.55	0.41
1:A:178:LYS:NZ	5:A:609:HOH:O	2.48	0.41
1:B:20:GLY:HA3	1:B:124:LEU:O	2.21	0.41
1:B:141:MET:SD	1:B:168:LYS:HA	2.60	0.41
1:A:143:TRP:CZ3	1:A:165:LEU:HG	2.55	0.41
1:A:20:GLY:HA3	1:A:124:LEU:O	2.20	0.41
1:B:7:ILE:HD13	1:B:88:PRO:HD3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	210/269 (78%)	208 (99%)	2 (1%)	0	100	100
1	B	210/269 (78%)	207 (99%)	3 (1%)	0	100	100
All	All	420/538 (78%)	415 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	179/226 (79%)	179 (100%)	0	100	100
1	B	179/226 (79%)	179 (100%)	0	100	100
All	All	358/452 (79%)	358 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	QYX	B	66[A]	1	22,24,25	3.23	5 (22%)	16,32,34	2.65	4 (25%)
1	ABA	A	215[B]	1	4,5,6	0.87	0	1,5,7	0.23	0
1	Q2K	B	66[B]	1	15,16,17	1.14	1 (6%)	8,21,23	1.34	0
1	QYX	A	66[A]	1	22,24,25	3.29	6 (27%)	16,32,34	2.71	4 (25%)
1	Q2K	A	66[B]	1	15,16,17	1.16	1 (6%)	8,21,23	1.33	0
1	ABA	B	215[B]	1	4,5,6	0.83	0	1,5,7	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	QYX	B	66[A]	1	-	4/9/31/32	0/2/2/2
1	ABA	A	215[B]	1	-	0/3/4/6	-
1	Q2K	B	66[B]	1	-	3/5/27/28	0/1/1/1
1	QYX	A	66[A]	1	-	5/9/31/32	0/2/2/2
1	Q2K	A	66[B]	1	-	1/5/27/28	0/1/1/1
1	ABA	B	215[B]	1	-	0/3/4/6	-

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66[A]	QYX	CB2-CA2	-13.36	1.36	1.50
1	B	66[A]	QYX	CB2-CA2	-13.17	1.36	1.50
1	A	66[A]	QYX	C1-N2	-3.98	1.29	1.35
1	B	66[A]	QYX	C1-N2	-3.93	1.29	1.35
1	A	66[A]	QYX	C1-N3	3.42	1.40	1.37
1	B	66[A]	QYX	CA2-N2	3.21	1.40	1.35
1	A	66[A]	QYX	CA2-N2	3.17	1.40	1.35

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	66[A]	QYX	CB2-CG2	-3.16	1.46	1.51
1	B	66[A]	QYX	C1-N3	3.08	1.40	1.37
1	B	66[A]	QYX	CB2-CG2	-3.06	1.46	1.51
1	A	66[B]	Q2K	C1-N3	2.87	1.43	1.38
1	B	66[B]	Q2K	C1-N3	2.71	1.42	1.38
1	A	66[A]	QYX	CA1-N1	2.07	1.32	1.27

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	66[A]	QYX	CG2-CB2-CA2	8.49	133.80	113.58
1	B	66[A]	QYX	CG2-CB2-CA2	8.10	132.87	113.58
1	B	66[A]	QYX	O2-C2-CA2	5.87	132.60	120.67
1	A	66[A]	QYX	O2-C2-CA2	5.74	132.35	120.67
1	A	66[A]	QYX	CB2-CA2-C2	-2.10	125.50	129.44
1	B	66[A]	QYX	O3-C3-CA3	-2.09	120.36	126.32
1	A	66[A]	QYX	O3-C3-CA3	-2.07	120.43	126.32
1	B	66[A]	QYX	CB2-CA2-C2	-2.01	125.67	129.44

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	66[A]	QYX	CA1-CB1-CG1-SD
1	A	66[A]	QYX	N2-CA2-CB2-CG2
1	A	66[A]	QYX	C3-CA3-N3-C2
1	B	66[A]	QYX	CA1-CB1-CG1-SD
1	B	66[A]	QYX	C1-CA1-CB1-CG1
1	B	66[A]	QYX	C2-CA2-CB2-CG2
1	A	66[A]	QYX	CB1-CG1-SD-CE
1	B	66[B]	Q2K	CB1-CG1-SD-CE
1	A	66[A]	QYX	C2-CA2-CB2-CG2
1	B	66[B]	Q2K	CA1-CB1-CG1-SD
1	B	66[A]	QYX	N2-CA2-CB2-CG2
1	A	66[B]	Q2K	C3-CA3-N3-C1
1	B	66[B]	Q2K	C3-CA3-N3-C1

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	EDO	B	302	-	3,3,3	0.49	0	2,2,2	0.33	0
3	1PE	B	301	-	15,15,15	0.14	0	14,14,14	0.16	0
3	1PE	A	302	-	15,15,15	0.14	0	14,14,14	0.06	0
2	PEG	A	301	-	6,6,6	0.12	0	5,5,5	0.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	302	-	-	0/1/1/1	-
3	1PE	B	301	-	-	7/13/13/13	-
3	1PE	A	302	-	-	9/13/13/13	-
2	PEG	A	301	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	302	1PE	OH6-C15-C25-OH5
3	B	301	1PE	OH5-C14-C24-OH4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	B	301	1PE	OH4-C13-C23-OH3
3	A	302	1PE	OH5-C14-C24-OH4
3	B	301	1PE	OH2-C12-C22-OH3
3	A	302	1PE	OH4-C13-C23-OH3
3	B	301	1PE	C14-C24-OH4-C13
3	A	302	1PE	OH2-C12-C22-OH3
3	A	302	1PE	OH7-C16-C26-OH6
3	B	301	1PE	C25-C15-OH6-C26
3	A	302	1PE	C25-C15-OH6-C26
3	B	301	1PE	C12-C22-OH3-C23
3	A	302	1PE	C14-C24-OH4-C13
3	A	302	1PE	C13-C23-OH3-C22
3	B	301	1PE	C23-C13-OH4-C24
3	A	302	1PE	C24-C14-OH5-C25

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	301	1PE	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	214/269 (79%)	-0.47	0 100 100	25, 34, 46, 56	0
1	B	214/269 (79%)	-0.41	0 100 100	24, 35, 52, 61	0
All	All	428/538 (79%)	-0.44	0 100 100	24, 34, 49, 61	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	QYX	B	66[A]	23/24	0.90	0.16	35,42,45,46	23
1	Q2K	B	66[B]	16/17	0.90	0.15	34,41,44,45	16
1	Q2K	A	66[B]	16/17	0.91	0.13	36,40,44,45	16
1	QYX	A	66[A]	23/24	0.92	0.14	35,41,42,43	23
1	ABA	B	215[B]	6/7	0.96	0.11	27,28,30,32	6
1	ABA	A	215[B]	6/7	0.97	0.10	26,29,30,32	6

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	1PE	A	302	16/16	0.76	0.20	27,37,42,44	16
4	EDO	B	302	4/4	0.86	0.12	41,45,45,55	0
3	1PE	B	301	16/16	0.88	0.13	35,42,57,57	0
2	PEG	A	301	7/7	0.89	0.18	46,47,49,49	7

## 6.5 Other polymers [i](#)

There are no such residues in this entry.