



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 10, 2024 – 12:58 pm GMT

PDB ID : 2BHG  
Title : 3C protease from type A10(61) foot-and-mouth disease virus  
Authors : Birtley, J.R.; Brick, P.; Curry, S.  
Deposited on : 2005-01-10  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	<div> <div>3%</div> <div>73%</div> <div>19%</div> <div>• 7%</div> </div>
1	B	209	<div> <div>8%</div> <div>72%</div> <div>16%</div> <div>12%</div> </div>

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2830 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FOOT-AND-MOUTH DISEASE VIRUS 3C PROTEASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	194	Total	C	N	O	S	Se	0	0	0
			1383	875	234	261	4	9			
1	B	184	Total	C	N	O	S	Se	0	0	0
			1349	854	236	246	4	9			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	95	LYS	CYS	engineered mutation	UNP P03306
A	163	ALA	CYS	engineered mutation	UNP P03306
B	95	LYS	CYS	engineered mutation	UNP P03306
B	163	ALA	CYS	engineered mutation	UNP P03306
A	78	GLY	ARG	conflict	UNP P03306
A	79	GLN	THR	conflict	UNP P03306
A	80	ASP	GLY	conflict	UNP P03306
A	81	MSE	HIS	conflict	UNP P03306
A	82	LEU	ALA	conflict	UNP P03306
A	83	SER	LEU	conflict	UNP P03306
A	84	ASP	ARG	conflict	UNP P03306
A	85	ALA	ARG	conflict	UNP P03306
A	86	ALA	GLY	conflict	UNP P03306
A	87	LEU	THR	conflict	UNP P03306
A	88	MSE	HIS	conflict	UNP P03306
A	89	VAL	TRP	conflict	UNP P03306
B	82	LEU	ALA	conflict	UNP P03306
B	83	SER	LEU	conflict	UNP P03306
B	84	ASP	ARG	conflict	UNP P03306
B	85	ALA	ARG	conflict	UNP P03306
B	86	ALA	GLY	conflict	UNP P03306
B	87	LEU	THR	conflict	UNP P03306
B	88	MSE	HIS	conflict	UNP P03306
B	89	VAL	TRP	conflict	UNP P03306


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	54	Total 54	O 54	0	0
2	B	44	Total 44	O 44	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: FOOT-AND-MOUTH DISEASE VIRUS 3C PROTEASE



#### • Molecule 1: FOOT-AND-MOUTH DISEASE VIRUS 3C PROTEASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	141.59Å 141.59Å 43.68Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	26.76 – 1.90 26.76 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (26.76-1.90) 100.0 (26.76-1.90)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.67 (at 1.89Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.218 , 0.245 0.215 , 0.242	Depositor DCC
$R_{free}$ test set	1310 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.7	Xtriage
Anisotropy	0.474	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 47.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.059 for h,-h-k,-l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	2830	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.35	0/1395	0.66	0/1875
1	B	0.35	0/1362	0.64	0/1822
All	All	0.35	0/2757	0.65	0/3697

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1383	0	1310	40	0
1	B	1349	0	1323	39	0
2	A	54	0	0	0	0
2	B	44	0	0	2	0
All	All	2830	0	2633	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:148:MSE:HE3	1:B:186:ASN:HB2	1.46	0.95

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:54:LYS:HE3	1:A:61:ALA:HB1	1.55	0.87
1:A:54:LYS:HE2	1:A:56:MSE:SE	2.28	0.83
1:A:147:THR:O	1:A:149:PRO:HD3	1.84	0.78
1:B:57:LEU:HB3	1:B:62:MSE:HE3	1.69	0.75
1:B:138:ASP:OD1	1:B:140:VAL:HG22	1.86	0.74
1:B:21:LEU:HD22	1:B:55:ILE:HD12	1.67	0.73
1:A:148:MSE:CE	1:B:186:ASN:HB2	2.21	0.70
1:B:106:THR:HG22	1:B:140:VAL:CG1	2.20	0.70
1:A:75:LYS:HA	1:A:80:ASP:OD1	1.93	0.68
1:A:8:LEU:HD12	1:A:11:MSE:HE2	1.74	0.68
1:B:92:ARG:HH11	1:B:92:ARG:HG3	1.58	0.68
1:A:8:LEU:HD12	1:A:11:MSE:CE	2.25	0.67
1:A:147:THR:C	1:A:149:PRO:HD3	2.15	0.66
1:A:134:LEU:HD21	1:B:134:LEU:HD21	1.79	0.65
1:B:21:LEU:CD2	1:B:55:ILE:HD12	2.28	0.63
1:A:138:ASP:O	1:A:140:VAL:HG23	2.03	0.59
1:B:62:MSE:HB2	1:B:66:ASP:OD2	2.03	0.58
1:A:76:VAL:HG12	1:A:77:LYS:N	2.19	0.58
1:B:134:LEU:HD13	1:B:155:LYS:HG2	1.86	0.57
1:B:138:ASP:OD1	1:B:140:VAL:CG2	2.50	0.57
1:A:147:THR:O	1:A:147:THR:HG22	2.06	0.56
1:A:116:VAL:HG13	1:A:128:ILE:HG23	1.87	0.56
1:B:126:ARG:NH1	2:B:2027:HOH:O	2.38	0.56
1:B:54:LYS:O	1:B:55:ILE:HD13	2.07	0.54
1:B:72:PHE:O	1:B:82:LEU:HD12	2.07	0.54
1:A:140:VAL:HG12	1:A:140:VAL:O	2.08	0.53
1:A:127:LEU:C	1:A:127:LEU:HD23	2.29	0.53
1:B:107:ALA:HB3	1:B:139:ILE:HG12	1.90	0.52
1:A:106:THR:HG22	1:A:141:VAL:HG22	1.92	0.51
1:B:137:LYS:NZ	1:B:148:MSE:O	2.43	0.51
1:B:148:MSE:O	1:B:148:MSE:HG2	2.10	0.51
1:B:104:ARG:HD3	1:B:196:ARG:HH21	1.76	0.51
1:B:92:ARG:HG3	1:B:92:ARG:NH1	2.26	0.50
1:A:45:ARG:NE	1:A:71:GLU:OE2	2.44	0.50
1:A:8:LEU:CD2	1:A:175:THR:HG22	2.43	0.49
1:A:55:ILE:O	1:A:55:ILE:HG13	2.07	0.49
1:A:137:LYS:NZ	1:A:148:MSE:O	2.45	0.49
1:A:116:VAL:HG13	1:A:128:ILE:CG2	2.43	0.49
1:B:60:ARG:HD3	1:B:62:MSE:CE	2.43	0.48
1:A:54:LYS:HE3	1:A:61:ALA:CB	2.36	0.48
1:B:60:ARG:HD3	1:B:62:MSE:HE1	1.97	0.47

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:151:LEU:HD22	1:A:190:TYR:CE1	2.51	0.46
1:B:60:ARG:CD	1:B:62:MSE:HE2	2.46	0.46
1:B:148:MSE:N	1:B:149:PRO:CD	2.79	0.46
1:B:60:ARG:CG	1:B:62:MSE:HE2	2.46	0.46
1:A:117:GLY:HA3	1:A:129:PHE:CE2	2.51	0.45
1:B:72:PHE:CE2	1:B:202:MSE:HE2	2.51	0.45
1:B:48:PHE:C	1:B:50:GLU:H	2.20	0.45
1:A:8:LEU:HD23	1:A:175:THR:HG22	1.99	0.44
1:B:57:LEU:HB3	1:B:62:MSE:CE	2.44	0.44
1:B:72:PHE:CG	1:B:198:MSE:HE3	2.53	0.44
1:A:117:GLY:HA3	1:A:129:PHE:CZ	2.53	0.44
1:A:8:LEU:HD23	1:A:168:LEU:HD13	1.98	0.44
1:A:141:VAL:HG12	1:A:141:VAL:O	2.16	0.44
1:A:171:ASP:HB2	1:A:176:PHE:CE2	2.53	0.44
1:A:55:ILE:HD11	1:A:62:MSE:CE	2.48	0.44
1:A:76:VAL:CG1	1:A:77:LYS:N	2.82	0.43
1:A:123:ASP:OD2	1:A:159:ARG:NH1	2.51	0.43
1:B:196:ARG:O	1:B:200:GLN:HG3	2.18	0.43
1:A:11:MSE:HE1	1:A:102:HIS:CD2	2.54	0.43
1:A:148:MSE:HE2	1:B:185:GLY:O	2.18	0.43
1:A:131:GLY:HA3	1:A:155:LYS:O	2.19	0.42
1:A:51:LYS:HD3	1:A:51:LYS:HA	1.88	0.42
1:B:117:GLY:HA3	1:B:129:PHE:CE2	2.54	0.42
1:B:106:THR:HG22	1:B:140:VAL:HG12	2.00	0.42
1:B:149:PRO:HA	2:B:2033:HOH:O	2.18	0.42
1:B:117:GLY:HA3	1:B:129:PHE:CZ	2.55	0.42
1:A:23:LEU:HD21	1:A:52:TYR:HA	2.02	0.41
1:A:140:VAL:O	1:A:141:VAL:C	2.58	0.41
1:B:72:PHE:CD2	1:B:198:MSE:HE3	2.55	0.41
1:B:106:THR:HG22	1:B:140:VAL:HG13	2.00	0.41
1:B:8:LEU:HD12	1:B:8:LEU:HA	1.95	0.41
1:A:151:LEU:HD22	1:A:190:TYR:CD1	2.56	0.41
1:B:60:ARG:HG2	1:B:62:MSE:HE2	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	190/209 (91%)	180 (95%)	10 (5%)	0	100	100
1	B	178/209 (85%)	172 (97%)	6 (3%)	0	100	100
All	All	368/418 (88%)	352 (96%)	16 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	132/153 (86%)	129 (98%)	3 (2%)	45	41
1	B	132/153 (86%)	130 (98%)	2 (2%)	60	59
All	All	264/306 (86%)	259 (98%)	5 (2%)	52	49

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	53	ASP
1	A	55	ILE
1	A	127	LEU
1	B	45	ARG
1	B	140	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	186	ASN
1	B	205	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	184/209 (88%)	0.32	7 (3%) 44 46	21, 32, 52, 59	0
1	B	175/209 (83%)	0.44	17 (9%) 15 15	20, 33, 55, 66	0
All	All	359/418 (85%)	0.38	24 (6%) 25 26	20, 32, 55, 66	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	141	VAL	4.3
1	A	136	TYR	4.1
1	A	140	VAL	3.5
1	B	204	ALA	3.4
1	B	160	ALA	3.2
1	A	139	ILE	3.2
1	B	172	GLY	3.1
1	B	72	PHE	3.0
1	B	149	PRO	3.0
1	A	160	ALA	2.9
1	A	173	ALA	2.9
1	B	23	LEU	2.9
1	B	49	ALA	2.8
1	A	78	GLY	2.6
1	B	7	ASP	2.6
1	B	106	THR	2.5
1	B	173	ALA	2.4
1	B	140	VAL	2.4
1	B	64	ASP	2.3
1	B	55	ILE	2.3
1	B	73	GLU	2.3
1	B	82	LEU	2.3
1	B	205	HIS	2.1
1	B	22	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.