



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 2, 2025 – 01:43 am BST

PDB ID : 2BJN / pdb_00002bjn
Title : X-ray Structure of human TPC6
Authors : Kummel, D.; Mueller, J.J.; Roske, Y.; Misselwitz, R.; Bussow, K.; Heinemann, U.
Deposited on : 2005-02-04
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5161 (1.70-1.70)
Clashscore	180529	5671 (1.70-1.70)
Ramachandran outliers	177936	5594 (1.70-1.70)
Sidechain outliers	177891	5594 (1.70-1.70)
RSRZ outliers	164620	5159 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	160	<div> <div>31%</div> <div>78%</div> <div>11%</div> <div>11%</div> </div>
1	B	160	<div> <div>27%</div> <div>79%</div> <div>11%</div> <div>7%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	1161	-	-	X	-
2	SO4	B	1160	-	-	X	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 2504 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRAFFICKING PROTEIN PARTICLE COMPLEX SUB-UNIT 6B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	143	Total	C	N	O	S	0	3	1
			1142	727	199	205	11			
1	B	149	Total	C	N	O	S	0	5	1
			1187	757	204	216	10			

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	58	Total	O	0	0
			58	58		
4	B	68	Total	O	0	0
			68	68		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: TRAFFICKING PROTEIN PARTICLE COMPLEX SUBUNIT 6B



• Molecule 1: TRAFFICKING PROTEIN PARTICLE COMPLEX SUBUNIT 6B



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	99.42Å 57.01Å 60.41Å 90.00° 117.16° 90.00°	Depositor
Resolution (Å)	22.98 – 1.70 22.98 – 1.70	Depositor EDS
% Data completeness (in resolution range)	98.6 (22.98-1.70) 98.5 (22.98-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.175 , 0.209 0.188 , 0.223	Depositor DCC
R_{free} test set	1697 reflections (5.19%)	wwPDB-VP
Wilson B-factor (Å ²)	26.5	Xtriage
Anisotropy	0.042	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 52.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2504	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	1/1177 (0.1%)	0.82	1/1572 (0.1%)
1	B	0.85	0/1227	0.92	5/1642 (0.3%)
All	All	0.81	1/2404 (0.0%)	0.87	6/3214 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	64	LEU	C-O	5.91	1.34	1.23

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	41	MET	CG-SD-CE	-10.74	83.02	100.20
1	A	41	MET	CG-SD-CE	-10.24	83.81	100.20
1	B	16	MET	CG-SD-CE	-8.93	85.92	100.20
1	B	10[A]	LEU	CA-CB-CG	-5.35	103.00	115.30
1	B	10[B]	LEU	CA-CB-CG	-5.35	103.00	115.30
1	B	156	MET	CG-SD-CE	5.10	108.36	100.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1142	0	1163	23	0
1	B	1187	0	1217	23	1
2	A	10	0	0	12	0
2	B	15	0	0	4	1
3	A	12	0	16	1	0
3	B	12	0	16	0	0
4	A	58	0	0	0	0
4	B	68	0	0	5	0
All	All	2504	0	2412	45	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:32:GLY:HA2	2:A:1161:SO4:O2	1.67	0.93
1:B:156:MET:HB3	2:B:1160:SO4:O3	1.71	0.89
1:B:156:MET:SD	4:B:2061:HOH:O	2.32	0.87
1:B:156:MET:HG3	4:B:2061:HOH:O	1.79	0.83
1:A:35:ILE:HG12	2:A:1161:SO4:O1	1.85	0.76
1:A:34:CYS:HB2	2:A:1161:SO4:O4	1.89	0.73
1:A:32:GLY:HA2	2:A:1161:SO4:S	2.31	0.70
1:B:154:GLN:NE2	1:B:156:MET:SD	2.65	0.69
1:B:156:MET:HB3	2:B:1160:SO4:S	2.34	0.67
1:B:31:ASN:HD22	1:B:31:ASN:C	2.01	0.64
1:B:61:LYS:HG2	1:B:62:ASP:N	2.13	0.63
1:A:29:VAL:O	2:A:1161:SO4:O2	2.18	0.62
1:A:35:ILE:CG1	2:A:1161:SO4:O1	2.48	0.62
1:B:85:ASN:OD1	1:B:87:ARG:HG3	2.01	0.60
1:A:34:CYS:CB	2:A:1161:SO4:O4	2.51	0.58
1:A:92:GLY:HA2	1:A:157:ILE:HG23	1.86	0.58
1:A:34:CYS:CA	2:A:1161:SO4:O4	2.53	0.57
1:A:29:VAL:O	2:A:1161:SO4:S	2.63	0.57
1:A:32:GLY:CA	2:A:1161:SO4:S	2.89	0.57
1:A:35:ILE:HD11	3:A:1163:GOL:H2	1.87	0.57
1:A:29:VAL:O	2:A:1161:SO4:O3	2.22	0.57
1:A:52:ARG:HG3	1:B:3:MET:HG2	1.88	0.55
1:B:31:ASN:ND2	1:B:33:ARG:H	2.05	0.54
1:A:32:GLY:HA2	2:A:1161:SO4:O1	2.08	0.54
1:B:31:ASN:HD22	1:B:32:GLY:N	2.06	0.53
1:B:31:ASN:C	1:B:31:ASN:ND2	2.63	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:156:MET:CB	2:B:1160:SO4:O3	2.53	0.51
1:A:92:GLY:CA	1:A:157:ILE:HG23	2.40	0.51
1:B:156:MET:CE	4:B:2061:HOH:O	2.58	0.49
1:B:116:HIS:CD2	1:B:116:HIS:C	2.87	0.47
1:A:92:GLY:HA2	1:A:157:ILE:CG2	2.46	0.46
1:A:87:ARG:NE	1:A:89:ASN:HD21	2.13	0.45
1:A:87:ARG:NE	1:A:89:ASN:ND2	2.65	0.45
1:B:156:MET:CG	4:B:2061:HOH:O	2.38	0.45
1:B:13:HIS:CE1	1:B:103:LEU:HD13	2.52	0.45
1:A:50:ILE:HD13	1:A:50:ILE:HA	1.82	0.44
1:B:140:ILE:N	1:B:140:ILE:HD13	2.34	0.43
1:A:87:ARG:HE	1:A:89:ASN:HD21	1.66	0.42
1:B:54:THR:N	2:B:1161:SO4:O4	2.36	0.41
1:B:55:LYS:HD3	1:B:60:PHE:CE2	2.55	0.41
1:B:10[A]:LEU:HD22	1:B:103:LEU:HD23	2.02	0.41
1:A:81:LYS:NZ	1:A:98:ASP:OD1	2.48	0.41
1:A:50:ILE:CG1	1:A:131:GLY:HA2	2.51	0.40
1:B:156:MET:HE2	4:B:2061:HOH:O	2.20	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:89:ASN:ND2	2:B:1160:SO4:O4[2_657]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	141/160 (88%)	135 (96%)	6 (4%)	0	100	100
1	B	150/160 (94%)	144 (96%)	5 (3%)	1 (1%)	19	7

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	291/320 (91%)	279 (96%)	11 (4%)	1 (0%)	30	23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	61	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	127/138 (92%)	123 (97%)	4 (3%)	35	18
1	B	133/138 (96%)	129 (97%)	4 (3%)	36	19
All	All	260/276 (94%)	252 (97%)	8 (3%)	34	18

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	85	ASN
1	A	106	GLN
1	A	146	SER
1	A	157	ILE
1	B	31	ASN
1	B	81	LYS
1	B	104	LEU
1	B	140	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	85	ASN
1	A	89	ASN
1	B	31	ASN
1	B	40	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	116	HIS
1	B	158	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	1161	3,1	4,4,4	0.26	0	6,6,6	0.27	0
3	GOL	B	1164	-	5,5,5	0.46	0	5,5,5	0.75	0
2	SO4	A	1160	-	4,4,4	0.55	0	6,6,6	0.36	0
2	SO4	B	1161	-	4,4,4	0.24	0	6,6,6	0.36	0
2	SO4	B	1160	-	4,4,4	0.19	0	6,6,6	0.62	0
2	SO4	B	1162	-	4,4,4	0.44	0	6,6,6	0.59	0
3	GOL	B	1163	-	5,5,5	0.49	0	5,5,5	0.22	0
3	GOL	A	1163	2	5,5,5	0.48	0	5,5,5	0.39	0
3	GOL	A	1162	-	5,5,5	0.28	0	5,5,5	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	1163	2	-	2/4/4/4	-
3	GOL	A	1162	-	-	4/4/4/4	-
3	GOL	B	1164	-	-	2/4/4/4	-
3	GOL	B	1163	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1162	GOL	C1-C2-C3-O3
3	A	1163	GOL	O1-C1-C2-C3
3	B	1164	GOL	O1-C1-C2-C3
3	A	1162	GOL	O2-C2-C3-O3
3	A	1163	GOL	O1-C1-C2-O2
3	A	1162	GOL	O1-C1-C2-O2
3	B	1164	GOL	O1-C1-C2-O2
3	A	1162	GOL	O1-C1-C2-C3

There are no ring outliers.

4 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1161	SO4	12	0
2	B	1161	SO4	1	0
2	B	1160	SO4	3	1
3	A	1163	GOL	1	0

5.7 Other polymers

There are no such residues in this entry.







5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	143/160 (89%)	1.66	50 (34%)  	12, 33, 40, 57	3 (2%)
1	B	149/160 (93%)	1.65	43 (28%)  	20, 33, 41, 48	5 (3%)
All	All	292/320 (91%)	1.66	93 (31%)  	12, 33, 41, 57	8 (2%)

All (93) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	58	ALA	5.6
1	A	129[C]	ARG	5.5
1	B	116	HIS	5.3
1	B	2	SER	5.1
1	A	159	LYS	4.9
1	A	137	ILE	4.9
1	A	105	THR	4.6
1	B	4	ALA	4.1
1	A	106	GLN	4.0
1	A	23	SER	4.0
1	B	146	SER	4.0
1	B	26	GLN	4.0
1	B	1	GLY	3.9
1	B	29	VAL	3.9
1	B	32	GLY	3.9
1	A	115	GLU	3.8
1	A	78	VAL	3.8
1	B	20	VAL	3.7
1	A	89	ASN	3.7
1	A	32	GLY	3.7
1	B	137	ILE	3.5
1	A	30	GLU	3.5
1	B	56	ASP	3.4
1	B	19	GLY	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	116	HIS	3.4
1	B	23[A]	SER	3.4
1	A	146	SER	3.3
1	A	136	GLY	3.3
1	A	90	HIS	3.2
1	B	3	MET	3.2
1	A	79	PHE	3.2
1	A	24	ALA	3.1
1	A	21	TYR	3.1
1	A	147	SER	3.0
1	B	60	PHE	3.0
1	B	25	GLU	3.0
1	B	156	MET	2.9
1	B	59	ARG	2.9
1	B	136	GLY	2.9
1	A	157	ILE	2.9
1	A	33	ARG	2.9
1	B	61	LYS	2.9
1	B	157	ILE	2.8
1	A	72	LYS	2.8
1	B	30	GLU	2.8
1	B	105	THR	2.8
1	A	107	MET	2.8
1	A	76	THR	2.8
1	A	27	GLY	2.8
1	A	75	TRP	2.7
1	B	27	GLY	2.7
1	A	88	THR	2.7
1	B	120	TYR	2.7
1	B	53	PHE	2.7
1	B	21	TYR	2.6
1	A	3	MET	2.6
1	B	7	ALA	2.6
1	A	74	PHE	2.5
1	A	81	LYS	2.5
1	B	140	ILE	2.5
1	B	57	THR	2.5
1	B	24	ALA	2.5
1	B	155	VAL	2.5
1	A	65	ASP	2.5
1	A	80	LYS	2.4
1	A	31	ASN	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	145	VAL	2.4
1	B	103	LEU	2.4
1	A	73	ASP	2.4
1	B	31	ASN	2.3
1	A	83	ILE	2.3
1	B	80	LYS	2.3
1	B	117	ALA	2.3
1	A	29	VAL	2.3
1	A	53	PHE	2.3
1	B	132	LEU	2.3
1	B	34	CYS	2.2
1	A	132	LEU	2.2
1	B	96	LEU	2.2
1	A	91	GLN	2.2
1	A	141	VAL	2.2
1	A	127[A]	LEU	2.2
1	B	131	GLY	2.2
1	A	139	SER	2.1
1	A	70	ILE	2.1
1	A	93	ILE	2.1
1	B	104	LEU	2.1
1	A	133	SER	2.1
1	A	22	LYS	2.1
1	A	138	LYS	2.1
1	A	151	CYS	2.1
1	B	138	LYS	2.1
1	A	26	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	A	1163	6/6	0.68	0.15	44,49,52,55	0
3	GOL	A	1162	6/6	0.71	0.18	43,49,50,53	0
2	SO4	B	1161	5/5	0.81	0.16	49,50,52,54	5
3	GOL	B	1163	6/6	0.81	0.15	35,41,42,44	0
3	GOL	B	1164	6/6	0.81	0.15	35,45,46,50	0
2	SO4	B	1160	5/5	0.85	0.22	30,33,38,38	5
2	SO4	A	1161	5/5	0.85	0.16	36,41,41,42	5
2	SO4	B	1162	5/5	0.98	0.13	24,26,28,30	0
2	SO4	A	1160	5/5	0.99	0.15	22,22,23,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.