



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2025 – 06:21 pm BST

PDB ID : 4BSG / pdb_00004bsg
Title : Crystal Structure of an H7N3 Avian Influenza Virus Haemagglutinin
Authors : Xiong, X.; Haire, L.F.; Martin, S.R.; Wharton, S.A.; Daniels, R.S.; Bennett, M.S.; McCauley, J.W.; Collins, P.J.; Walker, P.A.; Skehel, J.J.; Gamblin, S.J.
Deposited on : 2013-06-10
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.44


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	6234 (2.10-2.10)
Clashscore	180529	6893 (2.10-2.10)
Ramachandran outliers	177936	6839 (2.10-2.10)
Sidechain outliers	177891	6840 (2.10-2.10)
RSRZ outliers	164620	6234 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	321	<div> <div>12%</div> <div> <div></div> <div>89%</div> <div>10%</div> <div>.</div> </div> </div>
2	B	177	<div> <div>20%</div> <div> <div></div> <div>87%</div> <div>8%</div> <div>.</div> </div> </div>
3	C	2	<div> <div>50%</div> <div>50%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	A	404	X	-	-	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4215 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEMAGGLUTININ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	317	Total	C	N	O	S	0	0	1
			2418	1497	436	469	16			

- Molecule 2 is a protein called HEMAGGLUTININ.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	170	Total	C	N	O	S	0	0	0
			1380	850	243	279	8			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is SULFATE ION (CCD ID: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0
5	B	1	Total O S 5 4 1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	229	Total O 229 229	0	0
6	B	67	Total O 67 67	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: HEMAGGLUTININ



• Molecule 2: HEMAGGLUTININ



• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	116.06Å 116.06Å 296.31Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	32.40 – 2.10 32.40 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.0 (32.40-2.10) 99.0 (32.40-2.10)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.224 , 0.259 0.227 , 0.259	Depositor DCC
R_{free} test set	2273 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	37.3	Xtriage
Anisotropy	0.194	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.012 for $-1/3^*h+1/3^*k+1/3^*l,-k,8/3^*h+4/3^*k+1/3^*l$ 0.024 for $-2/3^*h-1/3^*k-1/3^*l,-1/3^*h-2/3^*k+1/3^*l,-4/3^*h+4/3^*k+1/3^*l$ 0.013 for $-h,1/3^*h-1/3^*k-1/3^*l,-4/3^*h-8/3^*k+1/3^*l$	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4215	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	1/2464 (0.0%)	0.70	0/3330
2	B	0.41	0/1404	0.69	0/1891
All	All	0.46	1/3868 (0.0%)	0.70	0/5221

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	316	GLU	C-N	-6.96	1.23	1.33

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2418	0	2363	20	0
2	B	1380	0	1281	10	0
3	C	28	0	25	2	0
4	A	28	0	26	0	0
5	A	50	0	0	0	0
5	B	15	0	0	0	0
6	A	229	0	0	7	0
6	B	67	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4215	0	3695	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:239:ASN:HB3	6:A:2150:HOH:O	1.74	0.85
6:B:2029:HOH:O	3:C:1:NAG:H83	1.94	0.68
1:A:228:LEU:HD22	1:A:234:VAL:HG23	1.77	0.66
2:B:98:LEU:HG	2:B:102:MET:HE2	1.83	0.60
2:B:75:LYS:HE3	3:C:1:NAG:H81	1.84	0.58
1:A:120:ILE:HG12	1:A:155:MET:HE1	1.88	0.56
1:A:158:SER:CB	6:A:2125:HOH:O	2.56	0.53
1:A:228:LEU:HD13	1:A:234:VAL:HG23	1.92	0.51
1:A:38:VAL:HG21	1:A:280:ILE:HD12	1.92	0.51
1:A:163:ARG:HD3	1:A:250:PHE:CZ	2.46	0.51
1:A:173:ILE:HD12	1:A:193:ILE:HD13	1.92	0.51
1:A:129:CYS:SG	6:A:2070:HOH:O	2.59	0.49
2:B:47:GLN:OE1	2:B:110:LEU:HD11	2.13	0.47
1:A:152:PHE:O	6:A:2122:HOH:O	2.20	0.47
2:B:129:ASN:HD22	2:B:129:ASN:N	2.10	0.47
1:A:44:LYS:HE2	1:A:269:GLU:HB2	1.97	0.46
1:A:228:LEU:HD13	1:A:234:VAL:CG2	2.45	0.46
1:A:226:LEU:HD12	1:A:226:LEU:C	2.41	0.45
2:B:55:LEU:HD22	2:B:99:LEU:HD21	2.00	0.43
1:A:228:LEU:CD2	1:A:234:VAL:HG23	2.45	0.43
1:A:298:ARG:HG2	2:B:92:TRP:CD2	2.53	0.43
2:B:56:ILE:HG22	2:B:56:ILE:O	2.17	0.43
1:A:235:THR:HG23	6:A:2125:HOH:O	2.18	0.43
1:A:170:ILE:CD1	1:A:225:TRP:HB3	2.49	0.42
1:A:194:THR:HG23	6:A:2151:HOH:O	2.19	0.42
1:A:60:ILE:CG2	1:A:170:ILE:HG13	2.51	0.41
1:A:170:ILE:O	1:A:245:PRO:HB3	2.20	0.41
2:B:88:MET:HE2	6:B:2045:HOH:O	2.22	0.41
2:B:62:GLN:HG2	2:B:64:GLU:OE2	2.21	0.40
6:A:2194:HOH:O	2:B:58:LYS:HB3	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	315/321 (98%)	305 (97%)	10 (3%)	0	100	100
2	B	168/177 (95%)	163 (97%)	5 (3%)	0	100	100
All	All	483/498 (97%)	468 (97%)	15 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/271 (98%)	261 (98%)	6 (2%)	47	53
2	B	145/152 (95%)	141 (97%)	4 (3%)	38	43
All	All	412/423 (97%)	402 (98%)	10 (2%)	44	49

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	HIS
1	A	94	ASN
1	A	122	THR
1	A	146	ASN
1	A	237	SER
1	A	256	MET
2	B	12	ASN
2	B	58	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	62	GLN
2	B	155	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	182	GLN
1	A	199	ASN
1	A	224	HIS
1	A	287	ASN
2	B	12	ASN
2	B	61	GLN
2	B	129	ASN
2	B	155	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	2,3	14,14,15	0.49	0	17,19,21	1.03	2 (11%)
3	NAG	C	2	3	14,14,15	0.54	0	17,19,21	0.99	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	2,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	NAG	C1-C2-N2	-2.30	106.56	110.49
3	C	1	NAG	C1-O5-C5	2.13	115.08	112.19
3	C	2	NAG	O5-C5-C6	2.08	110.47	107.20

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	2	NAG	O5-C5-C6-O6
3	C	2	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	1	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	SO4	A	1320	-	4,4,4	0.33	0	6,6,6	0.07	0
5	SO4	A	1321	-	4,4,4	0.39	0	6,6,6	0.16	0
4	NAG	A	403	1	14,14,15	0.41	0	17,19,21	1.35	1 (5%)
5	SO4	B	1171	-	4,4,4	0.33	0	6,6,6	0.08	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	A	404	1	14,14,15	0.43	0	17,19,21	1.37	2 (11%)
5	SO4	A	1326	-	4,4,4	0.33	0	6,6,6	0.10	0
5	SO4	A	1325	-	4,4,4	0.33	0	6,6,6	0.08	0
5	SO4	A	1324	-	4,4,4	0.35	0	6,6,6	0.05	0
5	SO4	A	1318	-	4,4,4	0.33	0	6,6,6	0.06	0
5	SO4	A	1323	-	4,4,4	0.32	0	6,6,6	0.12	0
5	SO4	B	1172	-	4,4,4	0.36	0	6,6,6	0.11	0
5	SO4	A	1319	-	4,4,4	0.35	0	6,6,6	0.09	0
5	SO4	A	1317	-	4,4,4	0.34	0	6,6,6	0.10	0
5	SO4	B	1173	-	4,4,4	0.34	0	6,6,6	0.09	0
5	SO4	A	1322	-	4,4,4	0.34	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	403	1	-	2/6/23/26	0/1/1/1
4	NAG	A	404	1	1/1/5/7	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	403	NAG	C1-O5-C5	4.96	118.91	112.19
4	A	404	NAG	C1-O5-C5	3.41	116.81	112.19
4	A	404	NAG	O5-C1-C2	-2.85	106.78	111.29

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	404	NAG	C1

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	404	NAG	O5-C5-C6-O6
4	A	403	NAG	C4-C5-C6-O6
4	A	404	NAG	C4-C5-C6-O6
4	A	403	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	317/321 (98%)	0.70	39 (12%) 9 10	24, 42, 73, 112	0
2	B	170/177 (96%)	1.07	35 (20%) 3 3	22, 70, 113, 131	0
All	All	487/498 (97%)	0.83	74 (15%) 6 7	22, 47, 103, 131	0

All (74) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	207	SER	6.3
1	A	192	LEU	5.0
1	A	4	CYS	4.5
2	B	58	LYS	4.4
1	A	129	CYS	4.3
1	A	205	VAL	4.2
2	B	34	THR	4.0
2	B	146	ASP	4.0
1	A	190	ASN	3.9
1	A	151	ALA	3.9
1	A	231	ASN	3.8
1	A	149	ASN	3.7
1	A	203	SER	3.6
2	B	59	THR	3.6
1	A	1	ASP	3.6
1	A	239	ASN	3.5
2	B	148	CYS	3.4
1	A	222	ASP	3.3
2	B	60	ASN	3.3
1	A	150	ALA	3.3
1	A	147	THR	3.3
2	B	170	ARG	3.2
2	B	153	ARG	3.2
1	A	206	PRO	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	204	PHE	3.1
1	A	153	PRO	3.1
2	B	141	PHE	3.1
1	A	198	SER	3.0
1	A	193	ILE	2.9
2	B	144	CYS	2.9
1	A	199	ASN	2.9
2	B	102	MET	2.9
2	B	110	LEU	2.9
1	A	189	GLY	2.9
1	A	191	LYS	2.9
1	A	194	THR	2.8
1	A	155	MET	2.8
2	B	161	ARG	2.7
2	B	29	ALA	2.7
1	A	13	GLY	2.7
1	A	183	THR	2.7
2	B	147	ASP	2.6
2	B	56	ILE	2.6
1	A	122	THR	2.5
1	A	152	PHE	2.5
2	B	127	ARG	2.5
2	B	160	SER	2.4
2	B	138	PHE	2.4
1	A	20	THR	2.4
2	B	156	THR	2.4
2	B	140	ILE	2.4
2	B	36	ALA	2.3
1	A	180	THR	2.3
1	A	188	SER	2.3
1	A	196	GLY	2.3
2	B	139	GLU	2.3
2	B	152	ILE	2.3
2	B	121	ARG	2.3
1	A	235	THR	2.3
2	B	31	GLY	2.2
2	B	155	ASN	2.2
1	A	201	GLN	2.2
2	B	143	LYS	2.2
1	A	237	SER	2.2
2	B	150	ALA	2.1
2	B	134	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	122	VAL	2.1
2	B	120	GLU	2.1
2	B	55	LEU	2.1
1	A	10	VAL	2.1
1	A	197	SER	2.1
1	A	185	LEU	2.1
2	B	169	ASN	2.0
2	B	27	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NAG	C	2	14/15	0.72	0.16	69,75,79,81	0
3	NAG	C	1	14/15	0.81	0.13	45,56,59,66	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	SO4	A	1317	5/5	0.46	0.13	115,118,118,119	0
4	NAG	A	403	14/15	0.58	0.17	61,73,76,76	0
4	NAG	A	404	14/15	0.65	0.15	65,74,76,78	0
5	SO4	B	1172	5/5	0.68	0.16	78,85,86,88	0
5	SO4	B	1173	5/5	0.68	0.20	91,93,94,96	0
5	SO4	A	1325	5/5	0.69	0.12	101,103,104,104	0
5	SO4	A	1318	5/5	0.70	0.14	96,96,98,99	0
5	SO4	A	1324	5/5	0.73	0.14	80,81,82,85	0
5	SO4	A	1320	5/5	0.74	0.13	118,120,121,122	0
5	SO4	A	1323	5/5	0.79	0.12	79,83,84,87	0
5	SO4	A	1322	5/5	0.80	0.11	89,91,93,94	0
5	SO4	A	1319	5/5	0.81	0.10	82,83,84,85	0
5	SO4	B	1171	5/5	0.83	0.10	87,87,89,90	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	SO4	A	1321	5/5	0.84	0.19	56,58,61,63	0
5	SO4	A	1326	5/5	0.92	0.07	72,73,75,76	0

6.5 Other polymers [i](#)

There are no such residues in this entry.