



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 21, 2025 – 06:09 PM EDT

PDB ID : 9C6U / pdb_00009c6u
Title : [d2d3] Tensegrity triangle with deazapurine center and helical strands (strands 2+3) with P63 symmetry
Authors : Vecchioni, S.; Sha, R.; Ohayon, Y.; Galindo, M.; Lopez-Chamorro, C.; Jong, M.; Woloszyn, K.
Deposited on : 2024-06-09
Resolution : 5.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.44


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 5.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1015 (6.06-3.94)
Clashscore	180529	1031 (6.04-3.96)
RSRZ outliers	164620	1011 (6.06-3.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	21	
2	B	7	
3	C	6	
4	D	8	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 855 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(* (7GU)P*(7DA)P*(7GU)P*CP*(7DA)P*(7GU)P*CP*CP*TP*(7GU)P*TP*(7DA)P*CP*(7GU)P*(7GU)P*(7DA)P*CP*(7DA)P*TP*CP*(7DA))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	21	Total	C	N	O	P	0	0	0
			429	214	74	121	20			

- Molecule 2 is a DNA chain called DNA (5'-D(P*CP*CP*(7GU)P*TP*(7DA)P*CP*(7DA))-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	7	Total	C	N	O	P	0	0	0
			141	70	23	41	7			

- Molecule 3 is a DNA chain called DNA (5'-D(P*GP*GP*CP*TP*GP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	6	Total	C	N	O	P	0	0	0
			124	58	23	37	6			


- Molecule 4 is a DNA chain called DNA (5'-D(*TP*CP*TP*GP*AP*TP*GP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	8	Total	C	N	O	P	0	0	0
			161	79	26	49	7			

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D*(7GU)P*(7DA)P*(7GU)P*CP*(7DA)P*(7GU)P*CP*CP*TP*(7GU)P*TP*(7DA)P*CP*(7GU)P*(7GU)P*(7DA)P*CP*(7DA)P*TP*CP*(7DA))-3')

Chain A: 




- Molecule 2: DNA (5'-D(P*CP*CP*(7GU)P*TP*(7DA)P*CP*(7DA))-3')

Chain B: 




- Molecule 3: DNA (5'-D(P*GP*GP*CP*TP*GP*C)-3')

Chain C: 



- Molecule 4: DNA (5'-D(*TP*CP*TP*GP*AP*TP*GP*T)-3')

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	125.39Å 125.39Å 59.37Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	33.76 – 5.00 33.76 – 5.00	Depositor EDS
% Data completeness (in resolution range)	79.1 (33.76-5.00) 70.2 (33.76-5.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.18 (at 5.07Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.247 , 0.295 0.248 , 0.294	Depositor DCC
R_{free} test set	128 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	175.6	Xtriage
Anisotropy	0.678	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.09 , 0.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.099 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	855	wwPDB-VP
Average B, all atoms (Å ²)	275.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 7GU, 7DA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/231	0.58	0/338
2	B	0.20	0/82	0.35	0/118
3	C	0.18	0/138	0.33	0/211
4	D	0.19	0/179	0.49	0/275
All	All	0.25	0/630	0.48	0/942

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	429	0	246	14	0
2	B	141	0	82	6	0
3	C	124	0	68	3	0
4	D	161	0	94	3	0
All	All	855	0	490	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:115:7GU:H2''	1:A:116:7DA:H8	1.64	0.78
1:A:102:DA:H2'	1:A:103:7GU:H8	1.75	0.69
1:A:109:DT:H2'	1:A:110:7GU:C8	2.24	0.68
2:B:120:DC:H2''	2:B:121:7GU:C7	2.28	0.64
2:B:124:DC:H2''	2:B:125:7DA:H5'	1.82	0.61
4:D:205:DA:H2''	4:D:206:DT:H71	1.83	0.61
1:A:120:DC:H2'	1:A:121:7DA:H8	1.88	0.54
2:B:119:DC:H2''	2:B:120:DC:H5'	1.89	0.54
2:B:124:DC:H2'	2:B:125:7DA:C8	2.38	0.54
4:D:202:DC:C2	4:D:203:DT:H72	2.43	0.53
2:B:119:DC:O4'	4:D:208:DT:H2''	2.09	0.52
1:A:120:DC:H2'	1:A:121:7DA:C8	2.40	0.51
3:C:211:DC:H2'	3:C:212:DT:H72	1.94	0.48
3:C:210:DG:H2'	3:C:211:DC:C6	2.49	0.48
1:A:102:DA:H2'	1:A:103:7GU:C8	2.41	0.47
3:C:211:DC:H5''	3:C:211:DC:H6	1.79	0.47
1:A:113:DC:H2''	1:A:114:7GU:C7	2.47	0.45
1:A:119:DT:H2''	1:A:120:DC:O5'	2.16	0.45
1:A:104:DC:H2''	1:A:105:7DA:H8	1.98	0.45
1:A:101:DG:N3	1:A:102:DA:C8	2.86	0.43
1:A:103:7GU:H2''	1:A:104:DC:O5'	2.19	0.43
1:A:108:DC:H2'	1:A:109:DT:C6	2.55	0.42
2:B:121:7GU:H2''	2:B:122:DT:O5'	2.20	0.41
1:A:107:DC:C2	1:A:108:DC:C5	3.08	0.41
1:A:115:7GU:H2''	1:A:116:7DA:C8	2.45	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

There are no protein molecules in this entry.

5.3.2 Protein sidechains ⓘ

There are no protein molecules in this entry.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

13 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	7DA	A	121	1,4	18,23,24	3.79	9 (50%)	18,33,36	4.24	6 (33%)
2	7DA	B	125	2,1	18,23,24	3.82	9 (50%)	18,33,36	4.12	6 (33%)
1	7GU	A	115	2,1	20,24,25	3.85	14 (70%)	21,35,38	5.08	5 (23%)
1	7GU	A	110	2,1	20,24,25	3.80	13 (65%)	21,35,38	5.03	4 (19%)
1	7DA	A	105	1,3	18,23,24	3.76	9 (50%)	18,33,36	4.19	6 (33%)
2	7DA	B	123	2,1	18,23,24	3.81	9 (50%)	18,33,36	4.03	5 (27%)
1	7DA	A	118	1,4	18,23,24	3.80	9 (50%)	18,33,36	4.12	5 (27%)
1	7DA	A	112	2,1	18,23,24	3.74	9 (50%)	18,33,36	4.31	7 (38%)
1	7GU	A	114	2,1	20,24,25	3.81	15 (75%)	21,35,38	5.03	4 (19%)
1	7GU	A	103	1,3	20,24,25	3.79	13 (65%)	21,35,38	5.15	5 (23%)
1	7GU	A	106	1,3	20,24,25	3.81	13 (65%)	21,35,38	4.98	4 (19%)
1	7DA	A	116	1,4	18,23,24	3.84	9 (50%)	18,33,36	3.82	7 (38%)
2	7GU	B	121	2,1	20,24,25	3.81	14 (70%)	21,35,38	5.27	5 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	7DA	A	121	1,4	-	0/3/21/22	0/3/3/3
2	7DA	B	125	2,1	-	2/3/21/22	0/3/3/3
1	7GU	A	115	2,1	-	1/3/21/22	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	7GU	A	110	2,1	-	3/3/21/22	0/3/3/3
1	7DA	A	105	1,3	-	0/3/21/22	0/3/3/3
2	7DA	B	123	2,1	-	0/3/21/22	0/3/3/3
1	7DA	A	118	1,4	-	2/3/21/22	0/3/3/3
1	7DA	A	112	2,1	-	2/3/21/22	0/3/3/3
1	7GU	A	114	2,1	-	0/3/21/22	0/3/3/3
1	7GU	A	103	1,3	-	0/3/21/22	0/3/3/3
1	7GU	A	106	1,3	-	0/3/21/22	0/3/3/3
1	7DA	A	116	1,4	-	0/3/21/22	0/3/3/3
2	7GU	B	121	2,1	-	2/3/21/22	0/3/3/3

All (145) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	114	7GU	C2'-C3'	-8.00	1.32	1.52
1	A	110	7GU	C2'-C3'	-7.91	1.32	1.52
2	B	121	7GU	C2'-C3'	-7.82	1.33	1.52
1	A	115	7GU	C2'-C3'	-7.76	1.33	1.52
1	A	103	7GU	C2'-C3'	-7.73	1.33	1.52
1	A	106	7GU	C2'-C3'	-7.66	1.33	1.52
1	A	116	7DA	C6-C5	-7.51	1.38	1.45
2	B	125	7DA	C6-C5	-7.50	1.38	1.45
2	B	123	7DA	C6-C5	-7.43	1.38	1.45
1	A	118	7DA	C6-C5	-7.31	1.38	1.45
1	A	112	7DA	C6-C5	-7.18	1.38	1.45
1	A	121	7DA	C6-C5	-7.00	1.38	1.45
1	A	105	7DA	C6-C5	-6.95	1.38	1.45
1	A	121	7DA	O4'-C4'	6.59	1.59	1.45
1	A	116	7DA	C3'-C4'	-6.52	1.35	1.53
2	B	123	7DA	O4'-C4'	6.48	1.59	1.45
2	B	125	7DA	C3'-C4'	-6.48	1.35	1.53
1	A	105	7DA	C3'-C4'	-6.48	1.36	1.53
1	A	116	7DA	O4'-C4'	6.47	1.59	1.45
1	A	121	7DA	C3'-C4'	-6.45	1.36	1.53
1	A	112	7DA	O4'-C4'	6.45	1.59	1.45
1	A	118	7DA	C3'-C4'	-6.44	1.36	1.53
2	B	125	7DA	O4'-C4'	6.44	1.59	1.45
2	B	123	7DA	C3'-C4'	-6.44	1.36	1.53
1	A	118	7DA	O4'-C4'	6.40	1.59	1.45
1	A	105	7DA	O4'-C4'	6.39	1.59	1.45
1	A	121	7DA	O4'-C1'	6.31	1.56	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	112	7DA	C3'-C4'	-6.16	1.36	1.53
1	A	105	7DA	O4'-C1'	6.15	1.56	1.42
1	A	118	7DA	O4'-C1'	6.14	1.56	1.42
1	A	116	7DA	O4'-C1'	6.14	1.56	1.42
2	B	123	7DA	O4'-C1'	6.07	1.55	1.42
2	B	125	7DA	O4'-C1'	5.99	1.55	1.42
1	A	115	7GU	C2-N3	5.87	1.47	1.33
1	A	106	7GU	C2-N3	5.83	1.47	1.33
1	A	112	7DA	O4'-C1'	5.82	1.55	1.42
1	A	110	7GU	C2-N3	5.80	1.47	1.33
1	A	114	7GU	C2-N3	5.76	1.47	1.33
1	A	103	7GU	C2-N3	5.72	1.47	1.33
2	B	121	7GU	C2-N3	5.70	1.47	1.33
1	A	106	7GU	O4'-C4'	5.56	1.57	1.45
2	B	121	7GU	O4'-C4'	5.55	1.57	1.45
2	B	123	7DA	C2'-C1'	-5.54	1.37	1.52
1	A	115	7GU	O4'-C4'	5.51	1.57	1.45
1	A	112	7DA	C2'-C1'	-5.50	1.37	1.52
1	A	103	7GU	O4'-C4'	5.50	1.57	1.45
2	B	125	7DA	C2'-C1'	-5.48	1.37	1.52
1	A	110	7GU	O4'-C4'	5.47	1.57	1.45
1	A	118	7DA	C2'-C1'	-5.45	1.37	1.52
1	A	114	7GU	O4'-C4'	5.44	1.57	1.45
1	A	116	7DA	C2'-C1'	-5.39	1.37	1.52
1	A	105	7DA	C2'-C1'	-5.36	1.37	1.52
1	A	115	7GU	C4-N3	5.31	1.50	1.37
1	A	106	7GU	C4-N3	5.29	1.50	1.37
1	A	121	7DA	C2'-C1'	-5.28	1.38	1.52
1	A	110	7GU	C4-N3	5.26	1.49	1.37
1	A	114	7GU	C4-N3	5.24	1.49	1.37
2	B	121	7GU	C4-N3	5.19	1.49	1.37
1	A	103	7GU	C4-N3	5.15	1.49	1.37
2	B	121	7GU	O4'-C1'	-5.13	1.30	1.42
1	A	115	7GU	O4'-C1'	-5.08	1.31	1.42
1	A	114	7GU	O4'-C1'	-5.05	1.31	1.42
1	A	106	7GU	O4'-C1'	-5.04	1.31	1.42
1	A	103	7GU	O4'-C1'	-4.99	1.31	1.42
1	A	110	7GU	O4'-C1'	-4.97	1.31	1.42
1	A	106	7GU	C2-N2	4.81	1.45	1.34
1	A	115	7GU	C2-N2	4.78	1.45	1.34
2	B	121	7GU	C2-N2	4.76	1.45	1.34
1	A	114	7GU	C2-N2	4.74	1.45	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	110	7GU	C2-N2	4.74	1.45	1.34
1	A	103	7GU	C2-N2	4.69	1.45	1.34
1	A	115	7GU	C2'-C1'	4.27	1.64	1.52
1	A	106	7GU	C2'-C1'	4.20	1.63	1.52
1	A	110	7GU	C2'-C1'	4.18	1.63	1.52
1	A	103	7GU	C2'-C1'	4.15	1.63	1.52
1	A	114	7GU	C2'-C1'	4.11	1.63	1.52
2	B	121	7GU	C2'-C1'	4.07	1.63	1.52
1	A	105	7DA	C2'-C3'	4.04	1.63	1.52
1	A	121	7DA	C2'-C3'	4.00	1.63	1.52
1	A	116	7DA	C2'-C3'	3.98	1.63	1.52
1	A	114	7GU	C5'-C4'	-3.92	1.39	1.51
2	B	125	7DA	C2'-C3'	3.90	1.62	1.52
1	A	115	7GU	C5'-C4'	-3.90	1.39	1.51
1	A	118	7DA	C2'-C3'	3.90	1.62	1.52
1	A	112	7DA	C2'-C3'	3.88	1.62	1.52
1	A	103	7GU	C5'-C4'	-3.87	1.39	1.51
2	B	123	7DA	C2'-C3'	3.87	1.62	1.52
1	A	106	7GU	C5'-C4'	-3.85	1.40	1.51
1	A	110	7GU	C5'-C4'	-3.85	1.40	1.51
2	B	121	7GU	C5'-C4'	-3.69	1.40	1.51
1	A	106	7GU	C5-C6	3.69	1.53	1.47
2	B	121	7GU	C5-C6	3.63	1.53	1.47
1	A	110	7GU	C5-C6	3.62	1.53	1.47
1	A	115	7GU	C5-C6	3.59	1.53	1.47
1	A	103	7GU	C5-C6	3.56	1.53	1.47
1	A	121	7DA	C6-N6	3.49	1.46	1.34
1	A	115	7GU	O3'-C3'	3.47	1.50	1.43
2	B	121	7GU	C6-N1	3.42	1.43	1.37
1	A	112	7DA	C6-N6	3.41	1.46	1.34
1	A	114	7GU	C5-C6	3.41	1.53	1.47
1	A	105	7DA	C6-N6	3.41	1.46	1.34
1	A	116	7DA	C6-N6	3.41	1.46	1.34
1	A	118	7DA	C6-N6	3.40	1.46	1.34
2	B	125	7DA	C6-N6	3.40	1.46	1.34
2	B	123	7DA	C6-N6	3.39	1.46	1.34
1	A	106	7GU	C6-N1	3.38	1.43	1.37
1	A	103	7GU	O3'-C3'	3.33	1.50	1.43
1	A	110	7GU	C6-N1	3.30	1.42	1.37
1	A	114	7GU	C6-N1	3.30	1.42	1.37
1	A	115	7GU	C6-N1	3.27	1.42	1.37
1	A	103	7GU	C6-N1	3.26	1.42	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	116	7DA	C2-N3	3.13	1.36	1.32
1	A	112	7DA	C2-N3	3.04	1.36	1.32
1	A	105	7DA	C2-N3	2.99	1.36	1.32
1	A	121	7DA	C2-N3	2.99	1.36	1.32
2	B	125	7DA	C2-N3	2.97	1.36	1.32
1	A	118	7DA	C2-N3	2.95	1.36	1.32
2	B	123	7DA	C2-N3	2.95	1.36	1.32
2	B	121	7GU	O3'-C3'	2.93	1.49	1.43
1	A	106	7GU	O3'-C3'	2.88	1.49	1.43
1	A	121	7DA	C5-C4	-2.84	1.36	1.43
1	A	118	7DA	C5-C4	-2.84	1.36	1.43
1	A	114	7GU	C2-N1	2.80	1.44	1.37
1	A	110	7GU	O3'-C3'	2.80	1.49	1.43
2	B	123	7DA	C5-C4	-2.79	1.36	1.43
1	A	106	7GU	C2-N1	2.79	1.44	1.37
1	A	112	7DA	C5-C4	-2.79	1.36	1.43
2	B	125	7DA	C5-C4	-2.78	1.36	1.43
2	B	121	7GU	C2-N1	2.78	1.44	1.37
1	A	110	7GU	C2-N1	2.77	1.44	1.37
1	A	103	7GU	C2-N1	2.76	1.44	1.37
1	A	115	7GU	C2-N1	2.76	1.44	1.37
1	A	105	7DA	C5-C4	-2.76	1.36	1.43
1	A	116	7DA	C5-C4	-2.69	1.36	1.43
1	A	114	7GU	O3'-C3'	2.54	1.48	1.43
2	B	121	7GU	C5-C4	-2.44	1.37	1.41
1	A	114	7GU	C5-C4	-2.41	1.38	1.41
1	A	115	7GU	C5-C4	-2.30	1.38	1.41
1	A	103	7GU	C5-C4	-2.26	1.38	1.41
1	A	110	7GU	C5-C4	-2.18	1.38	1.41
1	A	106	7GU	C5-C4	-2.16	1.38	1.41
1	A	114	7GU	C1'-N9	-2.13	1.43	1.49
2	B	121	7GU	C1'-N9	-2.11	1.43	1.49
1	A	114	7GU	O6-C6	-2.02	1.18	1.23
1	A	115	7GU	O6-C6	-2.01	1.18	1.23

All (69) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	121	7GU	C8-N9-C1'	22.83	147.93	125.37
1	A	103	7GU	C8-N9-C1'	22.37	147.46	125.37
1	A	110	7GU	C8-N9-C1'	21.89	146.99	125.37
1	A	115	7GU	C8-N9-C1'	21.81	146.91	125.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	114	7GU	C8-N9-C1'	21.76	146.87	125.37
1	A	106	7GU	C8-N9-C1'	21.50	146.61	125.37
1	A	105	7DA	C8-N9-C1'	11.33	136.56	125.37
1	A	118	7DA	C8-N9-C1'	11.10	136.34	125.37
1	A	121	7DA	C8-N9-C1'	11.09	136.33	125.37
2	B	125	7DA	C8-N9-C1'	11.01	136.25	125.37
1	A	112	7DA	C8-N9-C1'	10.95	136.19	125.37
2	B	123	7DA	C8-N9-C1'	10.51	135.75	125.37
1	A	121	7DA	C5-C6-N6	9.70	136.34	121.45
1	A	112	7DA	C5-C6-N6	9.54	136.11	121.45
1	A	105	7DA	C5-C6-N6	9.50	136.04	121.45
1	A	118	7DA	C5-C6-N6	9.32	135.77	121.45
2	B	125	7DA	C5-C6-N6	9.25	135.66	121.45
1	A	116	7DA	C5-C6-N6	9.25	135.65	121.45
2	B	123	7DA	C5-C6-N6	9.23	135.63	121.45
1	A	116	7DA	C8-N9-C1'	8.79	134.06	125.37
1	A	121	7DA	N6-C6-N1	-6.97	103.44	118.33
1	A	105	7DA	N6-C6-N1	-6.92	103.54	118.33
1	A	112	7DA	N6-C6-N1	-6.91	103.58	118.33
1	A	118	7DA	N6-C6-N1	-6.74	103.94	118.33
2	B	125	7DA	N6-C6-N1	-6.64	104.15	118.33
2	B	123	7DA	N6-C6-N1	-6.63	104.16	118.33
1	A	116	7DA	N6-C6-N1	-6.60	104.23	118.33
1	A	121	7DA	N3-C2-N1	-5.91	120.65	128.67
1	A	112	7DA	N3-C2-N1	-5.87	120.70	128.67
1	A	116	7DA	N3-C2-N1	-5.83	120.75	128.67
2	B	123	7DA	N3-C2-N1	-5.79	120.82	128.67
2	B	125	7DA	N3-C2-N1	-5.74	120.88	128.67
1	A	118	7DA	N3-C2-N1	-5.73	120.89	128.67
1	A	105	7DA	N3-C2-N1	-5.63	121.03	128.67
1	A	115	7GU	C5-C6-N1	4.92	120.12	114.93
1	A	110	7GU	C5-C6-N1	4.78	119.98	114.93
2	B	121	7GU	C5-C6-N1	4.71	119.91	114.93
1	A	106	7GU	C5-C6-N1	4.64	119.83	114.93
1	A	114	7GU	C5-C6-N1	4.62	119.81	114.93
1	A	103	7GU	C5-C6-N1	4.59	119.77	114.93
1	A	112	7DA	O4'-C1'-N9	3.64	113.57	108.75
1	A	112	7DA	C4'-O4'-C1'	-3.09	102.18	109.51
1	A	115	7GU	C3'-C2'-C1'	2.83	109.51	102.60
1	A	106	7GU	C3'-C2'-C1'	2.81	109.48	102.60
1	A	110	7GU	C2-N1-C6	-2.72	120.14	125.11
1	A	115	7GU	C2-N1-C6	-2.71	120.14	125.11

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	114	7GU	C2'-C1'-N9	-2.71	108.33	114.61
2	B	121	7GU	C3'-C2'-C1'	2.69	109.18	102.60
1	A	103	7GU	C2-N1-C6	-2.67	120.21	125.11
1	A	106	7GU	C2-N1-C6	-2.61	120.33	125.11
2	B	121	7GU	C2-N1-C6	-2.58	120.39	125.11
1	A	103	7GU	C3'-C2'-C1'	2.56	108.87	102.60
1	A	121	7DA	C2'-C3'-C4'	2.56	107.99	102.80
1	A	114	7GU	C2-N1-C6	-2.51	120.51	125.11
1	A	116	7DA	C2'-C1'-N9	-2.37	109.10	114.61
1	A	121	7DA	C3'-C2'-C1'	2.33	108.30	102.60
1	A	116	7DA	C3'-C2'-C1'	2.30	108.23	102.60
2	B	125	7DA	C2'-C3'-C4'	2.27	107.41	102.80
2	B	123	7DA	C2'-C1'-N9	-2.20	109.50	114.61
1	A	103	7GU	C2'-C3'-C4'	2.20	107.26	102.80
1	A	105	7DA	C3'-C2'-C1'	2.16	107.88	102.60
1	A	110	7GU	C3'-C2'-C1'	2.12	107.79	102.60
1	A	105	7DA	C2'-C3'-C4'	2.12	107.10	102.80
1	A	118	7DA	C2'-C1'-N9	-2.09	109.75	114.61
1	A	115	7GU	C2'-C3'-C4'	2.09	107.03	102.80
2	B	121	7GU	C2'-C1'-N9	-2.07	109.80	114.61
1	A	116	7DA	C2'-C3'-C4'	2.06	106.99	102.80
1	A	112	7DA	O4'-C1'-C2'	-2.01	102.48	106.24
2	B	125	7DA	C2'-C1'-N9	-2.01	109.94	114.61

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	118	7DA	C3'-C4'-C5'-O5'
1	A	118	7DA	O4'-C4'-C5'-O5'
1	A	110	7GU	C3'-C4'-C5'-O5'
2	B	125	7DA	O4'-C4'-C5'-O5'
1	A	112	7DA	C3'-C4'-C5'-O5'
1	A	110	7GU	O4'-C4'-C5'-O5'
1	A	112	7DA	O4'-C4'-C5'-O5'
1	A	110	7GU	C4'-C5'-O5'-P
2	B	125	7DA	C4'-C5'-O5'-P
2	B	121	7GU	O4'-C4'-C5'-O5'
2	B	121	7GU	C3'-C4'-C5'-O5'
1	A	115	7GU	C3'-C4'-C5'-O5'

There are no ring outliers.

9 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	121	7DA	2	0
2	B	125	7DA	2	0
1	A	115	7GU	2	0
1	A	110	7GU	1	0
1	A	105	7DA	1	0
1	A	114	7GU	1	0
1	A	103	7GU	3	0
1	A	116	7DA	2	0
2	B	121	7GU	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	11/21 (52%)	0.47	0 100 100	196, 281, 334, 337	0
2	B	4/7 (57%)	0.25	0 100 100	215, 251, 252, 267	0
3	C	6/6 (100%)	-0.04	0 100 100	245, 285, 314, 315	0
4	D	8/8 (100%)	0.03	0 100 100	246, 298, 338, 342	0
All	All	29/42 (69%)	0.21	0 100 100	196, 281, 337, 342	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	7DA	A	105	21/22	0.82	0.14	277,307,326,337	0
1	7DA	A	118	21/22	0.82	0.12	282,303,325,333	0
1	7GU	A	115	22/23	0.83	0.12	234,257,274,288	0
1	7DA	A	116	21/22	0.87	0.09	241,277,298,316	0
2	7GU	B	121	22/23	0.87	0.09	229,251,269,278	0
1	7GU	A	106	22/23	0.88	0.11	271,303,331,341	0
1	7DA	A	121	21/22	0.89	0.11	321,331,339,344	0
1	7DA	A	112	21/22	0.90	0.16	172,187,218,227	0
1	7GU	A	110	22/23	0.91	0.10	189,198,210,220	0
1	7GU	A	114	22/23	0.92	0.13	207,235,257,280	0
1	7GU	A	103	22/23	0.92	0.10	286,305,317,325	0
2	7DA	B	123	21/22	0.94	0.18	215,248,275,286	0
2	7DA	B	125	21/22	0.94	0.16	193,206,226,232	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.