



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 16, 2024 – 02:33 PM EDT

PDB ID : 5C9S
Title : Crystal Structure of the C-terminal domain of Rrp5
Authors : Khoshnevis, S.; Karbstein, K.
Deposited on : 2015-06-28
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	572	 2% 31% 14% 52%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2317 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called rRNA biogenesis protein RRP5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	276	2258	1449	378	426	5	0	0	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1158	GLY	-	expression tag	UNP Q05022
B	1159	PRO	-	expression tag	UNP Q05022
B	1160	LEU	-	expression tag	UNP Q05022
B	1161	GLY	-	expression tag	UNP Q05022
B	1162	SER	-	expression tag	UNP Q05022

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	B	1	6	3	3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	53	Total 53	O 53	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	114.14Å 114.14Å 67.84Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.67 – 2.70 43.67 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.8 (43.67-2.70) 99.9 (43.67-2.70)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.90 (at 2.69Å)	Xtriage
Refinement program	PHENIX 1.7.1_743	Depositor
R, R_{free}	0.221 , 0.283 0.225 , 0.263	Depositor DCC
R_{free} test set	719 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	34.7	Xtriage
Anisotropy	0.529	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 32.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.047 for -h,-k,l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	2317	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	B	1.04	5/2302 (0.2%)	0.73	4/3097 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1589	GLY	C-O	-5.42	1.15	1.23
1	B	1585	TRP	CE3-CZ3	-5.39	1.29	1.38
1	B	1658	TYR	CE1-CZ	-5.25	1.31	1.38
1	B	1585	TRP	CD2-CE2	-5.19	1.35	1.41
1	B	1585	TRP	CZ3-CH2	-5.01	1.32	1.40

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1654	LEU	CB-CG-CD2	-5.52	101.61	111.00
1	B	1612	LEU	C-N-CD	5.38	139.70	128.40
1	B	1590	ASP	CB-CG-OD1	5.30	123.07	118.30
1	B	1569	LEU	CA-CB-CG	5.17	127.19	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2258	0	2248	70	0
2	B	6	0	8	2	0
3	B	53	0	0	0	0
All	All	2317	0	2256	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1659:VAL:O	1:B:1663:VAL:HG13	1.78	0.84
1:B:1516:ILE:HG22	1:B:1520:MET:HE2	1.63	0.81
1:B:1653:ASP:O	1:B:1657:VAL:HG23	1.80	0.80
1:B:1689:LYS:O	1:B:1693:ASN:ND2	2.17	0.78
1:B:1622:ARG:NH1	1:B:1626:GLN:OE1	2.20	0.74
1:B:1517:TRP:HA	1:B:1520:MET:HE3	1.70	0.72
1:B:1410:THR:HG21	1:B:1480:ASN:OD1	1.89	0.72
1:B:1651:ARG:HH22	2:B:1801:GOL:H32	1.57	0.69
1:B:1608:ALA:O	1:B:1612:LEU:HB2	1.93	0.69
1:B:1622:ARG:HG2	1:B:1622:ARG:HH11	1.59	0.68
1:B:1500:ARG:O	1:B:1504:THR:HG23	1.93	0.68
1:B:1516:ILE:HG22	1:B:1520:MET:CE	2.25	0.67
1:B:1617:HIS:O	1:B:1621:VAL:HG23	1.95	0.65
1:B:1517:TRP:HA	1:B:1520:MET:CE	2.27	0.65
1:B:1578:GLY:O	1:B:1612:LEU:CD1	2.47	0.62
1:B:1667:ASP:OD1	1:B:1670:LYS:HB2	2.02	0.59
1:B:1705:GLU:HA	1:B:1708:ILE:HB	1.86	0.57
1:B:1579:GLY:O	1:B:1580:GLU:HG3	2.05	0.57
1:B:1409:TRP:CH2	1:B:1519:ALA:HB1	2.39	0.57
1:B:1716:THR:O	1:B:1719:VAL:HG12	2.05	0.56
1:B:1475:SER:O	1:B:1479:MET:HG3	2.06	0.55
1:B:1705:GLU:O	1:B:1709:GLU:HG2	2.06	0.55
1:B:1663:VAL:HG23	1:B:1663:VAL:O	2.06	0.55
1:B:1662:GLU:C	1:B:1664:LYS:H	2.11	0.53
1:B:1529:THR:OG1	1:B:1531:GLU:HG2	2.10	0.52
1:B:1622:ARG:NH1	1:B:1622:ARG:HG2	2.22	0.52
1:B:1707:THR:HA	1:B:1710:TYR:HB3	1.91	0.51
1:B:1683:ILE:HG23	1:B:1687:GLN:HB2	1.92	0.51
1:B:1539:ARG:O	1:B:1542:GLN:HB2	2.11	0.51
1:B:1534:GLU:O	1:B:1538:SER:HB2	2.12	0.50
1:B:1693:ASN:ND2	1:B:1693:ASN:H	2.09	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1665:ALA:O	1:B:1666:LYS:HG2	2.11	0.50
1:B:1582:VAL:HG22	1:B:1612:LEU:HD21	1.95	0.49
1:B:1693:ASN:H	1:B:1693:ASN:HD22	1.60	0.49
1:B:1598:GLU:OE2	1:B:1628:GLU:OE2	2.31	0.49
1:B:1652:ILE:O	1:B:1652:ILE:HG22	2.11	0.49
1:B:1673:ASP:N	1:B:1673:ASP:OD1	2.47	0.47
1:B:1517:TRP:CE3	1:B:1520:MET:HE1	2.49	0.47
1:B:1707:THR:O	1:B:1711:VAL:HG23	2.14	0.47
1:B:1573:THR:O	1:B:1573:THR:HG22	2.15	0.47
1:B:1684:THR:HG22	1:B:1687:GLN:H	1.80	0.47
1:B:1674:LEU:O	1:B:1678:ILE:HG13	2.14	0.46
1:B:1659:VAL:HG12	1:B:1674:LEU:HD23	1.99	0.45
1:B:1585:TRP:CD2	1:B:1608:ALA:HB2	2.51	0.45
1:B:1659:VAL:HG12	1:B:1674:LEU:HG	1.98	0.45
1:B:1579:GLY:C	1:B:1580:GLU:HG3	2.37	0.45
1:B:1648:ALA:C	1:B:1650:LYS:H	2.19	0.45
1:B:1409:TRP:HH2	1:B:1519:ALA:HB1	1.78	0.45
1:B:1576:LYS:HD3	1:B:1577:PHE:CZ	2.51	0.45
1:B:1535:GLU:O	1:B:1539:ARG:HG3	2.17	0.44
1:B:1474:SER:OG	1:B:1477:VAL:HG23	2.18	0.43
1:B:1517:TRP:CD2	1:B:1520:MET:HE1	2.54	0.43
1:B:1517:TRP:CA	1:B:1520:MET:HE3	2.42	0.43
1:B:1597:GLU:HB2	1:B:1600:GLU:HB2	2.01	0.43
1:B:1414:LEU:HD12	1:B:1414:LEU:HA	1.89	0.43
1:B:1479:MET:HA	1:B:1482:MET:HG2	2.01	0.42
1:B:1578:GLY:O	1:B:1612:LEU:HD13	2.19	0.42
1:B:1684:THR:O	1:B:1687:GLN:HB2	2.18	0.42
1:B:1699:GLU:HA	1:B:1702:GLU:HB2	2.00	0.42
1:B:1648:ALA:O	1:B:1650:LYS:N	2.51	0.42
1:B:1651:ARG:HH22	2:B:1801:GOL:C3	2.28	0.42
1:B:1486:LEU:HD21	1:B:1523:LEU:CD2	2.50	0.42
1:B:1704:ASP:O	1:B:1708:ILE:HG13	2.20	0.42
1:B:1502:LEU:HD11	1:B:1520:MET:CE	2.50	0.42
1:B:1491:ILE:HD11	1:B:1495:ARG:NH2	2.35	0.41
1:B:1667:ASP:O	1:B:1671:VAL:HG23	2.21	0.41
1:B:1696:LEU:HD12	1:B:1696:LEU:HA	1.88	0.41
1:B:1687:GLN:O	1:B:1690:PHE:HB3	2.21	0.40
1:B:1659:VAL:HG12	1:B:1674:LEU:CG	2.51	0.40
1:B:1659:VAL:HG12	1:B:1674:LEU:CD2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	272/572 (48%)	261 (96%)	10 (4%)	1 (0%)	34	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1663	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	238/502 (47%)	215 (90%)	23 (10%)	8	19

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1414	LEU
1	B	1418	GLN
1	B	1466	ARG
1	B	1487	GLN
1	B	1491	ILE
1	B	1510	GLU
1	B	1523	LEU
1	B	1531	GLU
1	B	1535	GLU
1	B	1538	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	1562	LYS
1	B	1570	PHE
1	B	1571	LYS
1	B	1575	LYS
1	B	1597	GLU
1	B	1604	ILE
1	B	1612	LEU
1	B	1645	VAL
1	B	1663	VAL
1	B	1667	ASP
1	B	1673	ASP
1	B	1684	THR
1	B	1719	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	1693	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	B	1801	-	5,5,5	0.30	0	5,5,5	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	1801	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1801	GOL	O1-C1-C2-C3
2	B	1801	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1801	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	276/572 (48%)	0.06	13 (4%) 31 30	19, 32, 85, 96	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1696	LEU	4.0
1	B	1711	VAL	3.6
1	B	1703	GLY	3.1
1	B	1708	ILE	3.0
1	B	1707	THR	2.7
1	B	1709	GLU	2.7
1	B	1718	TYR	2.6
1	B	1689	LYS	2.5
1	B	1415	ASP	2.4
1	B	1705	GLU	2.3
1	B	1691	PHE	2.3
1	B	1721	SER	2.1
1	B	1690	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	GOL	B	1801	6/6	0.83	0.15	42,44,51,52	0

6.5 Other polymers

There are no such residues in this entry.