



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 26, 2024 – 12:13 PM EDT

PDB ID : 6C99
Title : Crystal structure of FcRn bound to UCB-303
Authors : Fox III, D.; Abendroth, J.; Porter, J.; Deboves, H.
Deposited on : 2018-01-25
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	274	<div> <div>0%</div> <div>89%</div> <div>8%</div> <div>.</div> </div>
1	C	274	<div> <div>9%</div> <div>87%</div> <div>7%</div> <div>5%</div> </div>
2	B	99	<div> <div>90%</div> <div>10%</div> </div>
2	D	99	<div> <div>2%</div> <div>91%</div> <div>9%</div> </div>

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6054 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called IgG receptor FcRn large subunit p51.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	265	Total	C	N	O	S	0	11	0
			2064	1320	351	382	11			
1	C	260	Total	C	N	O	S	0	4	0
			1941	1247	324	360	10			

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C	N	O	S	0	3	0
			818	521	138	156	3			
2	D	99	Total	C	N	O	S	0	0	0
			760	489	133	135	3			

- Molecule 3 is methyl 7-(3,5-difluorophenyl)-5-(pyridin-3-yl)[1,2,4]triazolo[1,5-a]pyrimidine-6-carboxylate (three-letter code: EQY) (formula: C₁₈H₁₁F₂N₅O₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0	0
			27	18	2	5	2		
3	C	1	Total	C	F	N	O	0	0
			27	18	2	5	2		

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is CITRIC ACID (three-letter code: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	C	O	0	0
			13	6	7		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	151	Total	O	0	0
			151	151		
7	B	98	Total	O	0	0
			98	98		
7	C	65	Total	O	0	0
			65	65		
7	D	46	Total	O	0	0
			46	46		

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	41.77Å 76.08Å 140.04Å 90.00° 93.56° 90.00°	Depositor
Resolution (Å)	46.59 – 2.00 46.59 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (46.59-2.00) 99.8 (46.59-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.64 (at 2.00Å)	Xtriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.189 , 0.220 0.191 , 0.223	Depositor DCC
R_{free} test set	2896 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	26.4	Xtriage
Anisotropy	0.497	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 54.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6054	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EQY, NAG, CIT, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/2160	0.57	0/2946
1	C	0.35	0/2013	0.53	0/2754
2	B	0.42	0/850	0.59	0/1157
2	D	0.34	0/783	0.53	0/1071
All	All	0.39	0/5806	0.55	0/7928

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2064	0	1944	13	0
1	C	1941	0	1763	11	0
2	B	818	0	757	7	0
2	D	760	0	678	5	0
3	A	27	0	0	0	0
3	C	27	0	0	0	0
4	A	14	0	13	0	0
5	A	18	0	24	1	0
5	B	12	0	16	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	C	13	0	5	0	0
7	A	151	0	0	3	0
7	B	98	0	0	2	0
7	C	65	0	0	0	0
7	D	46	0	0	1	0
All	All	6054	0	5200	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53:TRP:HB3	1:C:181[A]:SER:OG	1.91	0.70
1:C:96:CYS:SG	1:C:159[B]:CYS:HB3	2.34	0.67
2:D:15:ALA:O	7:D:101:HOH:O	2.13	0.66
1:A:96:CYS:SG	1:A:159[B]:CYS:HB3	2.37	0.64
1:A:71:LYS:NZ	7:A:402:HOH:O	2.32	0.63
1:A:82:LEU:O	7:A:401:HOH:O	2.16	0.62
2:D:33:SER:HB2	2:D:54:LEU:HD21	1.80	0.62
1:A:181[A]:SER:OG	1:C:53:TRP:HB3	2.03	0.59
2:D:25:CYS:HB2	2:D:39:LEU:HD21	1.89	0.54
2:B:99:MET:HG3	5:B:101:GOL:H12	1.91	0.53
2:B:34[B]:ASP:OD1	7:B:201:HOH:O	2.18	0.52
1:C:189:SER:HB2	1:C:195:VAL:HG23	1.91	0.52
1:A:183:ARG:HB3	1:A:185:LYS:HE3	1.92	0.51
1:A:48:CYS:SG	7:A:533:HOH:O	2.58	0.51
1:A:116:GLU:HG2	5:A:305:GOL:H11	1.92	0.51
2:B:89:GLN:NE2	7:B:202:HOH:O	2.21	0.50
2:B:33:SER:HB2	2:B:54:LEU:HD21	1.96	0.48
1:C:25:TRP:HA	1:C:39:ASN:HA	1.95	0.47
1:C:235:HIS:NE2	2:D:99:MET:OXT	2.46	0.47
1:A:235:HIS:NE2	2:B:99:MET:O	2.44	0.46
1:A:140:ARG:O	1:A:144:GLN:HG2	2.15	0.46
1:C:71:LYS:NZ	1:C:151:GLU:OE2	2.43	0.45
1:C:47:PRO:HB3	1:C:61:TRP:CZ2	2.51	0.45
1:C:94:LEU:HD23	1:C:108:ALA:HA	1.99	0.45
2:B:24:ASN:HB3	2:B:65:LEU:HD11	1.99	0.44
2:B:95:TRP:CH2	2:B:97:ARG:HG2	2.52	0.44
1:A:186:ALA:HB2	1:A:196:LEU:HD23	1.98	0.44
1:A:157:PHE:C	1:A:160:PRO:HD2	2.39	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:154:PHE:HA	1:C:158:SER:HB2	2.02	0.41
1:A:4:HIS:HA	1:A:5:LEU:HA	1.67	0.41
1:C:106:PRO:HG2	1:C:122:LEU:HD13	2.03	0.41
2:D:33:SER:HB3	2:D:62:PHE:CE2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	274/274 (100%)	265 (97%)	8 (3%)	1 (0%)	30	27
1	C	260/274 (95%)	255 (98%)	5 (2%)	0	100	100
2	B	100/99 (101%)	100 (100%)	0	0	100	100
2	D	97/99 (98%)	97 (100%)	0	0	100	100
All	All	731/746 (98%)	717 (98%)	13 (2%)	1 (0%)	48	47

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	145	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/226 (95%)	213 (99%)	2 (1%)	75	81
1	C	190/226 (84%)	187 (98%)	3 (2%)	58	64
2	B	90/94 (96%)	89 (99%)	1 (1%)	70	76
2	D	75/94 (80%)	73 (97%)	2 (3%)	40	42
All	All	570/640 (89%)	562 (99%)	8 (1%)	62	68

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	226	PHE
1	A	244	SER
2	B	70	PHE
1	C	117	PHE
1	C	122	LEU
1	C	244	SER
2	D	57	SER
2	D	70	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	A	304	-	5,5,5	1.02	0	5,5,5	1.19	1 (20%)
5	GOL	A	303	-	5,5,5	0.97	0	5,5,5	1.27	1 (20%)
4	NAG	A	302	1	14,14,15	0.33	0	17,19,21	0.43	0
5	GOL	B	102	-	5,5,5	0.98	0	5,5,5	1.22	1 (20%)
3	EQY	A	301	-	27,30,30	2.00	7 (25%)	33,43,43	4.55	14 (42%)
5	GOL	B	101	-	5,5,5	1.04	0	5,5,5	1.50	1 (20%)
5	GOL	A	305	-	5,5,5	0.74	0	5,5,5	1.11	0
6	CIT	C	302	-	12,12,12	1.05	0	17,17,17	1.73	4 (23%)
3	EQY	C	301	-	27,30,30	1.87	6 (22%)	33,43,43	4.47	10 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	304	-	-	2/4/4/4	-
5	GOL	A	303	-	-	2/4/4/4	-
4	NAG	A	302	1	-	2/6/23/26	0/1/1/1
5	GOL	B	102	-	-	2/4/4/4	-
3	EQY	A	301	-	-	7/14/14/14	0/4/4/4
5	GOL	B	101	-	-	0/4/4/4	-
5	GOL	A	305	-	-	2/4/4/4	-
6	CIT	C	302	-	-	10/16/16/16	-
3	EQY	C	301	-	-	9/14/14/14	0/4/4/4

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	301	EQY	C05-C13	4.63	1.51	1.37
3	A	301	EQY	C06-N07	4.49	1.38	1.32
3	C	301	EQY	C05-C13	4.44	1.51	1.37
3	C	301	EQY	C06-N07	4.38	1.38	1.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	301	EQY	C14-C13	4.37	1.53	1.49
3	C	301	EQY	C05-C03	-3.79	1.41	1.50
3	A	301	EQY	C05-C03	-3.50	1.42	1.50
3	A	301	EQY	C08-N09	-3.23	1.31	1.35
3	A	301	EQY	C13-N12	3.12	1.47	1.41
3	C	301	EQY	C08-N09	-3.07	1.31	1.35
3	C	301	EQY	C14-C13	3.05	1.52	1.49
3	C	301	EQY	C13-N12	2.82	1.47	1.41
3	A	301	EQY	C05-C06	-2.01	1.37	1.41

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	301	EQY	C05-C13-N12	-20.27	109.74	119.53
3	C	301	EQY	C05-C13-N12	-19.21	110.25	119.53
3	C	301	EQY	C05-C13-C14	-10.62	110.65	122.40
3	A	301	EQY	C05-C13-C14	-9.96	111.38	122.40
3	C	301	EQY	C14-C13-N12	-8.25	106.50	117.83
3	A	301	EQY	C14-C13-N12	-7.48	107.57	117.83
3	C	301	EQY	C13-C05-C03	-6.13	114.02	122.21
3	A	301	EQY	C13-C05-C03	-5.16	115.31	122.21
6	C	302	CIT	O6-C6-C3	4.76	122.28	113.14
3	C	301	EQY	O02-C03-C05	4.47	122.78	112.28
3	A	301	EQY	O02-C03-C05	4.16	122.05	112.28
3	A	301	EQY	C22-C06-N07	-3.30	111.37	115.63
3	C	301	EQY	C25-N26-C27	3.11	122.29	116.85
3	A	301	EQY	C21-C19-C18	-2.92	119.96	123.50
3	A	301	EQY	C18-C16-C15	-2.73	120.18	123.50
3	A	301	EQY	C19-C18-C16	2.66	119.96	116.08
3	A	301	EQY	C25-N26-C27	2.65	121.49	116.85
3	C	301	EQY	C18-C16-C15	-2.63	120.31	123.50
3	C	301	EQY	C21-C19-C18	-2.58	120.36	123.50
5	B	101	GOL	C3-C2-C1	-2.51	102.61	111.80
6	C	302	CIT	O2-C1-C2	2.32	121.70	114.35
3	A	301	EQY	F17-C16-C15	2.32	121.57	118.28
6	C	302	CIT	O2-C1-O1	-2.30	117.41	123.33
3	C	301	EQY	O02-C03-O04	-2.30	118.99	123.46
5	A	303	GOL	C3-C2-C1	-2.23	103.60	111.80
3	C	301	EQY	C19-C18-C16	2.22	119.32	116.08
3	A	301	EQY	C01-O02-C03	2.22	120.07	115.81
5	B	102	GOL	C3-C2-C1	-2.22	103.67	111.80
3	A	301	EQY	C24-C23-C22	-2.14	118.10	120.54

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	C	302	CIT	O4-C5-C4	2.13	121.09	114.35
5	A	304	GOL	C3-C2-C1	-2.13	103.99	111.80
3	A	301	EQY	O02-C03-O04	-2.13	119.32	123.46

There are no chirality outliers.

All (36) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	301	EQY	C05-C06-C22-C27
3	A	301	EQY	N07-C06-C22-C27
3	C	301	EQY	C05-C13-C14-C21
3	C	301	EQY	C05-C06-C22-C27
3	C	301	EQY	N07-C06-C22-C27
5	A	303	GOL	O1-C1-C2-C3
6	C	302	CIT	C1-C2-C3-O7
6	C	302	CIT	C1-C2-C3-C4
6	C	302	CIT	O7-C3-C4-C5
6	C	302	CIT	C6-C3-C4-C5
6	C	302	CIT	O7-C3-C6-O5
6	C	302	CIT	O7-C3-C6-O6
6	C	302	CIT	C4-C3-C6-O5
6	C	302	CIT	C4-C3-C6-O6
6	C	302	CIT	C1-C2-C3-C6
6	C	302	CIT	C2-C3-C4-C5
4	A	302	NAG	C8-C7-N2-C2
4	A	302	NAG	O7-C7-N2-C2
5	A	304	GOL	O1-C1-C2-C3
5	A	305	GOL	C1-C2-C3-O3
5	B	102	GOL	O1-C1-C2-C3
3	C	301	EQY	C05-C06-C22-C23
5	A	303	GOL	O1-C1-C2-O2
3	A	301	EQY	C05-C06-C22-C23
5	A	305	GOL	O2-C2-C3-O3
3	C	301	EQY	N07-C06-C22-C23
3	A	301	EQY	N07-C06-C22-C23
3	A	301	EQY	C05-C13-C14-C21
5	A	304	GOL	O1-C1-C2-O2
5	B	102	GOL	O1-C1-C2-O2
3	C	301	EQY	O02-C03-C05-C06
3	C	301	EQY	O04-C03-C05-C06
3	A	301	EQY	O02-C03-C05-C13
3	C	301	EQY	O02-C03-C05-C13

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	301	EQY	O04-C03-C05-C13
3	C	301	EQY	O04-C03-C05-C13


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	101	GOL	1	0
5	A	305	GOL	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	265/274 (96%)	-0.19	4 (1%) 71 70	12, 26, 43, 63	12 (4%)
1	C	260/274 (94%)	0.58	24 (9%) 16 15	18, 39, 63, 72	4 (1%)
2	B	99/99 (100%)	-0.37	0 100 100	10, 23, 37, 43	4 (4%)
2	D	99/99 (100%)	0.44	2 (2%) 64 63	25, 39, 53, 59	0
All	All	723/746 (96%)	0.15	30 (4%) 42 40	10, 31, 56, 72	20 (2%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	84	GLY	3.7
1	C	217	LEU	3.4
1	C	4	HIS	3.1
1	A	84	GLY	3.0
1	C	248	HIS	3.0
2	D	1	ILE	3.0
1	A	58	SER	2.9
1	C	195	VAL	2.8
1	C	245	GLY	2.6
1	C	144	GLN	2.5
1	A	4	HIS	2.3
1	C	242	VAL	2.3
1	C	249	HIS	2.3
1	C	218	ALA	2.3
1	C	267	LEU	2.3
1	C	191	PRO	2.2
1	C	268	GLU	2.2
1	C	20	GLY	2.2
1	C	57	VAL	2.2
1	C	193	PHE	2.2
1	C	58	SER	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	59	TRP	2.1
1	C	83	GLY	2.1
1	C	187	ARG	2.1
1	C	77	GLU	2.1
1	C	246	ASP	2.1
2	D	2	GLN	2.1
1	C	139	GLN	2.1
1	C	215	ASN	2.0
1	A	99	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	A	302	14/15	0.78	0.13	48,52,66,68	0
6	CIT	C	302	13/13	0.81	0.14	41,55,63,67	0
5	GOL	B	102	6/6	0.83	0.13	32,43,46,48	0
5	GOL	A	305	6/6	0.87	0.12	39,43,48,51	0
5	GOL	A	303	6/6	0.90	0.12	28,39,47,47	0
5	GOL	A	304	6/6	0.91	0.11	26,36,37,43	0
5	GOL	B	101	6/6	0.94	0.08	16,24,31,39	0
3	EQY	C	301	27/27	0.96	0.06	20,26,34,37	0
3	EQY	A	301	27/27	0.98	0.04	12,17,24,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around EQY C 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around EQY A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.