



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 16, 2025 – 10:16 AM EDT

PDB ID : 9CBO / pdb_00009cbo
Title : Methionine synthase from *Thermus thermophilus* HB8, Folate and Cobalamin Domains, Pre-Catalytic Conformation (Pre-Fol-on)
Authors : Yamada, K.; Mendoza, J.; Koutmos, M.
Deposited on : 2024-06-19
Resolution : 3.34 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
buster-report : 1.1.7 (2018)
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.44


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1325 (3.38-3.30)
Clashscore	180529	1376 (3.38-3.30)
Ramachandran outliers	177936	1376 (3.38-3.30)
Sidechain outliers	177891	1375 (3.38-3.30)
RSRZ outliers	164620	1325 (3.38-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	511	<div> <div>%</div> <div> <div></div> <div>82%</div> <div>15%</div> <div>.</div> </div> </div>
1	B	511	<div> <div>%</div> <div> <div></div> <div>84%</div> <div>14%</div> <div>.</div> </div> </div>
1	C	511	<div> <div></div> <div> <div></div> <div>82%</div> <div>16%</div> <div>.</div> </div> </div>
1	D	511	<div> <div></div> <div> <div></div> <div>82%</div> <div>14%</div> <div>..</div> </div> </div>
1	E	511	<div> <div></div> <div> <div></div> <div>83%</div> <div>14%</div> <div>.</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	511	 A horizontal bar chart showing the quality of the chain. The bar is divided into three segments: a green segment representing 84%, a yellow segment representing 13%, and a small orange segment at the end. A small black dot is visible at the far right end of the bar.

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 24532 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Methionine synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	510	Total	C	N	O	S	0	0	0
			3978	2536	696	731	15			
1	B	510	Total	C	N	O	S	0	0	0
			3978	2536	696	731	15			
1	C	510	Total	C	N	O	S	0	0	0
			3978	2536	696	731	15			
1	D	507	Total	C	N	O	S	0	0	0
			3955	2523	692	725	15			
1	E	509	Total	C	N	O	S	0	0	0
			3969	2531	694	729	15			
1	F	510	Total	C	N	O	S	0	0	0
			3978	2536	696	731	15			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	759	ALA	ASP	engineered mutation	UNP Q5SKM5
B	759	ALA	ASP	engineered mutation	UNP Q5SKM5
C	759	ALA	ASP	engineered mutation	UNP Q5SKM5
D	759	ALA	ASP	engineered mutation	UNP Q5SKM5
E	759	ALA	ASP	engineered mutation	UNP Q5SKM5
F	759	ALA	ASP	engineered mutation	UNP Q5SKM5

- Molecule 2 is CO-METHYLCOBALAMIN (CCD ID: COB) (formula: $C_{63}H_{91}CoN_{13}O_{14}P$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	Co	N	O	P	0	0
			92	63	1	13	14	1		
2	B	1	Total	C	Co	N	O	P	0	0
			92	63	1	13	14	1		
2	C	1	Total	C	Co	N	O	P	0	0
			92	63	1	13	14	1		
2	D	1	Total	C	Co	N	O	P	0	0
			92	63	1	13	14	1		
2	E	1	Total	C	Co	N	O	P	0	0
			92	63	1	13	14	1		
2	F	1	Total	C	Co	N	O	P	0	0
			92	63	1	13	14	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		
3	F	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	30	Total	O	0	0
			30	30		
4	B	19	Total	O	0	0
			19	19		
4	C	26	Total	O	0	0
			26	26		
4	D	16	Total	O	0	0
			16	16		
4	E	6	Total	O	0	0
			6	6		
4	F	11	Total	O	0	0
			11	11		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Methionine synthase




- Molecule 1: Methionine synthase




- Molecule 1: Methionine synthase






● Molecule 1: Methionine synthase

Chain D: 82% 14%




● Molecule 1: Methionine synthase

Chain E: 83% 14%



● Molecule 1: Methionine synthase

Chain F: 84% 13%



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	187.77Å 187.77Å 325.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	163.18 – 3.34 162.65 – 3.34	Depositor EDS
% Data completeness (in resolution range)	99.9 (163.18-3.34) 99.9 (162.65-3.34)	Depositor EDS
R_{merge}	0.24	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.47 (at 3.33Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R_{free}	0.173 , 0.216 0.179 , 0.217	Depositor DCC
R_{free} test set	4229 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	100.1	Xtriage
Anisotropy	0.339	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 117.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	24532	wwPDB-VP
Average B, all atoms (Å ²)	119.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: COB, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.93	3/4044 (0.1%)	1.50	35/5458 (0.6%)
1	B	0.88	2/4044 (0.0%)	1.41	25/5458 (0.5%)
1	C	0.89	1/4044 (0.0%)	1.46	39/5458 (0.7%)
1	D	0.87	3/4021 (0.1%)	1.47	50/5427 (0.9%)
1	E	0.79	0/4035	1.41	27/5446 (0.5%)
1	F	0.83	1/4044 (0.0%)	1.43	34/5458 (0.6%)
All	All	0.87	10/24232 (0.0%)	1.45	210/32705 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	7
1	B	0	6
1	C	0	11
1	D	0	8
1	E	0	8
1	F	0	9
All	All	0	49

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	652	PRO	N-CD	8.31	1.59	1.47
1	D	496	GLU	CD-OE2	7.33	1.39	1.25
1	D	496	GLU	CD-OE1	6.66	1.38	1.25
1	B	485	HIS	CG-CD2	-6.65	1.28	1.35
1	B	503	ARG	NE-CZ	6.62	1.40	1.33

The worst 5 of 210 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	508	ARG	NE-CZ-NH1	-10.17	111.33	121.50
1	D	546	PRO	N-CA-C	10.07	126.90	114.92
1	D	841	ARG	NE-CZ-NH1	-9.39	112.11	121.50
1	D	546	PRO	N-CA-CB	-8.40	93.35	102.17
1	C	657	ASP	CA-CB-CG	8.34	120.94	112.60

There are no chirality outliers.

5 of 49 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	503	ARG	Sidechain
1	A	508	ARG	Sidechain
1	A	524	ARG	Sidechain
1	A	561	ARG	Sidechain
1	A	581	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3978	0	4096	45	0
1	B	3978	0	4096	34	0
1	C	3978	0	4096	42	0
1	D	3955	0	4077	34	0
1	E	3969	0	4088	41	0
1	F	3978	0	4096	33	0
2	A	92	0	87	13	0
2	B	92	0	87	8	0
2	C	92	0	85	11	0
2	D	92	0	87	8	0
2	E	92	0	86	5	0
2	F	92	0	86	9	0
3	A	6	0	8	0	0
3	B	6	0	8	1	0
3	C	12	0	16	1	0
3	E	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	6	0	8	0	0
4	A	30	0	0	6	0
4	B	19	0	0	3	0
4	C	26	0	0	2	0
4	D	16	0	0	1	0
4	E	6	0	0	1	0
4	F	11	0	0	1	0
All	All	24532	0	25115	270	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 270 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:901:COB:H362	2:E:901:COB:H351	1.40	1.01
1:A:651:ASP:HB3	1:A:654:LYS:HB3	1.41	0.98
2:B:901:COB:H362	2:B:901:COB:H351	1.47	0.96
1:D:545:ARG:HB2	1:D:546:PRO:HD3	1.48	0.95
2:A:901:COB:H362	2:A:901:COB:H351	1.52	0.89

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	508/511 (99%)	496 (98%)	12 (2%)	0	100	100
1	B	508/511 (99%)	493 (97%)	15 (3%)	0	100	100
1	C	508/511 (99%)	491 (97%)	17 (3%)	0	100	100
1	D	505/511 (99%)	495 (98%)	10 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	507/511 (99%)	494 (97%)	12 (2%)	1 (0%)	44	72
1	F	508/511 (99%)	492 (97%)	15 (3%)	1 (0%)	44	72
All	All	3044/3066 (99%)	2961 (97%)	81 (3%)	2 (0%)	48	76

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	367	SER
1	F	367	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	414/414 (100%)	411 (99%)	3 (1%)	81	88
1	B	414/414 (100%)	410 (99%)	4 (1%)	73	84
1	C	414/414 (100%)	412 (100%)	2 (0%)	86	91
1	D	412/414 (100%)	405 (98%)	7 (2%)	56	75
1	E	413/414 (100%)	410 (99%)	3 (1%)	81	88
1	F	414/414 (100%)	407 (98%)	7 (2%)	56	75
All	All	2481/2484 (100%)	2455 (99%)	26 (1%)	73	84

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	793	LYS
1	E	738	GLU
1	F	793	LYS
1	E	553	LEU
1	E	854	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	402	GLN
1	F	774	ASN
1	C	573	ASN
1	C	761	HIS
1	D	692	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	COB	D	901	-	86,102,102	1.36	11 (12%)	128,170,170	1.93	28 (21%)
2	COB	B	901	1	86,102,102	1.33	9 (10%)	128,170,170	1.78	34 (26%)
3	GOL	C	903	-	5,5,5	0.20	0	5,5,5	0.55	0
3	GOL	F	902	-	5,5,5	0.33	0	5,5,5	1.01	0
2	COB	C	901	-	86,102,102	1.72	21 (24%)	128,170,170	2.29	36 (28%)
2	COB	E	901	1	86,102,102	1.42	10 (11%)	128,170,170	1.98	28 (21%)
3	GOL	A	902	-	5,5,5	0.73	0	5,5,5	1.11	0
3	GOL	E	902	-	5,5,5	0.47	0	5,5,5	1.58	1 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	C	902	-	5,5,5	1.60	1 (20%)	5,5,5	0.90	0
2	COB	A	901	1	86,102,102	1.44	17 (19%)	128,170,170	2.05	31 (24%)
3	GOL	B	902	-	5,5,5	0.36	0	5,5,5	0.84	0
2	COB	F	901	1	86,102,102	1.34	11 (12%)	128,170,170	1.86	29 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COB	D	901	-	-	13/52/231/231	0/3/11/11
2	COB	B	901	1	-	13/52/231/231	0/3/11/11
3	GOL	C	903	-	-	0/4/4/4	-
3	GOL	F	902	-	-	2/4/4/4	-
2	COB	C	901	-	-	12/52/231/231	0/3/11/11
2	COB	E	901	1	-	9/52/231/231	0/3/11/11
3	GOL	A	902	-	-	2/4/4/4	-
3	GOL	E	902	-	-	2/4/4/4	-
3	GOL	C	902	-	-	1/4/4/4	-
2	COB	A	901	1	-	15/52/231/231	0/3/11/11
3	GOL	B	902	-	-	3/4/4/4	-
2	COB	F	901	1	-	13/52/231/231	0/3/11/11

The worst 5 of 80 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	901	COB	C36-C7	-6.13	1.44	1.54
2	C	901	COB	C1-C19	-4.92	1.45	1.51
2	C	901	COB	C8B-C9B	4.41	1.49	1.40
2	D	901	COB	C43-N45	4.30	1.46	1.32
2	A	901	COB	O6R-C1R	4.04	1.46	1.40

The worst 5 of 187 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	901	COB	C20-C1-C19	-11.03	95.77	109.42
2	D	901	COB	C20-C1-C19	-7.66	99.93	109.42
2	E	901	COB	C20-C1-C19	-7.36	100.31	109.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	901	COB	C12-C11-C10	-6.94	117.31	123.54
2	B	901	COB	C20-C1-C19	-6.73	101.09	109.42

There are no chirality outliers.

5 of 85 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	901	COB	C1-C2-C26-C27
2	A	901	COB	C25-C2-C26-C27
2	A	901	COB	C3-C2-C26-C27
2	A	901	COB	C54-C17-C55-C56
2	A	901	COB	C18-C17-C55-C56



There are no ring outliers.



8 monomers are involved in 56 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	901	COB	8	0
2	B	901	COB	8	0
3	C	903	GOL	1	0
2	C	901	COB	11	0
2	E	901	COB	5	0
2	A	901	COB	13	0
3	B	902	GOL	1	0
2	F	901	COB	9	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	510/511 (99%)	-0.58	3 (0%) 85 79	65, 106, 183, 249	0
1	B	510/511 (99%)	-0.70	3 (0%) 85 79	73, 106, 169, 204	0
1	C	510/511 (99%)	-0.77	1 (0%) 92 90	68, 100, 154, 223	0
1	D	507/511 (99%)	-0.79	0 100 100	72, 107, 157, 188	0
1	E	509/511 (99%)	-0.61	2 (0%) 89 85	91, 146, 196, 233	0
1	F	510/511 (99%)	-0.74	0 100 100	80, 117, 160, 250	0
All	All	3056/3066 (99%)	-0.70	9 (0%) 90 87	65, 113, 179, 250	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	651	ASP	6.2
1	A	650	GLU	4.2
1	B	695	LEU	2.7
1	C	648	HIS	2.6
1	B	715	ALA	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	C	902	6/6	0.70	0.20	84,93,185,197	0
3	GOL	F	902	6/6	0.70	0.17	132,162,180,194	0
3	GOL	E	902	6/6	0.87	0.24	96,106,117,133	0
3	GOL	A	902	6/6	0.92	0.17	95,103,125,140	0
3	GOL	C	903	6/6	0.94	0.14	106,111,141,148	0
2	COB	A	901	92/92	0.95	0.10	88,132,171,185	0
3	GOL	B	902	6/6	0.95	0.10	76,97,131,158	0
2	COB	B	901	92/92	0.97	0.08	86,111,138,160	0
2	COB	E	901	92/92	0.98	0.07	82,104,131,144	0
2	COB	F	901	92/92	0.98	0.07	74,90,119,129	0
2	COB	C	901	92/92	0.98	0.07	60,75,95,101	0
2	COB	D	901	92/92	0.98	0.07	61,82,101,125	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around COB A 901:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around COB B 901:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around COB E 901:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around COB F 901:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around COB C 901:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around COB D 901:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.