



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 21, 2025 – 11:33 PM EDT

PDB ID : 6CF6 / pdb\_00006cf6  
Title : RNF146 TBM-Tankyrase ARC2-3 complex  
Authors : Da Rosa, P.A.; Xu, W.  
Deposited on : 2018-02-13  
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1306 (1.94-1.94)
Clashscore	180529	1400 (1.94-1.94)
Ramachandran outliers	177936	1387 (1.94-1.94)
Sidechain outliers	177891	1387 (1.94-1.94)
RSRZ outliers	164620	1306 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	348	 3% 90% 8%
1	B	348	 9% 89% 8%
2	C	13	 38% 92% 8%
2	D	13	 23% 77% 23%

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5338 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tankyrase-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	320	Total	C	N	O	S	0	0	0
			2445	1527	450	454	14			
1	B	319	Total	C	N	O	S	0	2	0
			2453	1532	452	454	15			

- Molecule 2 is a protein called RNF146.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	12	Total	C	N	O	0	0	0
			78	42	15	21			
2	D	10	Total	C	N	O	0	0	0
			67	36	13	18			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	157	Total	O	0	0
			157	157		
3	B	128	Total	O	0	0
			128	128		
3	C	2	Total	O	0	0
			2	2		
3	D	8	Total	O	0	0
			8	8		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Tankyrase-1




#### • Molecule 1: Tankyrase-1



#### • Molecule 2: RNF146



#### • Molecule 2: RNF146



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	134.02Å 103.28Å 75.16Å 90.00° 106.93° 90.00°	Depositor
Resolution (Å)	50.00 – 1.93 50.00 – 1.93	Depositor EDS
% Data completeness (in resolution range)	97.0 (50.00-1.93) 97.0 (50.00-1.93)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.05 (at 1.94Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, $R_{free}$	0.195 , 0.228 0.203 , 0.235	Depositor DCC
$R_{free}$ test set	3620 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.7	Xtriage
Anisotropy	0.150	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 31.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5338	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.49	0/2489	0.69	0/3367
1	B	0.50	0/2501	0.67	2/3384 (0.1%)
2	C	0.43	0/77	0.72	0/102
2	D	0.47	0/66	0.77	0/87
All	All	0.50	0/5133	0.68	2/6940 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	590	ARG	NE-CZ-NH1	5.68	123.14	120.30
1	B	530	ARG	NE-CZ-NH1	5.17	122.88	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2445	0	2470	5	0
1	B	2453	0	2476	7	0
2	C	78	0	61	0	0
2	D	67	0	54	0	0
3	A	157	0	0	0	0
3	B	128	0	0	0	0
3	C	2	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	8	0	0	0	0
All	All	5338	0	5061	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:567:MET:CE	1:B:602:LEU:HD23	2.25	0.66
1:A:493:ALA:HB3	1:B:527[A]:HIS:HB3	1.81	0.62
1:B:567:MET:HE2	1:B:602:LEU:HD23	1.89	0.55
1:B:527[A]:HIS:CG	1:B:528:PRO:HD2	2.45	0.52
1:B:567:MET:HE3	1:B:602:LEU:HD23	1.93	0.51
1:B:441:VAL:HG12	1:B:473:LEU:HD13	1.94	0.49
1:A:493:ALA:HB3	1:B:527[B]:HIS:HB2	1.96	0.47
1:A:468:ALA:O	1:A:474:ARG:NH1	2.51	0.44
1:A:348:LEU:HD11	1:A:353:VAL:HG23	2.01	0.42
1:A:562:ALA:HB2	1:A:596:HIS:CG	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	318/348 (91%)	318 (100%)	0	0	100	100
1	B	319/348 (92%)	319 (100%)	0	0	100	100
2	C	10/13 (77%)	10 (100%)	0	0	100	100
2	D	8/13 (62%)	8 (100%)	0	0	100	100
All	All	655/722 (91%)	655 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	260/283 (92%)	260 (100%)	0	100	100
1	B	262/283 (93%)	262 (100%)	0	100	100
2	C	7/9 (78%)	7 (100%)	0	100	100
2	D	6/9 (67%)	6 (100%)	0	100	100
All	All	535/584 (92%)	535 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	389	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	320/348 (91%)	0.22	12 (3%) 44 49	20, 34, 52, 78	0
1	B	319/348 (91%)	0.66	31 (9%) 15 17	14, 40, 68, 100	2 (0%)
2	C	12/13 (92%)	1.83	5 (41%) 1 0	58, 74, 80, 81	0
2	D	10/13 (76%)	1.14	3 (30%) 1 1	45, 53, 64, 65	0
All	All	661/722 (91%)	0.48	51 (7%) 21 23	14, 37, 64, 100	2 (0%)

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	317	PRO	6.2
1	A	315	ALA	5.2
1	B	322	VAL	5.0
1	B	318	SER	4.7
1	B	498	VAL	4.5
1	B	496	ALA	4.4
1	B	323	LEU	4.2
1	B	327	TYR	4.2
2	C	191	LEU	4.1
1	B	321	ALA	3.9
1	B	319	ALA	3.7
1	B	497	LYS	3.7
1	B	324	THR	3.4
1	B	501	THR	3.3
2	D	192	ALA	3.3
1	B	504	LEU	3.2
1	A	318	SER	3.1
1	A	317	PRO	3.0
1	B	503	ALA	3.0
1	B	481	PHE	3.0
1	B	474	ARG	2.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	634	GLU	2.8
1	B	316	ASP	2.8
2	C	200	ALA	2.8
1	B	500	LYS	2.8
1	B	328	LYS	2.7
1	A	327	TYR	2.7
1	B	489	ALA	2.6
1	A	323	LEU	2.6
1	B	502	LEU	2.6
1	B	602	LEU	2.5
2	C	192	ALA	2.5
1	A	321	ALA	2.5
1	A	634	GLU	2.5
2	C	202	SER	2.4
1	A	515	HIS	2.4
1	B	443	VAL	2.4
1	B	325	GLY	2.4
1	B	499	LYS	2.3
1	A	319	ALA	2.3
1	A	438	LYS	2.3
1	B	490	ALA	2.2
1	A	533	VAL	2.2
2	C	195	SER	2.2
1	B	551	PHE	2.2
1	A	404	SER	2.1
1	B	495	LEU	2.1
2	D	201	ASP	2.1
1	B	358	SER	2.1
1	B	320	LYS	2.1
2	D	195	SER	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.