



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 26, 2024 – 07:17 AM EDT

PDB ID : 1CGV
Title : SITE DIRECTED MUTATIONS OF THE ACTIVE SITE RESIDUE TYROSINE 195 OF CYCLODEXTRIN GLYCOSYLTRANSFERASE FROM BACILLUS CIRCULANS STRAIN 251 AFFECTING ACTIVITY AND PRODUCT SPECIFICITY
Authors : Strokopytov, B.V.; Dijkstra, B.W.
Deposited on : 1994-08-05
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	6282 (2.50-2.50)
Ramachandran outliers	177936	6191 (2.50-2.50)
Sidechain outliers	177891	6193 (2.50-2.50)
RSRZ outliers	164620	5504 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	686	
2	B	2	
2	C	2	
2	D	2	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5505 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYCLOMALTODEXTRIN GLUCANOTRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	686	Total	C	N	O	S	0	0	0
			5263	3321	900	1026	16			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	195	PHE	TYR	engineered mutation	UNP P43379

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	B	2	Total	C	O	0	0	0
			23	12	11			
2	C	2	Total	C	O	0	0	0
			23	12	11			
2	D	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Ca	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	171	Total 171	O 171	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: CYCLOMALTODEXTRIN GLUCANOTRANSFERASE



• Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



• Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



• Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	121.03Å 111.11Å 66.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.50 20.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) (20.00-2.50) 89.2 (20.00-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.00 (at 2.51Å)	Xtriage
Refinement program	TNT	Depositor
R, R_{free}	0.155 , (Not available) 0.140 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	24.1	Xtriage
Anisotropy	0.044	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 79.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5505	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	0/5393	1.31	68/7350 (0.9%)

There are no bond length outliers.

All (68) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	331	ARG	NE-CZ-NH2	-17.06	111.77	120.30
1	A	331	ARG	NE-CZ-NH1	13.69	127.14	120.30
1	A	290	ARG	NE-CZ-NH1	12.78	126.69	120.30
1	A	63	ASP	CB-CG-OD2	-12.39	107.15	118.30
1	A	290	ARG	NE-CZ-NH2	-12.03	114.29	120.30
1	A	63	ASP	CB-CG-OD1	10.59	127.83	118.30
1	A	196	ASP	CB-CG-OD1	9.05	126.44	118.30
1	A	412	ARG	NE-CZ-NH1	8.99	124.80	120.30
1	A	196	ASP	CB-CG-OD2	-8.64	110.52	118.30
1	A	104	ASP	CB-CG-OD1	8.42	125.87	118.30
1	A	585	ASP	CB-CG-OD2	-8.39	110.75	118.30
1	A	170	ASP	CB-CG-OD2	-8.22	110.90	118.30
1	A	617	ASP	CB-CG-OD1	8.08	125.57	118.30
1	A	313	ASP	CB-CG-OD2	-7.74	111.34	118.30
1	A	436	ARG	NE-CZ-NH1	7.58	124.09	120.30
1	A	328	ASP	CB-CG-OD2	-7.51	111.54	118.30
1	A	213	ASP	CB-CG-OD2	-7.45	111.59	118.30
1	A	104	ASP	CB-CG-OD2	-7.43	111.61	118.30
1	A	617	ASP	CB-CG-OD2	-7.43	111.62	118.30
1	A	147	ASP	CB-CG-OD2	-7.19	111.83	118.30
1	A	103	ARG	NE-CZ-NH1	7.11	123.86	120.30
1	A	373	ASP	CB-CG-OD2	-7.04	111.96	118.30
1	A	208	ASP	CB-CG-OD1	6.92	124.53	118.30
1	A	199	ASP	CB-CG-OD2	-6.92	112.07	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	159	ASP	CB-CG-OD2	-6.80	112.18	118.30
1	A	220	ASP	CB-CG-OD2	-6.75	112.23	118.30
1	A	53	ASP	CB-CG-OD1	6.67	124.30	118.30
1	A	208	ASP	CB-CG-OD2	-6.58	112.38	118.30
1	A	3	ASP	CB-CG-OD1	6.55	124.19	118.30
1	A	577	ASP	CB-CG-OD2	-6.47	112.48	118.30
1	A	318	ASP	CB-CG-OD2	-6.29	112.64	118.30
1	A	27	ASP	CB-CG-OD2	-6.21	112.71	118.30
1	A	540	ASP	CB-CG-OD2	-6.15	112.77	118.30
1	A	325	ASP	CB-CG-OD1	6.12	123.81	118.30
1	A	147	ASP	CB-CG-OD1	6.04	123.73	118.30
1	A	543	SER	N-CA-CB	6.02	119.53	110.50
1	A	430	ALA	N-CA-CB	6.01	118.52	110.10
1	A	318	ASP	CB-CG-OD1	5.87	123.58	118.30
1	A	229	ASP	CB-CG-OD2	-5.77	113.11	118.30
1	A	213	ASP	CB-CG-OD1	5.76	123.48	118.30
1	A	313	ASP	CB-CG-OD1	5.74	123.47	118.30
1	A	325	ASP	CB-CG-OD2	-5.74	113.13	118.30
1	A	159	ASP	CB-CG-OD1	5.73	123.46	118.30
1	A	412	ARG	N-CA-CB	5.72	120.90	110.60
1	A	585	ASP	CB-CG-OD1	5.71	123.44	118.30
1	A	282	ASP	CB-CG-OD2	-5.66	113.20	118.30
1	A	577	ASP	CB-CG-OD1	5.65	123.39	118.30
1	A	328	ASP	CB-CG-OD1	5.63	123.36	118.30
1	A	170	ASP	CB-CG-OD1	5.59	123.33	118.30
1	A	336	ASN	N-CA-CB	-5.57	100.58	110.60
1	A	3	ASP	CB-CG-OD2	-5.50	113.35	118.30
1	A	353	ARG	NE-CZ-NH1	5.47	123.03	120.30
1	A	639	ASP	CB-CG-OD2	-5.46	113.39	118.30
1	A	182	ASP	CB-CG-OD2	-5.41	113.43	118.30
1	A	639	ASP	CB-CG-OD1	5.40	123.16	118.30
1	A	220	ASP	CB-CG-OD1	5.39	123.15	118.30
1	A	398	ARG	NE-CZ-NH1	5.36	122.98	120.30
1	A	436	ARG	NE-CZ-NH2	-5.32	117.64	120.30
1	A	24	ARG	NE-CZ-NH1	5.32	122.96	120.30
1	A	53	ASP	CB-CG-OD2	-5.31	113.52	118.30
1	A	199	ASP	CB-CG-OD1	5.30	123.07	118.30
1	A	282	ASP	CB-CG-OD1	5.29	123.06	118.30
1	A	590	ARG	NE-CZ-NH1	5.29	122.94	120.30
1	A	540	ASP	CB-CG-OD1	5.28	123.05	118.30
1	A	224	ASP	CB-CG-OD2	-5.12	113.69	118.30
1	A	23	ASP	CB-CG-OD2	-5.11	113.70	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	417	ASP	CB-CG-OD1	5.07	122.86	118.30
1	A	319	ASP	CB-CG-OD1	5.03	122.83	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5263	0	5025	73	0
2	B	23	0	21	2	0
2	C	23	0	21	0	0
2	D	23	0	21	2	0
3	A	2	0	0	0	0
4	A	171	0	0	0	0
All	All	5505	0	5088	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:88:ASN:HD21	1:A:91:GLY:H	1.08	0.90
1:A:560:ASN:HD21	1:A:578:ASN:HA	1.37	0.89
1:A:670:PHE:CE2	1:A:680:ILE:HD11	2.20	0.77
1:A:560:ASN:ND2	1:A:578:ASN:HA	2.01	0.74
1:A:643:PRO:HB2	1:A:646:LYS:HG3	1.70	0.73
1:A:333:HIS:HB2	1:A:342:LEU:HD12	1.72	0.72
1:A:142:SER:HB2	1:A:143:PRO:HD2	1.72	0.72
1:A:88:ASN:HD21	1:A:91:GLY:N	1.88	0.69
1:A:227:ARG:HG2	1:A:255:PHE:CE2	2.26	0.69
2:D:1:GLC:H62	2:D:2:GLC:O5	1.97	0.64
1:A:163:LEU:HD22	1:A:164:LEU:N	2.13	0.64
1:A:88:ASN:ND2	1:A:91:GLY:H	1.90	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:260:LEU:HB2	1:A:283:PHE:HB3	1.82	0.61
1:A:204:ASN:OD1	1:A:206:THR:HB	2.03	0.59
2:D:1:GLC:H62	2:D:2:GLC:C5	2.32	0.59
1:A:142:SER:HB2	1:A:143:PRO:CD	2.33	0.57
1:A:603:ASN:HB3	1:A:624:PRO:HB3	1.87	0.57
1:A:293:PHE:O	1:A:341:LYS:HD2	2.05	0.56
1:A:48:LEU:HD12	1:A:95:THR:HG23	1.89	0.54
1:A:147:ASP:C	1:A:149:PRO:HD3	2.27	0.54
1:A:414:ILE:HG12	1:A:415:ASN:N	2.23	0.54
1:A:148:GLN:N	1:A:149:PRO:HD3	2.23	0.53
1:A:330:GLU:HB3	1:A:369:GLY:HA2	1.91	0.53
1:A:511:PRO:HB3	1:A:555:ALA:HA	1.92	0.52
1:A:34:PRO:HG2	1:A:49:TYR:CG	2.44	0.52
1:A:163:LEU:HD22	1:A:165:GLY:N	2.26	0.51
1:A:669:THR:HG22	1:A:670:PHE:N	2.27	0.50
1:A:26:SER:O	1:A:56:GLY:HA3	2.12	0.49
1:A:227:ARG:HH11	1:A:257:GLU:HB2	1.77	0.49
1:A:417:ASP:O	1:A:436:ARG:HG3	2.12	0.49
1:A:300:MET:HB2	1:A:415:ASN:O	2.13	0.48
1:A:80:VAL:HA	1:A:107:LYS:O	2.13	0.48
1:A:562:ASN:HB3	1:A:575:VAL:HG13	1.95	0.48
1:A:333:HIS:HB2	1:A:342:LEU:CD1	2.40	0.48
1:A:413:TRP:CE2	2:B:1:GLC:H2	2.49	0.47
1:A:642:VAL:HB	1:A:643:PRO:HD2	1.96	0.47
1:A:227:ARG:NH1	1:A:257:GLU:HB2	2.30	0.47
1:A:607:THR:HG22	1:A:653:LEU:HD21	1.95	0.47
1:A:194:LEU:HD23	1:A:194:LEU:HA	1.66	0.46
1:A:340:ARG:O	1:A:340:ARG:HG3	2.15	0.46
1:A:304:LYS:O	1:A:308:GLU:HG2	2.14	0.46
1:A:651:LYS:HE3	1:A:663:GLU:O	2.14	0.46
1:A:60:LYS:HE2	1:A:381:PHE:CD2	2.50	0.46
1:A:594:ASN:HB2	1:A:683:ASN:OD1	2.16	0.46
1:A:29:ASN:HA	1:A:30:PRO:HD3	1.64	0.45
1:A:378:ILE:HG13	1:A:379:PRO:HD2	1.98	0.45
1:A:60:LYS:HA	1:A:60:LYS:HD2	1.64	0.45
1:A:283:PHE:O	1:A:287:GLN:HG2	2.16	0.44
1:A:633:TYR:CD2	1:A:634:PRO:HA	2.52	0.44
1:A:87:ILE:HG21	1:A:89:TYR:CE2	2.52	0.44
1:A:87:ILE:HD13	1:A:143:PRO:HG2	2.00	0.43
1:A:630:VAL:HG12	1:A:631:TYR:CE1	2.53	0.43
1:A:395:ALA:N	1:A:396:PRO:CD	2.82	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:562:ASN:HB3	1:A:575:VAL:CG1	2.48	0.42
1:A:371:ASP:HA	1:A:372:PRO:HA	1.71	0.42
1:A:397:LEU:HA	1:A:400:CYS:SG	2.60	0.42
1:A:1:ALA:HB1	1:A:2:PRO:HD2	2.02	0.42
1:A:617:ASP:OD1	1:A:618:PRO:HD2	2.19	0.42
1:A:633:TYR:CG	1:A:634:PRO:HA	2.55	0.42
1:A:87:ILE:CD1	1:A:143:PRO:HG2	2.50	0.41
1:A:664:GLY:O	1:A:685:GLN:HB2	2.21	0.41
1:A:235:PRO:O	1:A:239:GLN:HG3	2.21	0.41
1:A:168:THR:O	1:A:169:ASN:HB2	2.20	0.41
1:A:149:PRO:HA	1:A:154:ASN:ND2	2.35	0.41
1:A:266:SER:HA	1:A:267:PRO:HD3	1.83	0.41
1:A:290:ARG:NH2	1:A:330:GLU:O	2.53	0.41
1:A:617:ASP:OD2	1:A:620:LYS:HE2	2.21	0.41
1:A:262:VAL:H	1:A:262:VAL:HG23	1.68	0.41
1:A:361:GLY:HA3	1:A:366:MET:SD	2.60	0.41
1:A:395:ALA:HB3	1:A:396:PRO:HD3	2.02	0.41
1:A:452:PRO:HD2	1:A:456:TYR:OH	2.21	0.41
1:A:607:THR:HG22	1:A:653:LEU:CD2	2.50	0.41
1:A:479:ASN:N	1:A:479:ASN:ND2	2.70	0.40
1:A:413:TRP:CD1	2:B:1:GLC:H4	2.57	0.40
1:A:121:LEU:HD12	1:A:121:LEU:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	684/686 (100%)	656 (96%)	27 (4%)	1 (0%)	48 69

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	629	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	556/556 (100%)	528 (95%)	28 (5%)	20	41

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	PHE
1	A	48	LEU
1	A	75	TRP
1	A	92	VAL
1	A	103	ARG
1	A	107	LYS
1	A	146	SER
1	A	147	ASP
1	A	163	LEU
1	A	172	GLN
1	A	173	ASN
1	A	269	ASN
1	A	308	GLU
1	A	330	GLU
1	A	335	SER
1	A	353	ARG
1	A	375	ARG
1	A	412	ARG
1	A	436	ARG
1	A	438	LEU
1	A	443	SER
1	A	455	SER
1	A	525	SER
1	A	564	LYS
1	A	595	ASN
1	A	651	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	653	LEU
1	A	670	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	GLN
1	A	62	ASN
1	A	88	ASN
1	A	119	GLN
1	A	120	ASN
1	A	172	GLN
1	A	269	ASN
1	A	410	GLN
1	A	416	ASN
1	A	479	ASN
1	A	560	ASN
1	A	632	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	B	1	2	12,12,12	0.40	0	17,17,17	0.91	0
2	GLC	B	2	2	11,11,12	0.38	0	15,15,17	1.14	1 (6%)
2	GLC	C	1	2	12,12,12	0.36	0	17,17,17	1.05	1 (5%)
2	GLC	C	2	2	11,11,12	0.48	0	15,15,17	0.93	0
2	GLC	D	1	2	12,12,12	0.29	0	17,17,17	1.53	2 (11%)
2	GLC	D	2	2	11,11,12	0.32	0	15,15,17	1.88	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2	-	0/2/22/22	0/1/1/1
2	GLC	B	2	2	-	2/2/19/22	0/1/1/1
2	GLC	C	1	2	-	2/2/22/22	0/1/1/1
2	GLC	C	2	2	-	0/2/19/22	0/1/1/1
2	GLC	D	1	2	-	0/2/22/22	0/1/1/1
2	GLC	D	2	2	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	GLC	C1-O5-C5	5.90	120.09	112.19
2	D	1	GLC	O4-C4-C3	-3.70	101.65	110.38
2	D	1	GLC	O5-C1-C2	3.01	115.60	110.30
2	D	2	GLC	O5-C1-C2	3.01	117.97	110.79
2	C	1	GLC	O5-C5-C4	-2.97	104.34	109.70
2	B	2	GLC	C1-O5-C5	2.08	114.97	112.19

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	1	GLC	O5-C5-C6-O6
2	D	2	GLC	O5-C5-C6-O6
2	C	1	GLC	C4-C5-C6-O6
2	D	2	GLC	C4-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	B	2	GLC	C4-C5-C6-O6
2	B	2	GLC	O5-C5-C6-O6


There are no ring outliers.


3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1	GLC	2	0
2	B	1	GLC	2	0
2	D	2	GLC	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	686/686 (100%)	-0.98	0 100 100	10, 21, 44, 82	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	D	2	11/12	0.69	0.22	92,93,98,99	0
2	GLC	D	1	12/12	0.77	0.15	84,86,90,92	0
2	GLC	B	2	11/12	0.91	0.09	26,27,31,33	0
2	GLC	C	2	11/12	0.92	0.07	34,35,39,44	0
2	GLC	C	1	12/12	0.93	0.08	37,40,45,49	0
2	GLC	B	1	12/12	0.98	0.05	18,22,29,33	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain B:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	CA	A	691	1/1	0.99	0.06	20,20,20,20	0
3	CA	A	692	1/1	0.99	0.01	16,16,16,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.