



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 26, 2024 – 02:01 PM EDT

PDB ID : 6CJK  
Title : Anti HIV Fab 10A  
Authors : Hangartner, L.; Ward, A.B.; Wilson, I.A.; Oyen, D.  
Deposited on : 2018-02-26  
Resolution : 1.79 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	7108 (1.80-1.80)
Clashscore	180529	8162 (1.80-1.80)
Ramachandran outliers	177936	8077 (1.80-1.80)
Sidechain outliers	177891	8076 (1.80-1.80)
RSRZ outliers	164620	7108 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	217	<div> <div>96%</div> <div> <div></div> <div></div> <div></div> <div></div> </div> </div>
1	C	217	<div> <div>93%</div> <div>7%</div> </div>
2	B	223	<div> <div>2%</div> <div>91%</div> <div>6%</div> </div>
2	D	223	<div> <div>2%</div> <div>90%</div> <div>5%</div> </div>

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 13163 atoms, of which 6183 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Immunoglobulin Fab light chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	216	Total	C	H	N	O	S	10	6	0
			3176	1016	1543	267	341	9			
1	C	217	Total	C	H	N	O	S	0	5	0
			3172	1015	1549	263	336	9			

- Molecule 2 is a protein called Immunoglobulin Fab heavy chain.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	209	Total	C	H	N	O	S	0	7	0
			3086	984	1521	257	313	11			
2	D	213	Total	C	H	N	O	S	0	7	0
			3147	1004	1553	263	316	11			

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C H O 7 2 3 2	0	0
3	B	1	Total C H O 7 2 3 2	0	0
3	C	1	Total C H O 7 2 3 2	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	D	1	Total C H O 14 3 8 3	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	129	Total O 129 129	0	0
5	B	162	Total O 162 162	0	0
5	C	101	Total O 101 101	0	0
5	D	155	Total O 155 155	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Immunoglobulin Fab light chain




- Molecule 1: Immunoglobulin Fab light chain



- Molecule 2: Immunoglobulin Fab heavy chain



- Molecule 2: Immunoglobulin Fab heavy chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.17Å 63.56Å 72.65Å 67.99° 87.87° 76.73°	Depositor
Resolution (Å)	40.70 – 1.79 40.70 – 1.79	Depositor EDS
% Data completeness (in resolution range)	95.9 (40.70-1.79) 95.9 (40.70-1.79)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.36 (at 1.79Å)	Xtriage
Refinement program	PHENIX 1.12_2829	Depositor
R, $R_{free}$	0.163 , 0.206 0.164 , 0.206	Depositor DCC
$R_{free}$ test set	4000 reflections (5.20%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.3	Xtriage
Anisotropy	0.229	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.43 , 49.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	13163	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.34% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PCA, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.53	0/1666	0.66	0/2278
1	C	0.49	0/1657	0.62	0/2268
2	B	0.58	0/1594	0.67	0/2184
2	D	0.53	1/1639 (0.1%)	0.63	0/2250
All	All	0.53	1/6556 (0.0%)	0.64	0/8980

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	127	CYS	CB-SG	-5.29	1.73	1.81

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1633	1543	1533	3	0
1	C	1623	1549	1536	5	0
2	B	1565	1521	1510	4	0
2	D	1594	1553	1541	5	0
3	B	8	6	6	0	0
3	C	4	3	3	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	6	8	8	0	0
5	A	129	0	0	0	1
5	B	162	0	0	0	0
5	C	101	0	0	0	0
5	D	155	0	0	2	1
All	All	6980	6183	6137	15	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:117:LYS:NZ	5:D:403:HOH:O	2.39	0.54
1:A:12:GLU:HG2	1:A:107:LYS:HG3	1.90	0.53
1:C:6:GLN:HG3	1:C:23[B]:CYS:SG	2.48	0.53
1:A:137:ASN:OD1	2:B:164:ARG:NH2	2.41	0.53
1:A:137:ASN:OD1	2:B:164:ARG:NH1	2.44	0.51
2:B:160:THR:OG1	2:B:161:ASN:N	2.48	0.46
2:B:108:LEU:CD2	2:B:148[B]:GLU:HB2	2.46	0.46
2:D:108:LEU:HD21	2:D:148[B]:GLU:HB2	2.00	0.44
2:D:170[A]:ARG:HD2	2:D:176:TYR:CZ	2.54	0.42
2:D:147:PRO:HD2	2:D:199:ALA:CB	2.49	0.42
2:D:171:GLN:OE1	5:D:401:HOH:O	2.22	0.42
1:C:113:PRO:HB3	1:C:139:TYR:HB3	2.02	0.41
1:C:158:ILE:HA	1:C:177:THR:O	2.21	0.41
1:C:169:ASP:O	1:C:170:CYS:HB2	2.21	0.41
1:C:185:TYR:O	1:C:191:TYR:OH	2.37	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:310:HOH:O	5:D:415:HOH:O[1_564]	2.15	0.05



## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	219/217 (101%)	211 (96%)	7 (3%)	1 (0%)	25	14
1	C	220/217 (101%)	210 (96%)	10 (4%)	0	100	100
2	B	211/223 (95%)	209 (99%)	2 (1%)	0	100	100
2	D	218/223 (98%)	213 (98%)	5 (2%)	0	100	100
All	All	868/880 (99%)	843 (97%)	24 (3%)	1 (0%)	48	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	138	LYS

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	180/181 (99%)	177 (98%)	3 (2%)	56	47
1	C	178/181 (98%)	173 (97%)	5 (3%)	38	27
2	B	176/185 (95%)	176 (100%)	0	100	100
2	D	181/185 (98%)	179 (99%)	2 (1%)	70	65
All	All	715/732 (98%)	705 (99%)	10 (1%)	62	56

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	91	ASN
1	A	167	SER
1	A	207	ASN
1	C	10	SER
1	C	69	THR
1	C	91	ASN
1	C	183	THR
1	C	190	GLU
2	D	128	CYS
2	D	194	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PCA	D	3	2	7,8,9	2.10	2 (28%)	9,10,12	1.70	4 (44%)
2	PCA	B	3	2	7,8,9	2.45	2 (28%)	9,10,12	1.98	4 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCA	D	3	2	-	0/0/11/13	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCA	B	3	2	-	0/0/11/13	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	3	PCA	CD-N	5.12	1.47	1.34
2	D	3	PCA	CD-N	4.80	1.46	1.34
2	B	3	PCA	CA-N	3.59	1.50	1.46
2	D	3	PCA	CA-N	2.50	1.49	1.46

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3	PCA	OE-CD-CG	-3.19	121.02	126.72
2	B	3	PCA	CA-N-CD	-2.79	104.01	113.58
2	D	3	PCA	CA-N-CD	-2.60	104.69	113.58
2	D	3	PCA	CG-CD-N	2.33	114.10	108.39
2	B	3	PCA	CG-CD-N	2.32	114.07	108.39
2	B	3	PCA	CB-CA-N	2.32	109.63	103.24
2	D	3	PCA	CB-CA-N	2.27	109.48	103.24
2	D	3	PCA	OE-CD-CG	-2.18	122.83	126.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	GOL	D	301	-	5,5,5	0.39	0	5,5,5	0.52	0
3	ACT	B	302	-	3,3,3	0.78	0	3,3,3	1.19	0
3	ACT	B	301	-	3,3,3	0.76	0	3,3,3	1.24	0
3	ACT	C	301	-	3,3,3	0.73	0	3,3,3	1.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	D	301	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	215/217 (99%)	-0.31	3 (1%) 73 72	14, 36, 62, 103	5 (2%)
1	C	217/217 (100%)	-0.15	0 100 100	17, 39, 76, 104	5 (2%)
2	B	208/223 (93%)	-0.46	4 (1%) 66 64	11, 28, 65, 144	7 (3%)
2	D	212/223 (95%)	-0.33	4 (1%) 66 64	13, 35, 76, 105	6 (2%)
All	All	852/880 (96%)	-0.31	11 (1%) 74 74	11, 34, 72, 144	23 (2%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	160	THR	4.7
1	A	95	TYR	3.4
1	A	202	VAL	3.3
2	D	132	PRO	3.1
2	B	135	THR	3.0
1	A	199	THR	2.9
2	B	161	ASN	2.8
2	B	159	LEU	2.8
2	D	160	THR	2.8
2	D	210	PRO	2.2
2	D	131	THR	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	PCA	D	3	8/9	0.92	0.09	29,44,63,76	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PCA	B	3	8/9	0.95	0.08	28,45,65,79	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	ACT	B	301	4/4	0.86	0.12	35,42,58,62	0
3	ACT	C	301	4/4	0.89	0.13	40,58,70,70	0
3	ACT	B	302	4/4	0.90	0.14	35,60,72,72	0
4	GOL	D	301	6/6	0.95	0.08	28,37,63,76	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.