



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 22, 2025 – 01:07 AM EDT

PDB ID : 6CMN / pdb_00006cmn
Title : Co-Crystal Structure of HIV-1 TAR Bound to Lab-Evolved RRM TBP6.7
Authors : Belashov, I.A.; Wedekind, J.E.
Deposited on : 2018-03-05
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	2.0rc1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.006 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.42


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7108 (1.80-1.80)
Clashscore	180529	8162 (1.80-1.80)
Ramachandran outliers	177936	8077 (1.80-1.80)
Sidechain outliers	177891	8076 (1.80-1.80)
RSRZ outliers	164620	7108 (1.80-1.80)
RNA backbone	3690	1009 (2.24-1.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	119	<div> <div>11%</div> <div> <div></div> <div>76%</div> <div>24%</div> </div> </div>
2	D	27	<div> <div>96%</div> <div></div> </div>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2541 atoms, of which 1070 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TAR-Binding Protein 6.7.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	91	Total	C	H	N	O	S	0	1	0
			1523	480	777	135	129	2			

There are 33 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	expression tag	UNP G1TM83
A	-19	GLY	-	expression tag	UNP G1TM83
A	-18	SER	-	expression tag	UNP G1TM83
A	-17	SER	-	expression tag	UNP G1TM83
A	-16	HIS	-	expression tag	UNP G1TM83
A	-15	HIS	-	expression tag	UNP G1TM83
A	-14	HIS	-	expression tag	UNP G1TM83
A	-13	HIS	-	expression tag	UNP G1TM83
A	-12	HIS	-	expression tag	UNP G1TM83
A	-11	HIS	-	expression tag	UNP G1TM83
A	-10	SER	-	expression tag	UNP G1TM83
A	-9	SER	-	expression tag	UNP G1TM83
A	-8	GLY	-	expression tag	UNP G1TM83
A	-7	GLU	-	expression tag	UNP G1TM83
A	-6	ASN	-	expression tag	UNP G1TM83
A	-5	LEU	-	expression tag	UNP G1TM83
A	-4	TYR	-	expression tag	UNP G1TM83
A	-3	PHE	-	expression tag	UNP G1TM83
A	-2	GLN	-	expression tag	UNP G1TM83
A	-1	GLY	-	expression tag	UNP G1TM83
A	0	HIS	-	expression tag	UNP G1TM83
A	19	SER	GLU	engineered mutation	UNP G1TM83
A	31	HIS	TYR	engineered mutation	UNP G1TM83
A	36	ARG	GLN	engineered mutation	UNP G1TM83
A	46	PRO	SER	engineered mutation	UNP G1TM83
A	48	GLN	SER	engineered mutation	UNP G1TM83
A	49	ARG	LEU	engineered mutation	UNP G1TM83

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	50	THR	LYS	engineered mutation	UNP G1TM83
A	51	PRO	MET	engineered mutation	UNP G1TM83
A	83	ALA	ARG	engineered mutation	UNP G1TM83
A	91	LYS	SER	engineered mutation	UNP G1TM83
A	92	ARG	ASP	engineered mutation	UNP G1TM83
A	94	PRO	ILE	engineered mutation	UNP G1TM83

- Molecule 2 is a RNA chain called Trans-Activation Response RNA Element.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	D	27	Total	C	H	N	O	P	0	0	0
			865	256	293	101	189	26			



- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	64	Total	O	0	0
			64	64		
3	D	89	Total	O	0	0
			89	89		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: TAR-Binding Protein 6.7



- Molecule 2: Trans-Activation Response RNA Element



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	40.42Å 40.42Å 284.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.19 – 1.80 37.19 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.3 (37.19-1.80) 96.1 (37.19-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.97 (at 1.79Å)	Xtriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, R_{free}	0.189 , 0.221 0.194 , 0.223	Depositor DCC
R_{free} test set	22147 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	30.3	Xtriage
Anisotropy	0.009	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 53.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2541	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/765	0.54	0/1029
2	D	0.41	0/638	0.84	0/993
All	All	0.37	0/1403	0.70	0/2022

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	746	777	778	0	0
2	D	572	293	293	0	0
3	A	64	0	0	0	0
3	D	89	0	0	0	0
All	All	1471	1070	1071	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	90/119 (76%)	89 (99%)	1 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	82/105 (78%)	81 (99%)	1 (1%)	67	62

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	D	26/27 (96%)	1 (3%)	0

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	D	26	G

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	91/119 (76%)	0.67	13 (14%) 7 6	15, 36, 79, 121	1 (1%)
2	D	27/27 (100%)	0.09	0 100 100	34, 45, 65, 68	0
All	All	118/146 (80%)	0.54	13 (11%) 12 9	15, 39, 78, 121	1 (0%)

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	95	ALA	5.9
1	A	94	PRO	4.4
1	A	93	ILE	4.2
1	A	33	ILE	3.5
1	A	5	GLU	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.