



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 4, 2024 – 02:09 PM EST

PDB ID : 3D3W  
Title : Structure of L-Xylulose Reductase with bound coenzyme, phosphate and hydroxide.  
Authors : Zhao, H.-T.; El-Kabbani, O.  
Deposited on : 2008-05-12  
Resolution : 1.87 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.87 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1090 (1.88-1.88)
Clashscore	180529	1144 (1.88-1.88)
Ramachandran outliers	177936	1135 (1.88-1.88)
Sidechain outliers	177891	1135 (1.88-1.88)
RSRZ outliers	164620	1090 (1.88-1.88)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	244	<div> <div>9%</div> <div>86%</div> <div>12%</div> <div>.</div> </div>
2	B	245	<div> <div>5%</div> <div>83%</div> <div>15%</div> <div>..</div> </div>

2 Entry composition ⓘ

There are 5 unique types of molecules in this entry. The entry contains 4322 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-xylulose reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	244	1832	1153	327	340	12	0	4	0


- Molecule 2 is a protein called L-xylulose reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	244	1832	1152	328	339	13	0	4	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	138	CSO	CYS	microheterogeneity	UNP Q7Z4W1

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: C<sub>21</sub>H<sub>28</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
3	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	P	0	1
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	310	Total	O	0	0
			310	310		
5	B	247	Total	O	0	0
			247	247		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: L-xylulose reductase



- Molecule 2: L-xylulose reductase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	73.68Å 87.48Å 72.17Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.87 30.00 – 1.87	Depositor EDS
% Data completeness (in resolution range)	98.1 (30.00-1.87) 98.1 (30.00-1.87)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.38 (at 1.87Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.204 , 0.258 0.202 , 0.255	Depositor DCC
$R_{free}$ test set	1932 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.0	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 64.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.021 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4322	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 24.01 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.1691e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CSO, NAP, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.92	4/1865 (0.2%)	0.82	4/2532 (0.2%)
2	B	0.96	1/1862 (0.1%)	0.82	1/2528 (0.0%)
All	All	0.94	5/3727 (0.1%)	0.82	5/5060 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	166	GLU	CD-OE2	-5.67	1.19	1.25
2	B	166	GLU	CD-OE2	-5.46	1.19	1.25
1	A	166	GLU	CD-OE1	-5.43	1.19	1.25
1	A	105	GLU	CD-OE1	-5.39	1.19	1.25
1	A	105	GLU	CD-OE2	-5.30	1.19	1.25

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	47	LEU	CB-CG-CD2	5.95	121.12	111.00
1	A	173	ARG	NE-CZ-NH2	-5.89	117.36	120.30
1	A	203	ARG	NE-CZ-NH1	-5.66	117.47	120.30
1	A	69	ARG	NE-CZ-NH2	-5.23	117.68	120.30
2	B	139	SER	N-CA-CB	-5.12	102.81	110.50

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	138[B]	CSO	Mainchain
2	B	138[A]	CYS	Mainchain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1832	0	1892	30	0
2	B	1832	0	1888	39	0
3	A	48	0	25	3	0
3	B	48	0	25	2	0
4	B	5	0	0	1	0
5	A	310	0	0	4	0
5	B	247	0	0	7	0
All	All	4322	0	3830	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:37:VAL:HG22	2:B:57[B]:VAL:CG1	1.94	0.97
2:B:211:GLU:H	2:B:214:HIS:HD2	1.23	0.86
1:A:37:VAL:HG22	1:A:57[B]:VAL:CG1	2.06	0.85
2:B:48:VAL:HG23	5:B:1252:HOH:O	1.77	0.84
1:A:211:GLU:H	1:A:214:HIS:HD2	1.26	0.82
2:B:37:VAL:HG22	2:B:57[B]:VAL:HG13	1.63	0.81
2:B:203:ARG:HD2	5:B:1262:HOH:O	1.85	0.76
1:A:37:VAL:HG22	1:A:57[B]:VAL:HG13	1.73	0.71
2:B:48:VAL:CG2	5:B:1252:HOH:O	2.39	0.66
2:B:200:MET:HE3	2:B:209:PHE:CE1	2.31	0.65
2:B:33:ARG:HD2	5:B:1398:HOH:O	1.98	0.63
2:B:9:ARG:HH21	2:B:55:GLU:CD	2.06	0.59

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:14:GLY:HA2	3:B:1245:NAP:H1B	1.84	0.59
1:A:200:MET:HE3	1:A:209:PHE:CE1	2.38	0.58
1:A:9:ARG:HH21	1:A:55:GLU:CD	2.08	0.56
2:B:171:LYS:HE2	5:B:1269:HOH:O	2.05	0.56
1:A:89:LEU:HD12	1:A:190:THR:HG21	1.87	0.56
1:A:244:CYS:O	2:B:203:ARG:NH2	2.26	0.56
2:B:48:VAL:HG12	2:B:54:ILE:HG13	1.87	0.56
2:B:135:SER:O	3:B:1245:NAP:H6N	2.06	0.55
2:B:193:ASP:OD2	2:B:196:LYS:HD3	2.06	0.55
2:B:211:GLU:H	2:B:214:HIS:CD2	2.14	0.55
2:B:9:ARG:NH2	2:B:55:GLU:OE1	2.40	0.55
2:B:200:MET:CE	2:B:209:PHE:CE1	2.90	0.54
1:A:211:GLU:H	1:A:214:HIS:CD2	2.16	0.54
1:A:83:ASN:ND2	3:A:245:NAP:H4D	2.23	0.54
1:A:14:GLY:HA2	3:A:245:NAP:H1B	1.91	0.53
2:B:9:ARG:HD3	2:B:76:PRO:O	2.08	0.53
1:A:74:VAL:HG12	5:A:358:HOH:O	2.08	0.53
1:A:193:ASP:OD2	1:A:196:LYS:HD3	2.10	0.52
2:B:89:LEU:HD12	2:B:190:THR:HG21	1.92	0.51
2:B:37:VAL:HG22	2:B:57[B]:VAL:HG11	1.86	0.50
2:B:193:ASP:HB3	2:B:196:LYS:HB2	1.93	0.50
1:A:203:ARG:NH2	2:B:203:ARG:HD3	2.26	0.49
1:A:83:ASN:HD21	3:A:245:NAP:C5D	2.26	0.48
1:A:9:ARG:NH2	1:A:55:GLU:OE1	2.45	0.48
2:B:200:MET:HB3	2:B:200:MET:HE2	1.64	0.48
1:A:200:MET:CE	1:A:209:PHE:CE1	2.96	0.48
1:A:9:ARG:HD3	1:A:76:PRO:O	2.15	0.47
2:B:181:VAL:CG2	2:B:200:MET:HE3	2.44	0.47
2:B:10:VAL:HG22	2:B:79:LEU:HB2	1.97	0.46
1:A:195:HIS:HA	1:A:198:LYS:HB2	1.98	0.46
2:B:181:VAL:O	2:B:209:PHE:HB3	2.17	0.45
1:A:195:HIS:HA	1:A:198:LYS:CB	2.47	0.45
2:B:181:VAL:CG2	2:B:200:MET:CE	2.95	0.45
2:B:138[A]:CYS:HB2	4:B:245[A]:PO4:O2	2.18	0.44
1:A:200:MET:HB3	1:A:200:MET:HE2	1.69	0.43
2:B:30:THR:HG21	5:B:421:HOH:O	2.17	0.43
1:A:181:VAL:HG21	1:A:200:MET:CE	2.49	0.43
1:A:181:VAL:HG21	1:A:200:MET:HE1	2.00	0.43
1:A:55:GLU:HB3	5:A:335:HOH:O	2.19	0.43
2:B:181:VAL:HG21	2:B:200:MET:CE	2.50	0.42
1:A:45:ASP:O	1:A:48:VAL:HG22	2.19	0.42

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72:GLY:O	5:A:435:HOH:O	2.22	0.42
1:A:6:ALA:HB1	5:A:423:HOH:O	2.19	0.41
1:A:181:VAL:CG2	1:A:200:MET:HE3	2.49	0.41
2:B:78:ASP:HA	2:B:122:LEU:HD21	2.02	0.41
2:B:167:LEU:HB3	2:B:172:ILE:HB	2.02	0.41
2:B:177[A]:VAL:HG13	2:B:222:LEU:HD12	2.02	0.41
2:B:208:LYS:CE	5:B:1369:HOH:O	2.68	0.41
2:B:2:GLU:CD	2:B:4:PHE:H	2.24	0.41
1:A:181:VAL:CG2	1:A:200:MET:CE	2.99	0.41
1:A:181:VAL:O	1:A:209:PHE:HB3	2.21	0.41
2:B:200:MET:CE	2:B:209:PHE:HE1	2.32	0.40
1:A:180:THR:HB	1:A:238[B]:GLU:HA	2.03	0.40
2:B:181:VAL:HG21	2:B:200:MET:HE1	2.03	0.40
2:B:182:VAL:O	2:B:187:GLY:HA3	2.20	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	245/244 (100%)	238 (97%)	6 (2%)	1 (0%)	30	18
2	B	244/245 (100%)	236 (97%)	7 (3%)	1 (0%)	30	18
All	All	489/489 (100%)	474 (97%)	13 (3%)	2 (0%)	30	18

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	72	GLY
2	B	72	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/195 (102%)	192 (96%)	7 (4%)	31	15
2	B	199/196 (102%)	194 (98%)	5 (2%)	42	26
All	All	398/391 (102%)	386 (97%)	12 (3%)	38	19

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	THR
1	A	38	SER
1	A	47	LEU
1	A	196	LYS
1	A	201	LEU
1	A	238[A]	GLU
1	A	238[B]	GLU
2	B	30	THR
2	B	196	LYS
2	B	201	LEU
2	B	203	ARG
2	B	238	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	83	ASN
1	A	137	GLN
1	A	140	GLN
1	A	214	HIS
2	B	214	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CSO	A	138	-	3,6,7	1.11	0	1,6,8	1.34	0
2	CSO	B	138[B]	-	3,6,7	1.21	1 (33%)	1,6,8	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	138	-	-	0/1/5/7	-
2	CSO	B	138[B]	-	-	0/1/5/7	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	138[B]	CSO	O-C	2.09	1.27	1.20

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAP	A	245	-	46,52,52	1.80	8 (17%)	61,80,80	2.08	14 (22%)
3	NAP	B	1245	-	46,52,52	1.75	7 (15%)	61,80,80	2.05	14 (22%)
4	PO4	B	245[A]	-	4,4,4	1.02	0	6,6,6	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAP	A	245	-	-	11/31/67/67	0/5/5/5
3	NAP	B	1245	-	-	5/31/67/67	0/5/5/5

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	245	NAP	C2N-N1N	7.39	1.43	1.35
3	B	1245	NAP	C2N-N1N	6.74	1.42	1.35
3	B	1245	NAP	C6N-N1N	4.40	1.45	1.35
3	A	245	NAP	C6N-N1N	4.38	1.45	1.35
3	B	1245	NAP	C2A-N3A	3.46	1.37	1.32
3	B	1245	NAP	PA-O3	3.45	1.63	1.59
3	A	245	NAP	PN-O3	3.31	1.63	1.59
3	A	245	NAP	PA-O3	2.97	1.62	1.59
3	B	1245	NAP	PN-O3	2.93	1.62	1.59
3	A	245	NAP	P2B-O2B	2.73	1.64	1.59
3	B	1245	NAP	C2A-N1A	2.50	1.38	1.33
3	A	245	NAP	O4D-C1D	2.40	1.44	1.40
3	A	245	NAP	P2B-O3X	-2.40	1.45	1.54
3	A	245	NAP	C3N-C7N	2.10	1.53	1.50

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1245	NAP	C8A-N7A	-2.09	1.30	1.34

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	245	NAP	N3A-C2A-N1A	-6.86	119.36	128.67
3	B	1245	NAP	C5N-C4N-C3N	6.36	126.61	120.36
3	B	1245	NAP	O4B-C1B-N9A	6.35	117.17	108.75
3	A	245	NAP	C5N-C4N-C3N	6.30	126.56	120.36
3	B	1245	NAP	N3A-C2A-N1A	-5.74	120.88	128.67
3	A	245	NAP	O4B-C1B-N9A	4.53	114.75	108.75
3	B	1245	NAP	O3-PA-O1A	-3.89	98.99	110.70
3	B	1245	NAP	C6N-C5N-C4N	-3.85	113.89	119.45
3	A	245	NAP	C6N-C5N-C4N	-3.72	114.09	119.45
3	A	245	NAP	C5A-C6A-N6A	3.71	125.96	120.31
3	B	1245	NAP	O3D-C3D-C2D	3.39	122.70	111.82
3	A	245	NAP	C2B-C1B-N9A	-3.36	105.10	112.56
3	A	245	NAP	O3-PA-O1A	-3.17	101.15	110.70
3	A	245	NAP	C4A-C5A-N7A	3.14	112.65	109.34
3	B	1245	NAP	O2A-PA-O1A	3.13	126.98	112.44
3	A	245	NAP	O2A-PA-O1A	3.06	126.67	112.44
3	A	245	NAP	O4B-C1B-C2B	-2.93	101.60	106.61
3	B	1245	NAP	C2B-C1B-N9A	-2.89	106.14	112.56
3	A	245	NAP	O3D-C3D-C2D	2.59	120.13	111.82
3	A	245	NAP	O2X-P2B-O2B	-2.55	95.91	105.85
3	B	1245	NAP	O2N-PN-O3	2.52	114.08	107.27
3	A	245	NAP	O3X-P2B-O2X	2.50	117.17	107.80
3	A	245	NAP	C5A-C6A-N1A	-2.45	114.40	120.23
3	B	1245	NAP	O7N-C7N-N7N	2.21	125.81	122.62
3	B	1245	NAP	O3X-P2B-O2X	2.15	115.85	107.80
3	B	1245	NAP	C5A-C6A-N6A	2.12	123.54	120.31
3	B	1245	NAP	C1B-N9A-C4A	-2.08	122.99	126.64
3	B	1245	NAP	C6N-N1N-C2N	-2.02	120.16	121.88

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	245	NAP	C5D-O5D-PN-O3
3	A	245	NAP	C5D-O5D-PN-O1N
3	A	245	NAP	C5D-O5D-PN-O2N
3	A	245	NAP	O4D-C1D-N1N-C2N

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
3	B	1245	NAP	C5D-O5D-PN-O3
3	B	1245	NAP	C5D-O5D-PN-O1N
3	B	1245	NAP	C5D-O5D-PN-O2N
3	A	245	NAP	C2N-C3N-C7N-O7N
3	A	245	NAP	C4N-C3N-C7N-N7N
3	A	245	NAP	C2N-C3N-C7N-N7N
3	A	245	NAP	C4N-C3N-C7N-O7N
3	A	245	NAP	C2B-O2B-P2B-O3X
3	A	245	NAP	O4D-C1D-N1N-C6N
3	B	1245	NAP	O4D-C1D-N1N-C2N
3	B	1245	NAP	O4D-C1D-N1N-C6N
3	A	245	NAP	O4B-C4B-C5B-O5B

There are no ring outliers.


3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	245	NAP	3	0
3	B	1245	NAP	2	0
4	B	245[A]	PO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	243/244 (99%)	0.19	23 (9%) 15 18	11, 21, 52, 73	4 (1%)
2	B	243/245 (99%)	0.19	13 (5%) 33 34	11, 22, 52, 73	3 (1%)
All	All	486/489 (99%)	0.19	36 (7%) 22 23	11, 22, 55, 73	7 (1%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	189	ALA	4.4
1	A	191	TRP	4.0
1	A	190	THR	3.8
1	A	186	MET	3.7
1	A	184	THR	3.7
2	B	190	THR	3.7
2	B	191	TRP	3.4
2	B	189	ALA	3.4
1	A	187	GLY	3.3
1	A	196	LYS	3.1
1	A	183	MET	3.0
1	A	182	VAL	2.9
1	A	194	PRO	2.8
2	B	30	THR	2.8
2	B	182	VAL	2.7
2	B	199	THR	2.7
1	A	197	ALA	2.7
1	A	193	ASP	2.6
1	A	5	LEU	2.6
1	A	185	SER	2.6
1	A	4	PHE	2.5
1	A	192	SER	2.5
1	A	30	THR	2.5
2	B	195	HIS	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	6	ALA	2.4
2	B	197	ALA	2.4
2	B	4	PHE	2.3
1	A	195	HIS	2.3
2	B	194	PRO	2.3
2	B	184	THR	2.3
2	B	187	GLY	2.2
2	B	196	LYS	2.2
1	A	72	GLY	2.1
1	A	199	THR	2.1
1	A	200	MET	2.1
1	A	31	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

MODRES-RSR INFOmissingINFO

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	PO4	B	245[A]	5/5	0.78	0.21	49,50,50,51	5
3	NAP	B	1245	48/48	0.92	0.10	21,29,42,43	0
3	NAP	A	245	48/48	0.94	0.10	17,36,56,58	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around NAP B 1245:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around NAP A 245:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.