



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 11, 2025 – 10:12 AM EDT

PDB ID : 9D6O / pdb\_00009d6o  
Title : Loop-Deleted DNA Polymerase Theta Polymerase Domain in Complex with a dsDNA Overhang and an Allosteric Inhibitor  
Authors : Fried, W.; Chen, X.S.  
Deposited on : 2024-08-15  
Resolution : 3.31 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0rc1  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.006 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.45.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*




The reported resolution of this entry is 3.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	1066 (3.34-3.30)
Clashscore	180529	1111 (3.34-3.30)
Ramachandran outliers	177936	1109 (3.34-3.30)
Sidechain outliers	177891	1108 (3.34-3.30)
RSRZ outliers	164620	1066 (3.34-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	652	
2	E	24	
3	F	14	

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5489 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase theta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	627	Total	C	N	O	S	0	0	0
			4956	3153	849	925	29			

There are 141 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1893	GLY	-	insertion	UNP O75417
A	?	-	LEU	deletion	UNP O75417
A	?	-	THR	deletion	UNP O75417
A	?	-	SER	deletion	UNP O75417
A	?	-	SER	deletion	UNP O75417
A	?	-	LYS	deletion	UNP O75417
A	?	-	THR	deletion	UNP O75417
A	?	-	ALA	deletion	UNP O75417
A	?	-	THR	deletion	UNP O75417
A	?	-	ILE	deletion	UNP O75417
A	?	-	GLY	deletion	UNP O75417
A	?	-	SER	deletion	UNP O75417
A	?	-	ARG	deletion	UNP O75417
A	?	-	PHE	deletion	UNP O75417
A	?	-	LYS	deletion	UNP O75417
A	?	-	GLN	deletion	UNP O75417
A	?	-	ALA	deletion	UNP O75417
A	?	-	SER	deletion	UNP O75417
A	?	-	SER	deletion	UNP O75417
A	?	-	PRO	deletion	UNP O75417
A	?	-	GLN	deletion	UNP O75417
A	?	-	GLU	deletion	UNP O75417
A	?	-	ILE	deletion	UNP O75417
A	?	-	PRO	deletion	UNP O75417
A	?	-	ILE	deletion	UNP O75417
A	?	-	ARG	deletion	UNP O75417
A	?	-	ASP	deletion	UNP O75417

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	ASP	deletion	UNP O75417
A	?	-	GLY	deletion	UNP O75417
A	?	-	PHE	deletion	UNP O75417
A	?	-	PRO	deletion	UNP O75417
A	?	-	ILE	deletion	UNP O75417
A	?	-	LYS	deletion	UNP O75417
A	?	-	GLY	deletion	UNP O75417
A	1895	GLY	CYS	conflict	UNP O75417
A	?	-	GLN	deletion	UNP O75417
A	?	-	LYS	deletion	UNP O75417
A	?	-	GLU	deletion	UNP O75417
A	?	-	GLN	deletion	UNP O75417
A	?	-	LYS	deletion	UNP O75417
A	?	-	HIS	deletion	UNP O75417
A	?	-	SER	deletion	UNP O75417
A	?	-	GLU	deletion	UNP O75417
A	?	-	ILE	deletion	UNP O75417
A	?	-	SER	deletion	UNP O75417
A	?	-	ALA	deletion	UNP O75417
A	?	-	SER	deletion	UNP O75417
A	1918	GLY	LEU	conflict	UNP O75417
A	1931	GLY	VAL	conflict	UNP O75417
A	1932	SER	PRO	conflict	UNP O75417
A	1933	GLY	PRO	conflict	UNP O75417
A	1934	GLY	SER	conflict	UNP O75417
A	?	-	ASN	deletion	UNP O75417
A	?	-	ARG	deletion	UNP O75417
A	?	-	GLU	deletion	UNP O75417
A	?	-	MET	deletion	UNP O75417
A	?	-	LYS	deletion	UNP O75417
A	?	-	ASN	deletion	UNP O75417
A	?	-	GLN	deletion	UNP O75417
A	?	-	GLY	deletion	UNP O75417
A	?	-	SER	deletion	UNP O75417
A	?	-	LYS	deletion	UNP O75417
A	?	-	LYS	deletion	UNP O75417
A	?	-	THR	deletion	UNP O75417
A	?	-	LEU	deletion	UNP O75417
A	?	-	GLY	deletion	UNP O75417
A	?	-	SER	deletion	UNP O75417
A	?	-	THR	deletion	UNP O75417
A	?	-	ARG	deletion	UNP O75417

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	ARG	deletion	UNP O75417
A	?	-	GLY	deletion	UNP O75417
A	?	-	ILE	deletion	UNP O75417
A	?	-	ASP	deletion	UNP O75417
A	?	-	ASN	deletion	UNP O75417
A	?	-	GLY	deletion	UNP O75417
A	?	-	ARG	deletion	UNP O75417
A	?	-	LYS	deletion	UNP O75417
A	2171	GLY	LEU	conflict	UNP O75417
A	2172	GLY	ARG	conflict	UNP O75417
A	2173	SER	LEU	conflict	UNP O75417
A	2175	GLY	ARG	conflict	UNP O75417
A	?	-	PRO	deletion	UNP O75417
A	?	-	THR	deletion	UNP O75417
A	?	-	LEU	deletion	UNP O75417
A	?	-	VAL	deletion	UNP O75417
A	?	-	GLY	deletion	UNP O75417
A	?	-	GLU	deletion	UNP O75417
A	?	-	SER	deletion	UNP O75417
A	?	-	PRO	deletion	UNP O75417
A	?	-	PRO	deletion	UNP O75417
A	?	-	SER	deletion	UNP O75417
A	?	-	GLN	deletion	UNP O75417
A	?	-	ALA	deletion	UNP O75417
A	?	-	VAL	deletion	UNP O75417
A	?	-	GLY	deletion	UNP O75417
A	?	-	LYS	deletion	UNP O75417
A	?	-	GLY	deletion	UNP O75417
A	?	-	LEU	deletion	UNP O75417
A	?	-	LEU	deletion	UNP O75417
A	?	-	PRO	deletion	UNP O75417
A	?	-	MET	deletion	UNP O75417
A	?	-	GLY	deletion	UNP O75417
A	?	-	ARG	deletion	UNP O75417
A	?	-	GLY	deletion	UNP O75417
A	?	-	LYS	deletion	UNP O75417
A	?	-	TYR	deletion	UNP O75417
A	?	-	LYS	deletion	UNP O75417
A	?	-	LYS	deletion	UNP O75417
A	?	-	GLY	deletion	UNP O75417
A	?	-	PHE	deletion	UNP O75417
A	?	-	SER	deletion	UNP O75417

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	?	-	VAL	deletion	UNP O75417
A	?	-	ASN	deletion	UNP O75417
A	?	-	PRO	deletion	UNP O75417
A	?	-	ARG	deletion	UNP O75417
A	?	-	CYS	deletion	UNP O75417
A	?	-	GLN	deletion	UNP O75417
A	?	-	ALA	deletion	UNP O75417
A	?	-	GLN	deletion	UNP O75417
A	?	-	MET	deletion	UNP O75417
A	?	-	GLU	deletion	UNP O75417
A	?	-	GLU	deletion	UNP O75417
A	?	-	ARG	deletion	UNP O75417
A	2261	GLY	ALA	conflict	UNP O75417
A	2262	GLY	ALA	conflict	UNP O75417
A	2263	SER	ASP	conflict	UNP O75417
A	2264	GLY	ARG	conflict	UNP O75417
A	?	-	GLN	deletion	UNP O75417
A	?	-	THR	deletion	UNP O75417
A	?	-	GLY	deletion	UNP O75417
A	?	-	LEU	deletion	UNP O75417
A	?	-	SER	deletion	UNP O75417
A	?	-	ARG	deletion	UNP O75417
A	?	-	LYS	deletion	UNP O75417
A	?	-	ARG	deletion	UNP O75417
A	?	-	LYS	deletion	UNP O75417
A	?	-	LEU	deletion	UNP O75417
A	2522	GLY	GLN	conflict	UNP O75417
A	2524	SER	-	insertion	UNP O75417
A	2525	GLY	MET	conflict	UNP O75417
A	2526	GLY	PHE	conflict	UNP O75417

- Molecule 2 is a DNA chain called DNA Template.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	E	13	Total	C	N	O	P	0	0	0
			263	124	47	79	13			


- Molecule 3 is a DNA chain called DNA Primer.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	10	Total	C	N	O	P	0	0	0
			204	96	39	59	10			

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is 2'-3'-DIDEOXYGUANOSINE-5'-TRIPHOSPHATE (CCD ID: DG3) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>12</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			30	10	5	12	3		

- Molecule 6 is 2-[3-(2-hydroxyethyl)-2-oxoimidazolidin-1-yl]-4,6-bis(trifluoromethyl)phenyl (4-fluorophenyl)(methyl)carbamate (CCD ID: A1A2D) (formula: C<sub>21</sub>H<sub>18</sub>F<sub>7</sub>N<sub>3</sub>O<sub>4</sub>) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	F	N	O	0	0
			35	21	7	3	4		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase theta



- Molecule 2: DNA Template



- Molecule 3: DNA Primer



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	172.36Å 172.36Å 62.41Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	149.26 – 3.31 149.26 – 3.31	Depositor EDS
% Data completeness (in resolution range)	99.3 (149.26-3.31) 99.3 (149.26-3.31)	Depositor EDS
$R_{merge}$	0.22	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.91 (at 3.07Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.209 , 0.234 0.210 , 0.242	Depositor DCC
$R_{free}$ test set	908 reflections (4.51%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	108.4	Xtriage
Anisotropy	0.277	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 116.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.43$ , $\langle L^2 \rangle = 0.26$	Xtriage
Estimated twinning fraction	0.074 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	5489	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	142.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DG3, MG, A1A2D

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.08	0/5050	0.26	0/6809
2	E	0.16	0/293	0.30	0/449
3	F	0.12	0/228	0.27	0/349
All	All	0.09	0/5571	0.26	0/7607

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4956	0	4978	21	0
2	E	263	0	146	3	0
3	F	204	0	110	3	0
4	A	1	0	0	0	0
5	A	30	0	12	0	0
6	A	35	0	0	0	0
All	All	5489	0	5246	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 25 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2211:VAL:HG22	1:A:2314:MET:HE1	1.61	0.81
1:A:2091:GLY:O	1:A:2228:ARG:NH2	2.31	0.58
1:A:2338:ILE:HD11	1:A:2480:ILE:HD12	1.88	0.56
1:A:2483:ILE:HG23	1:A:2565:ALA:HB1	1.89	0.54
1:A:2315:ARG:NH2	3:F:13:DG:O3'	2.43	0.51

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	617/652 (95%)	579 (94%)	38 (6%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	547/561 (98%)	547 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	2221	ASN
1	A	2333	GLN
1	A	2463	HIS
1	A	2040	HIS
1	A	2035	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	A1A2D	A	2603	-	37,37,37	1.23	1 (2%)	53,56,56	1.06	3 (5%)
5	DG3	A	2602	4	28,32,32	0.79	0	33,50,50	1.21	4 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	A1A2D	A	2603	-	-	5/31/44/44	0/3/3/3
5	DG3	A	2602	4	-	7/18/31/31	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	2603	A1A2D	C08-N01	5.05	1.46	1.36

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	2603	A1A2D	O01-C08-N01	-2.63	120.56	124.06
5	A	2602	DG3	O2A-PA-O1A	2.21	122.71	112.44
5	A	2602	DG3	O2B-PB-O3B	2.15	113.09	107.27
5	A	2602	DG3	O2A-PA-O3A	2.13	113.02	107.27
6	A	2603	A1A2D	C07-N01-C04	2.09	120.50	117.01

There are no chirality outliers.

5 of 12 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	A	2603	A1A2D	C21-C20-N03-C19
6	A	2603	A1A2D	C21-C20-N03-C18
6	A	2603	A1A2D	N01-C08-O02-C09
5	A	2602	DG3	O4'-C4'-C5'-O5'
5	A	2602	DG3	PB-O3A-PA-O1A

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


## Ligand A1A2D A 2603




Bond lengths




Bond angles



Torsions



Rings



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	627/652 (96%)	-0.32	0 100 100	90, 138, 180, 235	0
2	E	13/24 (54%)	-0.73	0 100 100	118, 135, 241, 249	0
3	F	10/14 (71%)	-0.73	0 100 100	109, 148, 216, 217	0
All	All	650/690 (94%)	-0.33	0 100 100	90, 138, 183, 249	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	DG3	A	2602	30/30	0.89	0.09	110,130,154,161	0
6	A1A2D	A	2603	35/35	0.93	0.10	137,170,194,200	0
4	MG	A	2601	1/1	0.99	0.04	97,97,97,97	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around DG3 A 2602:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around A1A2D A 2603:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.