



## wwPDB EM Validation Summary Report ⓘ

Oct 26, 2024 – 05:03 PM EDT

PDB ID : 6DBL  
EMDB ID : EMD-7845  
Title : Cryo-EM structure of RAG in complex with 12-RSS and 23-RSS substrate DNAs  
Authors : Wu, H.; Liao, M.; Ru, H.; Mi, W.  
Deposited on : 2018-05-03  
Resolution : 5.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113  
MolProbity : 4.02b-467  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*









The reported resolution of this entry is 5.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1159	
1	C	1159	
2	B	533	
2	D	533	
3	E	50	
4	F	50	
5	G	61	
6	H	61	

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 19929 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Recombination activating gene 1 - MBP chimera.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	616	Total	C	N	O	S	0	0
			4972	3113	893	929	37		
1	C	616	Total	C	N	O	S	0	0
			4972	3113	893	929	37		

There are 68 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-127	MET	-	initiating methionine	UNP P0AEX9
A	-126	GLY	-	expression tag	UNP P0AEX9
A	-125	SER	-	expression tag	UNP P0AEX9
A	-124	SER	-	expression tag	UNP P0AEX9
A	-123	HIS	-	expression tag	UNP P0AEX9
A	-122	HIS	-	expression tag	UNP P0AEX9
A	-121	HIS	-	expression tag	UNP P0AEX9
A	-120	HIS	-	expression tag	UNP P0AEX9
A	-119	HIS	-	expression tag	UNP P0AEX9
A	-118	HIS	-	expression tag	UNP P0AEX9
A	-117	GLY	-	expression tag	UNP P0AEX9
A	-116	THR	-	expression tag	UNP P0AEX9
A	-115	LYS	-	expression tag	UNP P0AEX9
A	-114	THR	-	expression tag	UNP P0AEX9
A	251	GLY	-	linker	UNP P0AEX9
A	252	THR	-	linker	UNP P0AEX9
A	253	ASP	-	linker	UNP P0AEX9
A	254	TYR	-	linker	UNP P0AEX9
A	255	ASP	-	linker	UNP P0AEX9
A	256	ILE	-	linker	UNP P0AEX9
A	257	PRO	-	linker	UNP P0AEX9
A	258	THR	-	linker	UNP P0AEX9
A	259	THR	-	linker	UNP P0AEX9
A	260	LEU	-	linker	UNP P0AEX9
A	261	GLU	-	linker	UNP P0AEX9
A	262	VAL	-	linker	UNP P0AEX9

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	263	LEU	-	linker	UNP P0AEX9
A	264	PHE	-	linker	UNP P0AEX9
A	265	GLN	-	linker	UNP P0AEX9
A	266	GLY	-	linker	UNP P0AEX9
A	267	PRO	-	linker	UNP P0AEX9
A	268	LEU	-	linker	UNP P0AEX9
A	269	GLY	-	linker	UNP P0AEX9
A	270	SER	-	linker	UNP P0AEX9
C	-127	MET	-	initiating methionine	UNP P0AEX9
C	-126	GLY	-	expression tag	UNP P0AEX9
C	-125	SER	-	expression tag	UNP P0AEX9
C	-124	SER	-	expression tag	UNP P0AEX9
C	-123	HIS	-	expression tag	UNP P0AEX9
C	-122	HIS	-	expression tag	UNP P0AEX9
C	-121	HIS	-	expression tag	UNP P0AEX9
C	-120	HIS	-	expression tag	UNP P0AEX9
C	-119	HIS	-	expression tag	UNP P0AEX9
C	-118	HIS	-	expression tag	UNP P0AEX9
C	-117	GLY	-	expression tag	UNP P0AEX9
C	-116	THR	-	expression tag	UNP P0AEX9
C	-115	LYS	-	expression tag	UNP P0AEX9
C	-114	THR	-	expression tag	UNP P0AEX9
C	251	GLY	-	linker	UNP P0AEX9
C	252	THR	-	linker	UNP P0AEX9
C	253	ASP	-	linker	UNP P0AEX9
C	254	TYR	-	linker	UNP P0AEX9
C	255	ASP	-	linker	UNP P0AEX9
C	256	ILE	-	linker	UNP P0AEX9
C	257	PRO	-	linker	UNP P0AEX9
C	258	THR	-	linker	UNP P0AEX9
C	259	THR	-	linker	UNP P0AEX9
C	260	LEU	-	linker	UNP P0AEX9
C	261	GLU	-	linker	UNP P0AEX9
C	262	VAL	-	linker	UNP P0AEX9
C	263	LEU	-	linker	UNP P0AEX9
C	264	PHE	-	linker	UNP P0AEX9
C	265	GLN	-	linker	UNP P0AEX9
C	266	GLY	-	linker	UNP P0AEX9
C	267	PRO	-	linker	UNP P0AEX9
C	268	LEU	-	linker	UNP P0AEX9
C	269	GLY	-	linker	UNP P0AEX9
C	270	SER	-	linker	UNP P0AEX9

- Molecule 2 is a protein called Recombination activating gene 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	351	Total	C	N	O	S	0	0
			2714	1716	470	509	19		
2	D	351	Total	C	N	O	S	0	0
			2714	1716	470	509	19		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	GLY	-	expression tag	UNP Q1RLW7
B	-1	GLY	-	expression tag	UNP Q1RLW7
B	0	SER	-	expression tag	UNP Q1RLW7
D	-2	GLY	-	expression tag	UNP Q1RLW7
D	-1	GLY	-	expression tag	UNP Q1RLW7
D	0	SER	-	expression tag	UNP Q1RLW7

- Molecule 3 is a DNA chain called Molecule name: Forward strand of 12-RSS substrate DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	E	50	Total	C	N	O	P	0	0
			1023	486	192	295	50		

- Molecule 4 is a DNA chain called Molecule name: Reverse strand of 12-RSS substrate DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	F	50	Total	C	N	O	P	0	0
			1027	489	183	305	50		

- Molecule 5 is a DNA chain called Molecule name: Forward strand of 23-RSS substrate DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	G	61	Total	C	N	O	P	0	0
			1245	593	223	368	61		

- Molecule 6 is a DNA chain called Molecule name: Reverse strand of 23-RSS substrate DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	H	61	Total	C	N	O	P	0	0
			1256	596	235	364	61		

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
7	A	1	Total 1	Zn 1	0
7	C	1	Total 1	Zn 1	0

- Molecule 8 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		AltConf
8	A	1	Total 1	Ca 1	0
8	C	1	Total 1	Ca 1	0
8	E	1	Total 1	Ca 1	0
8	G	1	Total 1	Ca 1	0






THR	ALA	ASN	GLY	PRO	PHE	ALA	GLN	MET
LEU	VAL	ALA	LEU	TRP	LYS	VAL	VAL	GLY
SER	THR	ALA	GLU	ALA	TYR	GLU	ALA	SER
LEU	CYS	SER	VAL	TRP	GLU	ALA	THR	HIS
PRO	VAL	ARG	ASN	ASN	GLY	SER	GLY	HIS
LEU	CYS	GLN	LYS	ASN	LYS	LEU	ASP	HIS
LEU	ASP	THR	ASP	ASP	TYR	ILE	GLY	HIS
PRO	HIS	VAL	PRO	THR	ASP	ASN	PRO	HIS
PRO	LEU	ASP	PRO	SER	ILE	ASN	ASP	HIS
SER	LEU	GLU	LEU	VAL	LYS	LYS	ILE	GLY
GLU	SER	ALA	GLY	VAL	ASP	ASP	ILE	THR
GLU	ASP	LEU	ALA	ASN	VAL	LEU	PHE	LYS
CYS	PRO	LYS	VAL	GLY	GLY	LEU	THR	THR
SER	VAL	ASP	VAL	TYR	VAL	PRO	ALA	GLU
ASP	GLN	ALA	LEU	VAL	ASP	ASN	GLU	GLU
TRP	SER	GLN	LYS	THR	ASN	PRO	ASP	GLY
VAL	PRO	THR	SER	VAL	ALA	PRO	ARG	LYS
ARG	CYS	GLY	TYR	LEU	GLY	LYS	ARG	LEU
LEU	ARG	THR	GLU	PRO	ALA	THR	PHE	VAL
ASP	HIS	ASP	GLU	THR	LYS	TRP	GLY	ILE
SER	LEU	TYR	GLU	PHE	ALA	GLU	TYR	THR
PHE	PHE	ASP	LEU	LYS	GLY	GLU	ALA	ILE
ARG	CYS	ILE	LYS	GLN	LEU	ILE	GLN	ASN
GLU	ARG	PRO	LYS	GLN	THR	PRO	SER	GLY
HIS	LEU	THR	ASP	PRO	PHE	LEU	LEU	ASN
CYS	CYS	THR	PRO	THR	LEU	ALA	LYS	GLY
LEU	ILE	LEU	ARG	LYS	VAL	LYS	ASP	GLU
ASN	ILE	GLU	ILE	PRO	ASP	GLY	GLY	VAL
HIS	ARG	VAL	ALA	PHE	LEU	GLU	ALA	TYR
THR	THR	LEU	THR	VAL	ILE	ILE	ILE	ASN
ARG	HIS	GLN	MET	VAL	LYS	LYS	THR	GLY
GLU	GLU	GLY	GLU	LEU	ASN	ALA	PRO	ALA
LYS	ALA	GLY	GLU	LEU	LYS	LYS	GLU	GLU
GLU	LEU	PRO	ASN	SER	HIS	GLY	LYS	VAL
SER	GLY	LEU	ALA	ALA	MET	LYS	ALA	GLY
GLN	PRO	GLY	GLN	GLY	ASN	SER	PHE	LYS
GLU	ASN	SER	GLY	ILE	ALA	GLN	GLN	GLY
CYS	CYS	ARG	GLY	ASN	ASP	LEU	ASP	PHE
GLU	PRO	GLN	GLU	ALA	THR	MET	GLY	LYS
THR	THR	GLN	ILE	ALA	ASP	PHE	LEU	ASN
PRO	CYS	ARG	MET	SER	TYR	ASN	TYR	ASP
ASN	ASN	ASP	PRO	PRO	THR	LEU	PRO	THR
GLU	GLN	HIS	ASN	ASN	ILE	GLN	PHE	GLY
GLN	HIS	LEU	ILE	LYS	ALA	GLU	THR	ILE
ASN	LEU	SER	PRO	GLU	GLU	PRO	TRP	LYS
LEU	ASN	THR	GLN	LEU	ALA	TYR	ASP	THR
ASP	PRO	LYS	MET	ALA	ALA	PHE	ALA	VAL
GLY	SER	LEU	SER	LYS	PHE	THR	VAL	THR
TYR	HIS	ILE	PHE	GLU	ASN	TRP	VAL	VAL
LEU	LEU	PRO	ALA	PHE	LYS	GLU	ARG	HIS
PRO	ILE	THR	TRP	LEU	GLY	ILE	TYR	PRO
VAL	LYS	GLU	TYR	GLU	THR	ALA	GLY	ASN
ASN	PRO	VAL	ALA	ASN	THR	ALA	LYS	LYS
LYS	ALA	VAL	VAL	TYR	THR	ASP	LEU	LYS
GLN	ALA	ARG	ARG	THR	ALA	ILE	LEU	GLU
CYS	LYS	ALA	ARG	LEU	ALA	ASP	ILE	LEU
G408	GLY	ASN	THR	ASN	PHE	THR	PRO	PRO
G409	LYS	ASN	THR	THR	THR	THR	THR	GLY
R410	PHE	THR	THR	THR	THR	THR	THR	THR
R411	ALA	ASN	THR	GLU	GLY	THR	PRO	THR








## • Molecule 2: Recombination activating gene 2

Chain B: 27% 35% 34%







• Molecule 2: Recombination activating gene 2




• Molecule 3: Molecule name: Forward strand of 12-RSS substrate DNA




• Molecule 4: Molecule name: Reverse strand of 12-RSS substrate DNA



• Molecule 5: Molecule name: Forward strand of 23-RSS substrate DNA



• Molecule 6: Molecule name: Reverse strand of 23-RSS substrate DNA



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	23176	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	47	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.122	Depositor
Minimum map value	-0.050	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.007	Depositor
Recommended contour level	0.03	Depositor
Map size (Å)	237.69601, 237.69601, 237.69601	wwPDB
Map dimensions	192, 192, 192	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.238, 1.238, 1.238	Depositor

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.83	9/5067 (0.2%)	1.22	26/6812 (0.4%)
1	C	0.81	5/5067 (0.1%)	1.21	43/6812 (0.6%)
2	B	0.76	2/2784 (0.1%)	1.29	25/3784 (0.7%)
2	D	0.76	2/2784 (0.1%)	1.17	15/3784 (0.4%)
3	E	1.34	9/1148 (0.8%)	1.18	8/1768 (0.5%)
4	F	1.21	5/1150 (0.4%)	1.27	10/1774 (0.6%)
5	G	1.27	11/1394 (0.8%)	1.10	3/2148 (0.1%)
6	H	1.10	6/1410 (0.4%)	1.13	4/2175 (0.2%)
All	All	0.92	49/20804 (0.2%)	1.21	134/29057 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	C	0	3
2	B	0	3
2	D	0	2
All	All	0	12

The worst 5 of 49 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	F	40	DC	C1'-N1	-8.10	1.35	1.47
5	G	18	DA	C3'-O3'	8.07	1.54	1.44
6	H	33	DT	C1'-N1	7.67	1.59	1.49
1	A	851	TRP	CB-CG	7.56	1.63	1.50
3	E	14	DT	C1'-N1	6.90	1.58	1.49

The worst 5 of 134 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	666	GLN	CG-CD-OE1	24.10	169.80	121.60
2	B	100	ASN	CB-CG-ND2	-16.20	77.82	116.70
1	A	666	GLN	CG-CD-NE2	-16.18	77.87	116.70
1	C	544	ASN	N-CA-CB	-15.62	82.49	110.60
2	B	298	HIS	CB-CG-CD2	-13.79	88.04	130.80

There are no chirality outliers.

5 of 12 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	478	GLY	Peptide
1	A	512	GLY	Peptide
1	A	549	TRP	Mainchain
1	A	999	ARG	Peptide
2	B	61	SER	Mainchain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4972	0	4935	388	0
1	C	4972	0	4937	425	0
2	B	2714	0	2665	226	0
2	D	2714	0	2665	218	0
3	E	1023	0	561	74	0
4	F	1027	0	566	70	0
5	G	1245	0	688	78	0
6	H	1256	0	686	62	0
7	A	1	0	0	0	0
7	C	1	0	0	0	0
8	A	1	0	0	0	0
8	C	1	0	0	0	0
8	E	1	0	0	0	0
8	G	1	0	0	0	0
All	All	19929	0	17703	1392	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 37.

The worst 5 of 1392 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:543:LYS:C	1:C:544:ASN:HD22	1.33	1.31
1:C:543:LYS:O	1:C:544:ASN:ND2	1.78	1.17
1:C:903:LYS:HZ1	1:C:937:GLN:NE2	1.45	1.15
1:C:731:GLU:OE2	1:C:960:LYS:NZ	1.80	1.13
1:C:820:ILE:HG21	1:C:870:ARG:HH12	1.13	1.11

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	612/1159 (53%)	576 (94%)	31 (5%)	5 (1%)	16	54
1	C	612/1159 (53%)	576 (94%)	27 (4%)	9 (2%)	8	39
2	B	349/533 (66%)	331 (95%)	18 (5%)	0	100	100
2	D	349/533 (66%)	314 (90%)	33 (10%)	2 (1%)	22	60
All	All	1922/3384 (57%)	1797 (94%)	109 (6%)	16 (1%)	19	54

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	480	GLY
1	C	973	GLY
1	C	976	GLY
1	A	464	HIS
1	A	994	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	545/1000 (54%)	534 (98%)	11 (2%)	50	68
1	C	545/1000 (54%)	524 (96%)	21 (4%)	27	48
2	B	303/465 (65%)	297 (98%)	6 (2%)	50	68
2	D	303/465 (65%)	298 (98%)	5 (2%)	56	72
All	All	1696/2930 (58%)	1653 (98%)	43 (2%)	43	62

5 of 43 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	870	ARG
1	C	999	ARG
1	C	878	ARG
1	C	944	SER
1	C	1029	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	544	ASN
1	C	936	GLN
2	D	253	HIS
1	C	937	GLN
1	C	475	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.



## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-7845. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map



X Index: 96



Y Index: 96




Z Index: 96


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 82



Y Index: 98




Z Index: 80


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.03. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 185 nm<sup>3</sup>; this corresponds to an approximate mass of 167 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)



\*Reported resolution corresponds to spatial frequency of 0.200 Å<sup>-1</sup>

## 8 Fourier-Shell correlation


This section was not generated. No FSC curve or half-maps provided.




## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-7845 and PDB model 6DBL. Per-residue inclusion information can be found in section 3 on page 7.


### 9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.03 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.03).

## 9.4 Atom inclusion [i](#)



At the recommended contour level, 87% of all backbone atoms, 69% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.03) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div></div> 0.6860	<div></div> 0.2300
A	<div></div> 0.6840	<div></div> 0.2300
B	<div></div> 0.7710	<div></div> 0.2610
C	<div></div> 0.7010	<div></div> 0.2360
D	<div></div> 0.7630	<div></div> 0.2530
E	<div></div> 0.6620	<div></div> 0.2140
F	<div></div> 0.6490	<div></div> 0.2200
G	<div></div> 0.5180	<div></div> 0.1740
H	<div></div> 0.5100	<div></div> 0.1670

1.0

0.0

<0.0