



## wwPDB EM Validation Summary Report ⓘ

May 19, 2025 – 06:36 PM EDT

PDB ID : 8DBW / pdb\_00008dbw  
EMDB ID : EMD-27311  
Title : E. coli ATP synthase imaged in 10mM MgATP State3 "down" Fo classified  
Authors : Sobti, M.; Stewart, A.G.  
Deposited on : 2022-06-14  
Resolution : 4.10 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev118  
Mogul : 2022.3.0, CSD as543be (2022)  
MolProbity : 4-5-2 with Phenix2.0rc1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.43.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 4.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	512	97%
1	B	512	94% 5%
1	C	512	93% 6%
2	D	460	96% .
2	E	460	95% 5%
2	F	460	95% .
3	G	284	89% 9% .
4	H	134	79% 16% . .

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
5	I	77	
5	J	77	
5	L	77	
5	M	77	
5	N	77	
5	O	77	
5	P	77	
5	Q	77	
5	R	77	
5	S	77	
6	W	173	
7	X	155	
7	Y	155	
8	a	266	

## 2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 36835 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP synthase subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	510	Total	C	N	O	S	0	0
			3852	2420	679	744	9		
1	B	505	Total	C	N	O	S	0	0
			3810	2395	672	734	9		
1	C	510	Total	C	N	O	S	0	0
			3845	2416	678	742	9		

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	ALA	CYS	conflict	UNP A0A7U9G3U3
A	90	ALA	CYS	conflict	UNP A0A7U9G3U3
A	193	ALA	CYS	conflict	UNP A0A7U9G3U3
A	243	ALA	CYS	conflict	UNP A0A7U9G3U3
A	419	ASN	LYS	conflict	UNP A0A7U9G3U3
B	47	ALA	CYS	conflict	UNP A0A7U9G3U3
B	90	ALA	CYS	conflict	UNP A0A7U9G3U3
B	193	ALA	CYS	conflict	UNP A0A7U9G3U3
B	243	ALA	CYS	conflict	UNP A0A7U9G3U3
B	419	ASN	LYS	conflict	UNP A0A7U9G3U3
C	47	ALA	CYS	conflict	UNP A0A7U9G3U3
C	90	ALA	CYS	conflict	UNP A0A7U9G3U3
C	193	ALA	CYS	conflict	UNP A0A7U9G3U3
C	243	ALA	CYS	conflict	UNP A0A7U9G3U3
C	419	ASN	LYS	conflict	UNP A0A7U9G3U3

- Molecule 2 is a protein called ATP synthase subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	D	460	Total	C	N	O	S	0	0
			3534	2227	604	688	15		
2	E	460	Total	C	N	O	S	0	0
			3534	2227	604	688	15		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					AltConf	Trace
2	F	458	Total	C	N	O	S	0	0
			3521	2219	602	686	14		

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	137	ALA	CYS	conflict	UNP A0A192CEZ8
E	137	ALA	CYS	conflict	UNP A0A192CEZ8
F	137	ALA	CYS	conflict	UNP A0A192CEZ8

- Molecule 3 is a protein called ATP synthase gamma chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	G	284	Total	C	N	O	S	0	0
			2187	1373	382	420	12		

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	5	ASP	GLU	conflict	UNP C3SLA2
G	87	ALA	CYS	conflict	UNP C3SLA2
G	112	ALA	CYS	conflict	UNP C3SLA2

- Molecule 4 is a protein called ATP synthase epsilon chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	H	134	Total	C	N	O	S	0	0
			1022	641	178	200	3		

- Molecule 5 is a protein called ATP synthase subunit c.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	I	77	Total	C	N	O	S	0	0
			558	370	88	93	7		
5	J	77	Total	C	N	O	S	0	0
			558	370	88	93	7		
5	L	77	Total	C	N	O	S	0	0
			558	370	88	93	7		
5	M	77	Total	C	N	O	S	0	0
			558	370	88	93	7		
5	N	77	Total	C	N	O	S	0	0
			558	370	88	93	7		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					AltConf	Trace
5	O	77	Total	C	N	O	S	0	0
			558	370	88	93	7		
5	P	77	Total	C	N	O	S	0	0
			558	370	88	93	7		
5	Q	77	Total	C	N	O	S	0	0
			558	370	88	93	7		
5	R	76	Total	C	N	O	S	0	0
			552	367	87	91	7		
5	S	77	Total	C	N	O	S	0	0
			558	370	88	93	7		

- Molecule 6 is a protein called ATP synthase subunit delta.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	W	173	Total	C	N	O	S	0	0
			1318	826	233	253	6		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
W	64	ALA	CYS	conflict	UNP V0ZA15
W	140	ALA	CYS	conflict	UNP V0ZA15

- Molecule 7 is a protein called ATP synthase subunit b.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	X	153	Total	C	N	O	S	0	0
			1179	737	213	226	3		
7	Y	155	Total	C	N	O	S	0	0
			1172	733	212	224	3		

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	21	ALA	CYS	conflict	UNP D6IFY0
Y	21	ALA	CYS	conflict	UNP D6IFY0


- Molecule 8 is a protein called ATP synthase subunit a.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	a	266	Total	C	N	O	S	0	0
			2104	1418	326	350	10		

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
a	27	ALA	-	insertion	UNP C3SL77
a	?	-	THR	deletion	UNP C3SL77
a	38	TYR	PHE	conflict	UNP C3SL77
a	46	PHE	MET	conflict	UNP C3SL77
a	47	MET	PHE	conflict	UNP C3SL77


- Molecule 9 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula:  $C_{10}H_{16}N_5O_{13}P_3$ ).



Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
10	C	1	Total	Mg	0
			1	1	
10	D	1	Total	Mg	0
			1	1	
10	F	1	Total	Mg	0
			1	1	

- Molecule 11 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).



Mol	Chain	Residues	Atoms					AltConf
11	D	1	Total	C	N	O	P	0
			27	10	5	10	2	
11	E	1	Total	C	N	O	P	0
			27	10	5	10	2	




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ATP synthase subunit alpha

Chain A:  97%





- Molecule 1: ATP synthase subunit alpha

Chain B:  94% 5%




- Molecule 1: ATP synthase subunit alpha


Chain C:  93% 6%





- Molecule 2: ATP synthase subunit beta

Chain D:  96% 7%







- Molecule 2: ATP synthase subunit beta






- Molecule 2: ATP synthase subunit beta




- Molecule 3: ATP synthase gamma chain




- Molecule 4: ATP synthase epsilon chain





- Molecule 5: ATP synthase subunit c








• Molecule 5: ATP synthase subunit c





• Molecule 5: ATP synthase subunit c




• Molecule 5: ATP synthase subunit c




• Molecule 5: ATP synthase subunit c




- Molecule 5: ATP synthase subunit c




- Molecule 5: ATP synthase subunit c




- Molecule 5: ATP synthase subunit c




- Molecule 5: ATP synthase subunit c




- Molecule 5: ATP synthase subunit c










- Molecule 6: ATP synthase subunit delta






- Molecule 7: ATP synthase subunit b




- Molecule 7: ATP synthase subunit b



- Molecule 8: ATP synthase subunit a





## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	8757	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	48	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.053	Depositor
Minimum map value	-0.017	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.015	Depositor
Map size (Å)	379.80798, 379.80798, 379.80798	wwPDB
Map dimensions	352, 352, 352	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.079, 1.079, 1.079	Depositor

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, ADP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.17	0/3908	0.39	0/5288
1	B	0.16	0/3864	0.42	0/5228
1	C	0.20	0/3899	0.45	0/5276
2	D	0.17	0/3591	0.40	0/4858
2	E	0.16	0/3591	0.37	0/4858
2	F	0.17	0/3578	0.39	0/4841
3	G	0.28	0/2218	0.52	0/2990
4	H	0.44	0/1037	0.83	5/1398 (0.4%)
5	I	0.22	0/566	0.48	0/764
5	J	0.21	0/566	0.47	0/764
5	L	0.19	0/566	0.48	0/764
5	M	0.33	0/566	0.57	0/764
5	N	0.26	0/566	0.49	0/764
5	O	0.24	0/566	0.48	0/764
5	P	0.22	0/566	0.48	0/764
5	Q	0.24	0/566	0.60	2/764 (0.3%)
5	R	0.32	0/560	0.66	0/757
5	S	0.30	0/566	0.57	0/764
6	W	0.23	0/1331	0.45	0/1794
7	X	0.28	0/1187	0.51	0/1596
7	Y	0.23	0/1180	0.43	0/1589
8	a	0.27	0/2164	0.54	0/2954
All	All	0.22	0/37202	0.46	7/50303 (0.0%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	39	ALA	N-CA-C	7.19	116.59	108.25
4	H	73	PRO	N-CA-C	5.82	120.14	111.41

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	42	LEU	N-CA-C	5.53	117.67	108.99
5	Q	45	LEU	CA-C-N	5.44	123.66	120.24
5	Q	45	LEU	C-N-CA	5.44	123.66	120.24

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3852	0	3896	6	0
1	B	3810	0	3863	13	0
1	C	3845	0	3896	17	0
2	D	3534	0	3548	10	0
2	E	3534	0	3548	11	0
2	F	3521	0	3530	9	0
3	G	2187	0	2238	49	0
4	H	1022	0	1027	52	0
5	I	558	0	598	7	0
5	J	558	0	598	6	0
5	L	558	0	598	2	0
5	M	558	0	598	4	0
5	N	558	0	598	7	0
5	O	558	0	598	6	0
5	P	558	0	598	6	0
5	Q	558	0	598	6	0
5	R	552	0	593	13	0
5	S	558	0	598	8	0
6	W	1318	0	1343	3	0
7	X	1179	0	1223	9	0
7	Y	1172	0	1198	3	0
8	a	2104	0	2194	17	0
9	A	31	0	12	0	0
9	B	31	0	12	0	0
9	C	31	0	12	0	0
9	F	31	0	12	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
10	A	1	0	0	0	0
10	B	1	0	0	0	0
10	C	1	0	0	0	0
10	D	1	0	0	0	0
10	F	1	0	0	0	0
11	D	27	0	12	0	0
11	E	27	0	12	0	0
All	All	36835	0	37551	213	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 213 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:38:HIS:HD2	5:R:42:GLN:HA	1.38	0.87
3:G:204:ASP:HB2	4:H:41:LEU:HB2	1.56	0.86
4:H:42:LEU:HD12	4:H:42:LEU:H	1.50	0.76
3:G:203:TRP:O	4:H:71:VAL:HG11	1.87	0.75
5:Q:63:ILE:HG23	5:Q:64:PRO:HD3	1.69	0.75

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	508/512 (99%)	499 (98%)	9 (2%)	0	100	100
1	B	503/512 (98%)	488 (97%)	15 (3%)	0	100	100
1	C	508/512 (99%)	496 (98%)	12 (2%)	0	100	100
2	D	458/460 (100%)	450 (98%)	8 (2%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	E	458/460 (100%)	446 (97%)	12 (3%)	0	100	100
2	F	456/460 (99%)	448 (98%)	7 (2%)	1 (0%)	44	77
3	G	282/284 (99%)	276 (98%)	5 (2%)	1 (0%)	30	67
4	H	132/134 (98%)	125 (95%)	7 (5%)	0	100	100
5	I	75/77 (97%)	74 (99%)	1 (1%)	0	100	100
5	J	75/77 (97%)	73 (97%)	2 (3%)	0	100	100
5	L	75/77 (97%)	74 (99%)	1 (1%)	0	100	100
5	M	75/77 (97%)	74 (99%)	1 (1%)	0	100	100
5	N	75/77 (97%)	73 (97%)	2 (3%)	0	100	100
5	O	75/77 (97%)	74 (99%)	1 (1%)	0	100	100
5	P	75/77 (97%)	74 (99%)	1 (1%)	0	100	100
5	Q	75/77 (97%)	74 (99%)	1 (1%)	0	100	100
5	R	74/77 (96%)	74 (100%)	0	0	100	100
5	S	75/77 (97%)	74 (99%)	1 (1%)	0	100	100
6	W	171/173 (99%)	165 (96%)	5 (3%)	1 (1%)	22	59
7	X	151/155 (97%)	148 (98%)	3 (2%)	0	100	100
7	Y	153/155 (99%)	151 (99%)	2 (1%)	0	100	100
8	a	264/266 (99%)	253 (96%)	11 (4%)	0	100	100
All	All	4793/4853 (99%)	4683 (98%)	107 (2%)	3 (0%)	50	82

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	G	205	TYR
6	W	148	ALA
2	F	265	VAL

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	400/402 (100%)	396 (99%)	4 (1%)	73	81
1	B	395/402 (98%)	392 (99%)	3 (1%)	79	84
1	C	399/402 (99%)	394 (99%)	5 (1%)	65	77
2	D	380/380 (100%)	377 (99%)	3 (1%)	79	84
2	E	380/380 (100%)	377 (99%)	3 (1%)	79	84
2	F	379/380 (100%)	375 (99%)	4 (1%)	70	80
3	G	236/236 (100%)	230 (98%)	6 (2%)	42	62
4	H	107/107 (100%)	102 (95%)	5 (5%)	22	46
5	I	54/54 (100%)	54 (100%)	0	100	100
5	J	54/54 (100%)	54 (100%)	0	100	100
5	L	54/54 (100%)	54 (100%)	0	100	100
5	M	54/54 (100%)	54 (100%)	0	100	100
5	N	54/54 (100%)	54 (100%)	0	100	100
5	O	54/54 (100%)	53 (98%)	1 (2%)	52	69
5	P	54/54 (100%)	54 (100%)	0	100	100
5	Q	54/54 (100%)	54 (100%)	0	100	100
5	R	54/54 (100%)	53 (98%)	1 (2%)	52	69
5	S	54/54 (100%)	53 (98%)	1 (2%)	52	69
6	W	135/136 (99%)	134 (99%)	1 (1%)	81	86
7	X	117/120 (98%)	117 (100%)	0	100	100
7	Y	112/120 (93%)	111 (99%)	1 (1%)	75	83
8	a	232/232 (100%)	228 (98%)	4 (2%)	56	72
All	All	3812/3837 (99%)	3770 (99%)	42 (1%)	69	80

5 of 42 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	H	25	VAL
5	S	8	LEU
4	H	34	ILE
4	H	76	VAL
7	Y	46	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
3	G	135	GLN
8	a	14	HIS
4	H	38	HIS
8	a	20	GLN
7	X	64	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 5 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
11	ADP	E	500	-	24,29,29	0.90	0	29,45,45	1.17	3 (10%)
9	ATP	B	600	10	28,33,33	1.47	2 (7%)	34,52,52	0.76	1 (2%)
9	ATP	A	600	10	28,33,33	0.67	0	34,52,52	0.70	1 (2%)
11	ADP	D	501	10	24,29,29	0.96	1 (4%)	29,45,45	1.29	4 (13%)
9	ATP	C	600	10	28,33,33	0.69	0	34,52,52	0.60	1 (2%)
9	ATP	F	502	10	28,33,33	0.74	0	34,52,52	0.77	2 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	ADP	E	500	-	-	4/12/32/32	0/3/3/3
9	ATP	B	600	10	-	0/18/38/38	0/3/3/3
9	ATP	A	600	10	-	3/18/38/38	0/3/3/3
11	ADP	D	501	10	-	1/12/32/32	0/3/3/3
9	ATP	C	600	10	-	1/18/38/38	0/3/3/3
9	ATP	F	502	10	-	6/18/38/38	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	B	600	ATP	PB-O3B	5.19	1.65	1.59
9	B	600	ATP	PA-O3A	4.64	1.64	1.59
11	D	501	ADP	O4'-C1'	2.37	1.44	1.40

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	D	501	ADP	N3-C2-N1	-3.76	123.56	128.67
11	E	500	ADP	N3-C2-N1	-3.69	123.67	128.67
11	D	501	ADP	C4-C5-N7	-2.62	106.57	109.34
11	E	500	ADP	C4-C5-N7	-2.47	106.72	109.34
9	B	600	ATP	C5-C6-N6	2.45	124.05	120.31

There are no chirality outliers.

5 of 15 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
9	A	600	ATP	C5'-O5'-PA-O2A
9	C	600	ATP	C5'-O5'-PA-O1A
9	F	502	ATP	C5'-O5'-PA-O1A
11	E	500	ADP	C5'-O5'-PA-O1A
9	F	502	ATP	O4'-C4'-C5'-O5'


There are no ring outliers.

1 monomer is involved in 1 short contact:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	F	502	ATP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-27311. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 176




Y Index: 176




Z Index: 176


### 6.2.2 Raw map



X Index: 176



Y Index: 176




Z Index: 176

The images above show central slices of the map in three orthogonal directions.




## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 176




Y Index: 135




Z Index: 178


### 6.3.2 Raw map



X Index: 178



Y Index: 136




Z Index: 178


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.015. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

## 6.6 Mask visualisation [i](#)


This section was not generated. No masks/segmentation were deposited.



## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 257 nm<sup>3</sup>; this corresponds to an approximate mass of 232 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of 0.244  $\text{\AA}^{-1}$

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.244  $\text{\AA}^{-1}$

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	4.10	-	-
Author-provided FSC curve	4.13	7.19	4.24
Unmasked-calculated*	7.82	13.18	8.12

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 7.82 differs from the reported value 4.1 by more than 10 %


## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-27311 and PDB model 8DBW. Per-residue inclusion information can be found in section [3](#) on page [9](#).


### 9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.015 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.015).

## 9.4 Atom inclusion [i](#)

















































At the recommended contour level, 77% of all backbone atoms, 58% of all non-hydrogen atoms, are inside the map.



## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.015) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.5830	 0.2910
A	 0.7340	 0.3760
B	 0.7140	 0.3650
C	 0.7280	 0.3770
D	 0.6870	 0.3520
E	 0.7290	 0.3810
F	 0.7500	 0.3950
G	 0.6450	 0.3060
H	 0.3180	 0.1450
I	 0.1590	 0.0580
J	 0.2370	 0.0430
L	 0.2030	 0.0660
M	 0.3210	 0.1080
N	 0.2920	 0.0810
O	 0.2500	 0.0890
P	 0.1960	 0.0640
Q	 0.2250	 0.0590
R	 0.2980	 0.0930
S	 0.3060	 0.1440
W	 0.6520	 0.3020
X	 0.3310	 0.2060
Y	 0.4120	 0.2100
a	 0.2550	 0.1160

