



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2024 – 04:29 PM EDT

PDB ID : 5DF7  
Title : CRYSTAL STRUCTURE OF PENICILLIN-BINDING PROTEIN 3 FROM PSEUDOMONAS AERUGINOSA IN COMPLEX WITH AZLOCILLIN  
Authors : Ren, J.; Nettleship, J.E.; Males, A.; Stuart, D.I.; Owens, R.J.  
Deposited on : 2015-08-26  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	564	<div> <div>3%</div> <div>84%</div> <div>7%</div> <div>9%</div> </div>
1	B	564	<div> <div>7%</div> <div>79%</div> <div>7%</div> <div>13%</div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8127 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cell division protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	515	Total	C	N	O	S	0	0	0
			3927	2477	713	725	12			
1	B	490	Total	C	N	O	S	0	0	0
			3722	2352	667	691	12			

There are 38 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	16	MET	-	engineered mutation	UNP Q51504
A	17	ALA	-	expression tag	UNP Q51504
A	18	HIS	-	expression tag	UNP Q51504
A	19	HIS	-	expression tag	UNP Q51504
A	20	HIS	-	expression tag	UNP Q51504
A	21	HIS	-	expression tag	UNP Q51504
A	22	HIS	-	expression tag	UNP Q51504
A	23	HIS	-	expression tag	UNP Q51504
A	24	SER	-	expression tag	UNP Q51504
A	25	SER	-	expression tag	UNP Q51504
A	26	GLY	-	expression tag	UNP Q51504
A	27	LEU	-	expression tag	UNP Q51504
A	28	GLU	-	expression tag	UNP Q51504
A	29	VAL	-	expression tag	UNP Q51504
A	30	LEU	-	expression tag	UNP Q51504
A	31	PHE	-	expression tag	UNP Q51504
A	32	GLN	-	expression tag	UNP Q51504
A	33	GLY	-	expression tag	UNP Q51504
A	34	PRO	-	expression tag	UNP Q51504
B	16	MET	-	engineered mutation	UNP Q51504
B	17	ALA	-	expression tag	UNP Q51504
B	18	HIS	-	expression tag	UNP Q51504
B	19	HIS	-	expression tag	UNP Q51504
B	20	HIS	-	expression tag	UNP Q51504
B	21	HIS	-	expression tag	UNP Q51504

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	22	HIS	-	expression tag	UNP Q51504
B	23	HIS	-	expression tag	UNP Q51504
B	24	SER	-	expression tag	UNP Q51504
B	25	SER	-	expression tag	UNP Q51504
B	26	GLY	-	expression tag	UNP Q51504
B	27	LEU	-	expression tag	UNP Q51504
B	28	GLU	-	expression tag	UNP Q51504
B	29	VAL	-	expression tag	UNP Q51504
B	30	LEU	-	expression tag	UNP Q51504
B	31	PHE	-	expression tag	UNP Q51504
B	32	GLN	-	expression tag	UNP Q51504
B	33	GLY	-	expression tag	UNP Q51504
B	34	PRO	-	expression tag	UNP Q51504

- Molecule 2 is (2R,4S)-5,5-dimethyl-2-[(1R)-2-oxo-1-[[[(2R)-{[(2-oxoimidazolidin-1-yl)carbonyl]amino}-2-phenylacetyl]amino}ethyl]-1,3-thiazolidine-4-carboxylic acid (three-letter code: 59H) (formula: C<sub>20</sub>H<sub>25</sub>N<sub>5</sub>O<sub>6</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			32	20	5	6	1		
2	B	1	Total	C	N	O	S	0	0
			32	20	5	6	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		
4	B	3	Total	Cl	0	0
			3	3		

- Molecule 5 is IMIDAZOLE (three-letter code: IMD) (formula: C<sub>3</sub>H<sub>5</sub>N<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	N	0	0
			5	3	2		
5	B	1	Total	C	N	0	0
			5	3	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	218	Total	O	0	0
			218	218		
6	B	164	Total	O	0	0
			164	164		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Cell division protein



#### • Molecule 1: Cell division protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	57.25Å 74.92Å 82.72Å 71.26° 85.99° 85.69°	Depositor
Resolution (Å)	47.19 – 2.00 47.19 – 2.00	Depositor EDS
% Data completeness (in resolution range)	86.7 (47.19-2.00) 86.7 (47.19-2.00)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.89 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.198 , 0.237 0.197 , 0.236	Depositor DCC
$R_{free}$ test set	3803 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.2	Xtriage
Anisotropy	0.287	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 55.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	8127	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IMD, 59H, CL, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.34	0/4004	0.54	0/5433
1	B	0.33	0/3797	0.52	0/5154
All	All	0.34	0/7801	0.53	0/10587

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3927	0	3988	21	0
1	B	3722	0	3760	26	0
2	A	32	0	23	0	0
2	B	32	0	23	0	0
3	A	6	0	8	2	0
3	B	12	0	16	5	0
4	A	1	0	0	0	0
4	B	3	0	0	0	0
5	A	5	0	5	0	0
5	B	5	0	5	0	0
6	A	218	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	164	0	0	1	0
All	All	8127	0	7828	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:269:ASN:H	3:B:603:GOL:H31	1.56	0.70
1:B:321:GLY:HA3	1:B:335:ARG:HD3	1.78	0.66
1:B:562:THR:HG21	6:B:803:HOH:O	1.99	0.62
1:A:51:ARG:HH11	1:A:198:ASP:HB2	1.65	0.62
1:A:478:GLY:H	1:A:562:THR:HG23	1.66	0.60
1:A:518:ILE:HD12	1:A:550:LEU:HD23	1.84	0.59
1:B:478:GLY:H	1:B:562:THR:HG23	1.67	0.59
1:A:262:MET:SD	1:A:416:ILE:HD11	2.44	0.57
1:A:169:ASP:HB3	1:A:175:ARG:HG3	1.88	0.56
1:B:268:TYR:HB2	3:B:603:GOL:H32	1.88	0.55
1:B:322:THR:HG22	1:B:331:ARG:HG2	1.87	0.55
1:A:168:THR:HB	1:A:172:ASP:HA	1.88	0.54
1:A:301:MET:HG2	1:A:460:MET:HE1	1.90	0.53
1:B:86:PRO:O	1:B:90:MET:HG2	2.10	0.52
1:B:259:ILE:HD11	1:B:416:ILE:HD11	1.91	0.51
1:B:191:GLY:HA2	1:B:211:LYS:HG2	1.92	0.51
1:B:354:ILE:HG21	1:B:407:TYR:HB3	1.93	0.51
1:A:120:ARG:NH1	1:B:338:ARG:HD3	2.26	0.50
1:B:560:LEU:N	1:B:561:PRO:HD2	2.27	0.49
1:A:181:ALA:HB2	1:A:384:PRO:HD3	1.95	0.48
1:B:181:ALA:HB2	1:B:384:PRO:HD3	1.96	0.48
1:A:525:ASP:OD1	3:A:602:GOL:H11	2.14	0.47
1:B:173:ARG:HH11	1:B:175:ARG:HH11	1.62	0.47
1:A:354:ILE:HG21	1:A:407:TYR:HB3	1.98	0.46
1:B:523:VAL:HG11	3:B:602:GOL:H11	1.97	0.46
1:A:525:ASP:CG	3:A:602:GOL:H11	2.35	0.46
1:B:58:ILE:HD11	1:B:193:ARG:HB3	1.97	0.46
1:B:83:TRP:CE2	1:B:146:TYR:HB2	2.51	0.45
1:B:343:THR:HA	1:B:460:MET:HE2	1.98	0.45
1:A:51:ARG:NH1	1:A:198:ASP:HB2	2.31	0.44
1:A:86:PRO:O	1:A:90:MET:HG2	2.17	0.44
1:A:518:ILE:HD11	1:A:553:MET:SD	2.57	0.44

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:342:LEU:HA	1:B:345:ILE:HD12	2.00	0.44
1:B:256:THR:HG22	1:B:442:VAL:HG22	1.99	0.44
1:B:67:ASP:HB2	1:B:223:ILE:O	2.18	0.43
1:B:248:SER:HB2	3:B:602:GOL:H12	1.99	0.43
1:A:186:LEU:O	1:A:216:GLY:HA3	2.19	0.43
1:A:95:ARG:NH1	1:A:140:LEU:O	2.43	0.43
1:A:283:ASN:ND2	1:A:286:MET:HG3	2.34	0.42
1:A:222:SER:HB2	1:A:258:GLU:HB3	2.02	0.42
1:A:560:LEU:N	1:A:561:PRO:HD2	2.35	0.41
1:B:298:PRO:HG3	1:B:461:LEU:HD21	2.01	0.41
1:B:86:PRO:HG2	1:B:120:ARG:O	2.20	0.41
1:B:442:VAL:HA	1:B:443:PRO:HD3	1.88	0.40
1:A:88:GLU:CD	1:A:202:ARG:HB3	2.41	0.40
1:B:269:ASN:H	3:B:603:GOL:C3	2.28	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	513/564 (91%)	500 (98%)	12 (2%)	1 (0%)	47	44
1	B	486/564 (86%)	474 (98%)	11 (2%)	1 (0%)	47	44
All	All	999/1128 (89%)	974 (98%)	23 (2%)	2 (0%)	47	44

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	172	ASP
1	B	93	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	410/448 (92%)	402 (98%)	8 (2%)	55	58
1	B	387/448 (86%)	386 (100%)	1 (0%)	92	95
All	All	797/896 (89%)	788 (99%)	9 (1%)	73	78

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	67	ASP
1	A	161	VAL
1	A	171	ASP
1	A	173	ARG
1	A	349	SER
1	A	354	ILE
1	A	416	ILE
1	A	441	ARG
1	B	175	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	212	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
5	IMD	B	607	-	3,5,5	0.31	0	4,5,5	0.65	0
2	59H	B	601	1	29,34,34	2.85	9 (31%)	36,49,49	2.28	9 (25%)
3	GOL	B	602	-	5,5,5	0.18	0	5,5,5	0.61	0
3	GOL	B	603	-	5,5,5	0.30	0	5,5,5	0.38	0
2	59H	A	601	1	29,34,34	2.78	9 (31%)	36,49,49	2.11	10 (27%)
5	IMD	A	604	-	3,5,5	0.32	0	4,5,5	0.62	0
3	GOL	A	602	-	5,5,5	0.32	0	5,5,5	0.96	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	IMD	B	607	-	-	-	0/1/1/1
2	59H	B	601	1	-	3/24/55/55	0/3/3/3
3	GOL	B	602	-	-	2/4/4/4	-
3	GOL	B	603	-	-	4/4/4/4	-
2	59H	A	601	1	-	3/24/55/55	0/3/3/3
5	IMD	A	604	-	-	-	0/1/1/1
3	GOL	A	602	-	-	2/4/4/4	-

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	59H	CX2-CX1	8.77	1.53	1.39
2	A	601	59H	CX2-CX1	8.72	1.53	1.39
2	B	601	59H	CX5-CX6	5.84	1.51	1.38
2	A	601	59H	CX5-CX6	5.76	1.51	1.38
2	B	601	59H	C-N1	4.82	1.44	1.34
2	A	601	59H	C-N1	4.67	1.44	1.34
2	A	601	59H	CX7-N	4.56	1.45	1.35
2	B	601	59H	CX7-N	4.47	1.45	1.35
2	B	601	59H	CXD-S4	-4.44	1.74	1.84
2	B	601	59H	CX4-CX3	4.12	1.48	1.38
2	A	601	59H	CX4-CX3	3.98	1.48	1.38
2	A	601	59H	CXD-S4	-3.96	1.75	1.84
2	B	601	59H	CXC-S4	-3.34	1.78	1.85
2	A	601	59H	CXC-S4	-2.93	1.79	1.85
2	A	601	59H	CXA-NX2	2.83	1.43	1.34
2	B	601	59H	CXA-NX2	2.82	1.43	1.34
2	B	601	59H	CX6-CX1	-2.19	1.35	1.39
2	A	601	59H	CX6-CX1	-2.08	1.35	1.39

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	59H	CX9-NX2-CXA	-6.38	107.06	113.60
2	B	601	59H	CX8-NX1-CX7	5.68	127.30	118.66
2	A	601	59H	CX9-NX2-CXA	-4.96	108.51	113.60
2	A	601	59H	CX8-NX1-CX7	4.94	126.17	118.66
2	B	601	59H	CX1-CA-C	4.81	118.96	108.16
2	A	601	59H	CXD-CA1-N1	4.47	119.71	109.98
2	B	601	59H	CXD-CA1-N1	3.94	118.57	109.98
2	A	601	59H	CX1-CA-C	3.82	116.75	108.16
2	B	601	59H	CA3-CXC-S4	-3.39	97.09	103.81
2	A	601	59H	N-CX7-NX1	3.31	118.85	114.04
2	B	601	59H	CA1-N1-C	-3.27	119.55	123.12
2	A	601	59H	CX9-CX8-NX1	3.26	105.48	102.93
2	A	601	59H	CA3-CXC-S4	-2.89	98.09	103.81
2	B	601	59H	N-CX7-NX1	2.87	118.22	114.04
2	B	601	59H	CXC-S4-CXD	2.52	99.38	93.99
2	A	601	59H	CX8-CX9-NX2	2.47	105.28	102.54
2	B	601	59H	CX9-CX8-NX1	2.41	104.82	102.93
2	A	601	59H	C3-CA3-N3	-2.27	107.16	110.70
2	A	601	59H	CXC-S4-CXD	2.24	98.79	93.99

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	602	GOL	C1-C2-C3-O3
3	A	602	GOL	O2-C2-C3-O3
3	B	602	GOL	O1-C1-C2-O2
3	B	602	GOL	O1-C1-C2-C3
3	B	603	GOL	O1-C1-C2-C3
3	B	603	GOL	O2-C2-C3-O3
2	B	601	59H	OXT-C3-CA3-N3
2	A	601	59H	C1-CA1-N1-C
3	B	603	GOL	O1-C1-C2-O2
2	B	601	59H	O3-C3-CA3-N3
3	B	603	GOL	C1-C2-C3-O3
2	A	601	59H	O3-C3-CA3-N3
2	A	601	59H	OXT-C3-CA3-N3
2	B	601	59H	N-CA-CX1-CX2


There are no ring outliers.

3 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	602	GOL	2	0
3	B	603	GOL	3	0
3	A	602	GOL	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

## Ligand 59H B 601







## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	515/564 (91%)	-0.02	16 (3%) 49 48	15, 33, 68, 111	0
1	B	490/564 (86%)	0.29	40 (8%) 11 11	16, 37, 79, 104	0
All	All	1005/1128 (89%)	0.14	56 (5%) 24 23	15, 34, 76, 111	0

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	213	ALA	6.6
1	A	50	ALA	6.1
1	A	564	THR	5.9
1	B	192	LYS	4.8
1	B	146	TYR	4.8
1	B	89	LEU	4.5
1	B	211	LYS	4.3
1	B	139	ALA	4.2
1	B	83	TRP	4.2
1	B	328	TYR	4.2
1	B	119	GLU	4.1
1	B	58	ILE	4.1
1	B	122	PHE	4.0
1	B	135	GLU	3.9
1	B	57	ALA	3.7
1	B	120	ARG	3.7
1	A	199	ARG	3.7
1	B	441	ARG	3.6
1	B	212	ASN	3.6
1	B	190	PRO	3.5
1	B	92	ALA	3.5
1	B	143	PRO	3.5
1	B	124	TYR	3.4
1	A	441	ARG	3.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	563	ALA	3.2
1	A	51	ARG	3.1
1	B	193	ARG	3.1
1	A	200	ARG	3.0
1	B	84	ALA	3.0
1	B	560	LEU	2.9
1	B	118	ALA	2.8
1	B	87	LYS	2.8
1	A	560	LEU	2.7
1	A	561	PRO	2.7
1	A	444	ASP	2.7
1	A	440	ASP	2.6
1	B	442	VAL	2.6
1	B	189	VAL	2.6
1	A	139	ALA	2.5
1	B	91	THR	2.5
1	B	59	PRO	2.5
1	B	140	LEU	2.4
1	A	395	ARG	2.4
1	A	140	LEU	2.3
1	A	442	VAL	2.3
1	B	116	GLN	2.3
1	A	124	TYR	2.3
1	B	327	ARG	2.3
1	B	123	ILE	2.3
1	B	514	THR	2.1
1	B	440	ASP	2.1
1	B	90	MET	2.1
1	B	102	ALA	2.0
1	B	88	GLU	2.0
1	A	148	ILE	2.0
1	B	129	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	GOL	B	603	6/6	0.85	0.17	47,53,57,59	0
3	GOL	B	602	6/6	0.91	0.19	29,39,48,48	0
5	IMD	B	607	5/5	0.92	0.13	48,55,62,63	0
5	IMD	A	604	5/5	0.94	0.07	49,52,57,59	0
3	GOL	A	602	6/6	0.94	0.16	21,26,32,38	0
4	CL	A	603	1/1	0.95	0.10	55,55,55,55	0
4	CL	B	604	1/1	0.96	0.06	47,47,47,47	0
2	59H	A	601	32/32	0.96	0.10	23,29,35,38	0
2	59H	B	601	32/32	0.96	0.11	28,34,37,41	0
4	CL	B	605	1/1	0.98	0.06	39,39,39,39	0
4	CL	B	606	1/1	0.99	0.07	35,35,35,35	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




### Electron density around 59H A 601:

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around 59H B 601:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.