



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 23, 2024 – 02:02 AM EDT

PDB ID : 6DHQ
Title : Bovine glutamate dehydrogenase complexed with NADPH, glutamate, and GTP
Authors : Smith, T.J.
Deposited on : 2018-05-20
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	501	<div> <div>10%</div> <div>68%</div> <div>25%</div> <div>6%</div> </div>
1	B	501	<div> <div>9%</div> <div>69%</div> <div>27%</div> <div>.</div> </div>
1	C	501	<div> <div>12%</div> <div>67%</div> <div>27%</div> <div>5%</div> </div>
1	D	501	<div> <div>9%</div> <div>70%</div> <div>26%</div> <div>.</div> </div>
1	E	501	<div> <div>12%</div> <div>70%</div> <div>24%</div> <div>5%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	501	<div><div></div><div>11%</div><div>67%</div><div>28%</div><div></div></div>

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 24419 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate dehydrogenase 1, mitochondrial.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	501	Total	C	N	O	S	0	0	0
			3916	2473	687	737	19			
1	B	501	Total	C	N	O	S	0	0	0
			3916	2473	687	737	19			
1	C	501	Total	C	N	O	S	0	0	0
			3916	2473	687	737	19			
1	D	501	Total	C	N	O	S	0	0	0
			3916	2473	687	737	19			
1	E	501	Total	C	N	O	S	0	0	0
			3916	2473	687	737	19			
1	F	501	Total	C	N	O	S	0	0	0
			3916	2473	687	737	19			

- Molecule 2 is GLUTAMIC ACID (three-letter code: GLU) (formula: C₅H₉NO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 10	C 5	N 1	O 4	0	0
2	B	1	Total 10	C 5	N 1	O 4	0	0
2	C	1	Total 10	C 5	N 1	O 4	0	0
2	D	1	Total 10	C 5	N 1	O 4	0	0
2	E	1	Total 10	C 5	N 1	O 4	0	0
2	F	1	Total 10	C 5	N 1	O 4	0	0

- Molecule 3 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 48	C 21	N 7	O 17	P 3	0	0
3	B	1	Total 48	C 21	N 7	O 17	P 3	0	0
3	C	1	Total 48	C 21	N 7	O 17	P 3	0	0
3	D	1	Total 48	C 21	N 7	O 17	P 3	0	0
3	E	1	Total 48	C 21	N 7	O 17	P 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	F	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 4 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
4	B	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
4	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
4	D	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
4	E	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
4	F	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	62	Total O 62 62	0	0
5	B	83	Total O 83 83	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	50	Total 50	O 50	0	0
5	D	66	Total 66	O 66	0	0
5	E	58	Total 58	O 58	0	0
5	F	64	Total 64	O 64	0	0


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamate dehydrogenase 1, mitochondrial




- Molecule 1: Glutamate dehydrogenase 1, mitochondrial







- Molecule 1: Glutamate dehydrogenase 1, mitochondrial




- Molecule 1: Glutamate dehydrogenase 1, mitochondrial



- Molecule 1: Glutamate dehydrogenase 1, mitochondrial



- Molecule 1: Glutamate dehydrogenase 1, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	123.25Å 101.75Å 167.85Å 90.00° 102.22° 90.00°	Depositor
Resolution (Å)	29.55 – 2.30 29.55 – 2.30	Depositor EDS
% Data completeness (in resolution range)	60.2 (29.55-2.30) 60.2 (29.55-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 2.31Å)	Xtriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, R_{free}	0.224 , 0.262 0.224 , 0.262	Depositor DCC
R_{free} test set	1998 reflections (1.84%)	wwPDB-VP
Wilson B-factor (Å ²)	45.9	Xtriage
Anisotropy	0.666	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 57.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	24419	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.03	19/3999 (0.5%)	1.22	54/5396 (1.0%)
1	B	0.79	10/3999 (0.3%)	1.05	37/5396 (0.7%)
1	C	0.83	17/3999 (0.4%)	1.16	49/5396 (0.9%)
1	D	0.80	12/3999 (0.3%)	1.20	42/5396 (0.8%)
1	E	1.21	34/3999 (0.9%)	1.30	59/5396 (1.1%)
1	F	0.83	16/3999 (0.4%)	1.18	45/5396 (0.8%)
All	All	0.93	108/23994 (0.5%)	1.19	286/32376 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	3
1	B	0	2
1	C	0	7
1	D	0	3
1	E	0	4
1	F	0	6
All	All	0	25

The worst 5 of 108 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	401	TYR	CE2-CZ	-21.36	1.10	1.38
1	A	401	TYR	CE1-CZ	-21.11	1.11	1.38
1	A	401	TYR	CG-CD1	-20.11	1.13	1.39
1	A	401	TYR	CG-CD2	-19.33	1.14	1.39
1	E	78	TYR	CE1-CZ	-18.69	1.14	1.38

The worst 5 of 286 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	44	ARG	NE-CZ-NH2	30.45	135.53	120.30
1	E	42	ARG	NE-CZ-NH2	25.93	133.26	120.30
1	F	439	ARG	NE-CZ-NH2	23.18	131.89	120.30
1	D	396	ARG	NE-CZ-NH1	-23.13	108.73	120.30
1	B	67	ARG	NE-CZ-NH2	23.05	131.83	120.30

There are no chirality outliers.

5 of 25 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	39	GLU	Peptide
1	A	499	THR	Peptide
1	A	86	ARG	Peptide
1	B	37	THR	Peptide
1	B	86	ARG	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3916	0	3880	161	0
1	B	3916	0	3880	120	0
1	C	3916	0	3880	141	0
1	D	3916	0	3880	129	0
1	E	3916	0	3880	137	0
1	F	3916	0	3880	134	0
2	A	10	0	5	1	0
2	B	10	0	5	2	0
2	C	10	0	5	2	0
2	D	10	0	5	1	0
2	E	10	0	5	1	0
2	F	10	0	5	1	0
3	A	48	0	26	4	0
3	B	48	0	26	3	0
3	C	48	0	26	4	0
3	D	48	0	26	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	48	0	26	4	0
3	F	48	0	26	5	0
4	A	32	0	12	0	0
4	B	32	0	12	1	0
4	C	32	0	12	0	0
4	D	32	0	12	1	0
4	E	32	0	12	0	0
4	F	32	0	12	1	0
5	A	62	0	0	10	0
5	B	83	0	0	8	1
5	C	50	0	0	10	0
5	D	66	0	0	17	0
5	E	58	0	0	5	1
5	F	64	0	0	20	0
All	All	24419	0	23538	756	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 756 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:363:ARG:HH21	1:A:365:ILE:CD1	1.25	1.48
1:A:363:ARG:NH2	1:A:365:ILE:HD11	1.36	1.40
1:E:458:GLU:CD	1:E:462:ARG:HH21	1.31	1.33
1:C:333:LYS:HE2	1:C:355:GLU:CB	1.63	1.26
1:C:333:LYS:HE2	1:C:355:GLU:CG	1.65	1.24

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:783:HOH:O	5:E:757:HOH:O[1_655]	1.81	0.39

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	499/501 (100%)	467 (94%)	25 (5%)	7 (1%)	11	11
1	B	499/501 (100%)	468 (94%)	23 (5%)	8 (2%)	9	9
1	C	499/501 (100%)	463 (93%)	25 (5%)	11 (2%)	6	5
1	D	499/501 (100%)	469 (94%)	22 (4%)	8 (2%)	9	9
1	E	499/501 (100%)	464 (93%)	24 (5%)	11 (2%)	6	5
1	F	499/501 (100%)	465 (93%)	26 (5%)	8 (2%)	9	9
All	All	2994/3006 (100%)	2796 (93%)	145 (5%)	53 (2%)	8	7

5 of 53 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	4	GLU
1	A	38	GLU
1	A	500	PHE
1	B	35	ARG
1	B	38	GLU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	417/417 (100%)	395 (95%)	22 (5%)	22	31
1	B	417/417 (100%)	403 (97%)	14 (3%)	37	51
1	C	417/417 (100%)	401 (96%)	16 (4%)	33	47

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	417/417 (100%)	399 (96%)	18 (4%)	29	40
1	E	417/417 (100%)	397 (95%)	20 (5%)	25	36
1	F	417/417 (100%)	404 (97%)	13 (3%)	40	55
All	All	2502/2502 (100%)	2399 (96%)	103 (4%)	30	43

5 of 103 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	86	ARG
1	E	9	PHE
1	F	405	SER
1	D	204	SER
1	D	419	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	205	GLN
1	E	205	GLN
1	F	484	ASN
1	E	414	GLN
1	F	205	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NDP	D	602	-	47,52,52	2.23	10 (21%)	61,80,80	1.81	19 (31%)
3	NDP	B	602	-	47,52,52	2.49	8 (17%)	61,80,80	1.62	13 (21%)
4	GTP	F	603	-	29,34,34	1.21	2 (6%)	35,54,54	1.59	8 (22%)
2	GLU	B	601	-	8,9,9	1.05	1 (12%)	8,11,11	1.76	2 (25%)
2	GLU	C	601	-	8,9,9	1.01	0	8,11,11	1.51	2 (25%)
4	GTP	C	603	-	29,34,34	1.31	5 (17%)	35,54,54	1.42	8 (22%)
2	GLU	E	601	-	8,9,9	1.24	2 (25%)	8,11,11	1.21	1 (12%)
4	GTP	A	603	-	29,34,34	1.20	2 (6%)	35,54,54	1.56	5 (14%)
3	NDP	A	602	-	47,52,52	2.78	11 (23%)	61,80,80	1.55	14 (22%)
2	GLU	A	601	-	8,9,9	1.19	2 (25%)	8,11,11	1.27	1 (12%)
4	GTP	B	603	-	29,34,34	1.46	5 (17%)	35,54,54	1.41	5 (14%)
4	GTP	E	603	-	29,34,34	1.06	2 (6%)	35,54,54	1.39	5 (14%)
3	NDP	C	602	-	47,52,52	2.53	13 (27%)	61,80,80	1.69	14 (22%)
3	NDP	E	602	-	47,52,52	2.98	12 (25%)	61,80,80	1.55	13 (21%)
3	NDP	F	602	-	47,52,52	2.49	10 (21%)	61,80,80	1.70	18 (29%)
2	GLU	F	601	-	8,9,9	1.05	0	8,11,11	1.76	1 (12%)
4	GTP	D	603	-	29,34,34	1.30	2 (6%)	35,54,54	1.49	7 (20%)
2	GLU	D	601	-	8,9,9	1.07	0	8,11,11	1.47	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NDP	D	602	-	-	10/30/77/77	0/5/5/5
3	NDP	B	602	-	-	6/30/77/77	0/5/5/5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GTP	F	603	-	-	5/18/38/38	0/3/3/3
2	GLU	B	601	-	-	2/9/9/9	-
2	GLU	C	601	-	-	2/9/9/9	-
4	GTP	C	603	-	-	2/18/38/38	0/3/3/3
2	GLU	E	601	-	-	4/9/9/9	-
4	GTP	A	603	-	-	2/18/38/38	0/3/3/3
3	NDP	A	602	-	-	3/30/77/77	0/5/5/5
2	GLU	A	601	-	-	4/9/9/9	-
4	GTP	B	603	-	-	2/18/38/38	0/3/3/3
4	GTP	E	603	-	-	4/18/38/38	0/3/3/3
3	NDP	C	602	-	-	6/30/77/77	0/5/5/5
3	NDP	E	602	-	-	11/30/77/77	0/5/5/5
3	NDP	F	602	-	-	5/30/77/77	0/5/5/5
2	GLU	F	601	-	-	2/9/9/9	-
4	GTP	D	603	-	-	2/18/38/38	0/3/3/3
2	GLU	D	601	-	-	4/9/9/9	-

The worst 5 of 87 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	NDP	P2B-O2B	13.88	1.83	1.59
3	E	602	NDP	P2B-O2B	13.25	1.82	1.59
3	F	602	NDP	P2B-O2B	12.94	1.82	1.59
3	C	602	NDP	P2B-O2B	11.87	1.80	1.59
3	D	602	NDP	P2B-O2B	11.64	1.79	1.59

The worst 5 of 137 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	602	NDP	O4B-C1B-N9A	-5.13	101.95	108.75
3	B	602	NDP	C4B-O4B-C1B	-4.36	105.93	109.92
4	A	603	GTP	O4'-C1'-N9	4.23	114.36	108.75
3	F	602	NDP	O2N-PN-O3	4.19	118.59	107.27
3	D	602	NDP	P2B-O2B-C2B	-4.02	112.69	123.43

There are no chirality outliers.

5 of 76 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	602	NDP	C5D-O5D-PN-O3
3	B	602	NDP	C5D-O5D-PN-O2N
3	B	602	NDP	O4D-C4D-C5D-O5D
3	C	602	NDP	O4D-C4D-C5D-O5D
3	D	602	NDP	C5B-O5B-PA-O1A


There are no ring outliers.

15 monomers are involved in 31 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	602	NDP	7	0
3	B	602	NDP	3	0
4	F	603	GTP	1	0
2	B	601	GLU	2	0
2	C	601	GLU	2	0
2	E	601	GLU	1	0
3	A	602	NDP	4	0
2	A	601	GLU	1	0
4	B	603	GTP	1	0
3	C	602	NDP	4	0
3	E	602	NDP	4	0
3	F	602	NDP	5	0
2	F	601	GLU	1	0
4	D	603	GTP	1	0
2	D	601	GLU	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.








Ligand GTP F 603






Ligand GTP C 603




















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	501/501 (100%)	0.48	51 (10%) 6 9	34, 55, 86, 115	0
1	B	501/501 (100%)	0.49	45 (8%) 9 12	34, 54, 86, 118	0
1	C	501/501 (100%)	0.53	62 (12%) 4 5	40, 57, 88, 112	0
1	D	501/501 (100%)	0.45	45 (8%) 9 12	39, 56, 89, 112	0
1	E	501/501 (100%)	0.63	61 (12%) 4 6	37, 62, 93, 113	0
1	F	501/501 (100%)	0.57	56 (11%) 5 7	38, 60, 90, 117	0
All	All	3006/3006 (100%)	0.52	320 (10%) 6 8	34, 57, 91, 118	0

The worst 5 of 320 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1	ALA	8.4
1	F	1	ALA	8.3
1	B	1	ALA	8.0
1	B	3	ARG	7.6
1	F	5	ASP	6.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NDP	F	602	48/48	0.92	0.16	51,68,78,204	0
2	GLU	D	601	10/10	0.95	0.19	48,54,60,62	0
2	GLU	E	601	10/10	0.95	0.14	50,56,64,67	0
2	GLU	F	601	10/10	0.95	0.18	50,57,65,66	0
2	GLU	C	601	10/10	0.95	0.20	43,50,63,65	0
3	NDP	D	602	48/48	0.96	0.13	38,55,61,67	0
3	NDP	E	602	48/48	0.96	0.10	52,63,74,81	0
3	NDP	C	602	48/48	0.96	0.13	44,54,62,67	0
2	GLU	A	601	10/10	0.97	0.16	48,52,60,67	0
2	GLU	B	601	10/10	0.97	0.17	42,50,56,57	0
3	NDP	A	602	48/48	0.97	0.13	39,51,60,63	0
3	NDP	B	602	48/48	0.97	0.13	38,50,59,64	0
4	GTP	A	603	32/32	0.97	0.12	41,53,60,65	0
4	GTP	C	603	32/32	0.97	0.12	48,56,65,69	0
4	GTP	E	603	32/32	0.97	0.15	55,62,71,76	0
4	GTP	D	603	32/32	0.98	0.10	46,54,63,66	0
4	GTP	B	603	32/32	0.98	0.11	42,49,59,62	0
4	GTP	F	603	32/32	0.98	0.12	53,60,67,73	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NDP F 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NDP D 602:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NDP E 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NDP C 602:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NDP A 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NDP B 602:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP A 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GTP C 603:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP E 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GTP D 603:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP B 603:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GTP F 603:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.