



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 22, 2025 – 02:47 PM EDT

PDB ID : 5DVZ
Title : Holo TrpB from *Pyrococcus furiosus*
Authors : Buller, A.R.; Arnold, F.H.
Deposited on : 2015-09-21
Resolution : 1.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.21
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.004 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.41.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	5161 (1.70-1.70)
Clashscore	180529	5671 (1.70-1.70)
Ramachandran outliers	177936	5594 (1.70-1.70)
Sidechain outliers	177891	5594 (1.70-1.70)
RSRZ outliers	164620	5159 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	396	<div> <div>%</div> <div>94%</div> <div>• •</div> </div>
1	B	396	<div> <div>%</div> <div>95%</div> <div>• •</div> </div>
1	C	396	<div> <div>4%</div> <div>94%</div> <div>• •</div> </div>
1	D	396	<div> <div>%</div> <div>94%</div> <div>• •</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12445 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tryptophan synthase beta chain 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	384	Total	C	N	O	P	S	0	2	0
			2952	1888	507	544	1	12			
1	B	382	Total	C	N	O	P	S	0	2	0
			2915	1867	498	537	1	12			
1	C	384	Total	C	N	O	P	S	0	3	0
			2936	1880	499	544	1	12			
1	D	385	Total	C	N	O	P	S	0	3	0
			2964	1895	506	550	1	12			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	389	LEU	-	expression tag	UNP Q8U093
A	390	GLU	-	expression tag	UNP Q8U093
A	391	HIS	-	expression tag	UNP Q8U093
A	392	HIS	-	expression tag	UNP Q8U093
A	393	HIS	-	expression tag	UNP Q8U093
A	394	HIS	-	expression tag	UNP Q8U093
A	395	HIS	-	expression tag	UNP Q8U093
A	396	HIS	-	expression tag	UNP Q8U093
B	389	LEU	-	expression tag	UNP Q8U093
B	390	GLU	-	expression tag	UNP Q8U093
B	391	HIS	-	expression tag	UNP Q8U093
B	392	HIS	-	expression tag	UNP Q8U093
B	393	HIS	-	expression tag	UNP Q8U093
B	394	HIS	-	expression tag	UNP Q8U093
B	395	HIS	-	expression tag	UNP Q8U093
B	396	HIS	-	expression tag	UNP Q8U093
C	389	LEU	-	expression tag	UNP Q8U093
C	390	GLU	-	expression tag	UNP Q8U093
C	391	HIS	-	expression tag	UNP Q8U093
C	392	HIS	-	expression tag	UNP Q8U093
C	393	HIS	-	expression tag	UNP Q8U093

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	394	HIS	-	expression tag	UNP Q8U093
C	395	HIS	-	expression tag	UNP Q8U093
C	396	HIS	-	expression tag	UNP Q8U093
D	389	LEU	-	expression tag	UNP Q8U093
D	390	GLU	-	expression tag	UNP Q8U093
D	391	HIS	-	expression tag	UNP Q8U093
D	392	HIS	-	expression tag	UNP Q8U093
D	393	HIS	-	expression tag	UNP Q8U093
D	394	HIS	-	expression tag	UNP Q8U093
D	395	HIS	-	expression tag	UNP Q8U093
D	396	HIS	-	expression tag	UNP Q8U093

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Na 1 1	0	0
2	B	1	Total Na 1 1	0	0
2	C	1	Total Na 1 1	0	0
2	D	1	Total Na 1 1	0	0

- Molecule 3 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O P 5 4 1	0	0
3	B	1	Total O P 5 4 1	0	0
3	C	1	Total O P 5 4 1	0	0
3	D	1	Total O P 5 4 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	164	Total O 164 164	0	0
4	B	158	Total O 158 158	0	0
4	C	156	Total O 156 156	0	1
4	D	176	Total O 176 176	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Tryptophan synthase beta chain 1



- Molecule 1: Tryptophan synthase beta chain 1



- Molecule 1: Tryptophan synthase beta chain 1



- Molecule 1: Tryptophan synthase beta chain 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	87.05Å 111.88Å 160.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.33 – 1.69 39.33 – 1.69	Depositor EDS
% Data completeness (in resolution range)	97.0 (39.33-1.69) 97.1 (39.33-1.69)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.46 (at 1.69Å)	Xtriage
Refinement program	REFMAC 5.8.0124	Depositor
R, R_{free}	0.202 , 0.228 0.209 , 0.234	Depositor DCC
R_{free} test set	8573 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	26.2	Xtriage
Anisotropy	0.242	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 25.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	12445	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP, NA, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/2994	0.56	0/4049
1	B	0.34	0/2956	0.56	0/4000
1	C	0.35	0/2978	0.55	0/4029
1	D	0.36	0/3009	0.58	0/4070
All	All	0.35	0/11937	0.56	0/16148

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2952	0	2936	6	0
1	B	2915	0	2889	2	0
1	C	2936	0	2893	4	0
1	D	2964	0	2941	4	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	5	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	0	0
4	A	164	0	0	1	0
4	B	158	0	0	0	0
4	C	156	0	0	0	0
4	D	176	0	0	0	0
All	All	12445	0	11659	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:283:GLN:NE2	4:A:501:HOH:O	2.36	0.57
1:A:211:ILE:HG21	1:A:219:PRO:HD3	1.96	0.47
1:D:211:ILE:HG21	1:D:219:PRO:HD3	1.97	0.47
1:D:10:TYR:O	1:D:276:GLY:HA2	2.16	0.46
1:A:382:LEU:HD21	1:B:138:LYS:HE2	1.99	0.44
1:C:17:GLU:HB3	1:C:18:PRO:HD3	2.00	0.44
1:A:351:ALA:HA	1:A:354:MET:HE2	2.00	0.43
1:A:188:GLY:HA2	1:A:275:HIS:O	2.18	0.43
1:B:351:ALA:HA	1:B:354:MET:HE2	2.01	0.42
1:C:128:TYR:OH	1:C:171:ASP:OD2	2.31	0.42
1:A:73:GLU:O	1:A:76[B]:VAL:HG22	2.20	0.42
1:D:110:HIS:CE1	1:D:184:GLY:HA2	2.55	0.42
1:C:351:ALA:HA	1:C:354:MET:CE	2.51	0.41
1:C:102:ILE:HA	1:C:126:ASP:O	2.21	0.41
1:D:188:GLY:HA2	1:D:275:HIS:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	383/396 (97%)	376 (98%)	6 (2%)	1 (0%)	37	23
1	B	379/396 (96%)	373 (98%)	5 (1%)	1 (0%)	37	23
1	C	382/396 (96%)	373 (98%)	8 (2%)	1 (0%)	37	23
1	D	385/396 (97%)	376 (98%)	9 (2%)	0	100	100
All	All	1529/1584 (96%)	1498 (98%)	28 (2%)	3 (0%)	44	29

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	186	VAL
1	B	186	VAL
1	C	186	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	295/315 (94%)	294 (100%)	1 (0%)	91	88
1	B	290/315 (92%)	289 (100%)	1 (0%)	91	88
1	C	291/315 (92%)	290 (100%)	1 (0%)	91	88
1	D	297/315 (94%)	294 (99%)	3 (1%)	73	64
All	All	1173/1260 (93%)	1167 (100%)	6 (0%)	86	82

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	104	GLU
1	B	270	GLN
1	C	150	ASN
1	D	104	GLU
1	D	283	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	285	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	270	GLN
1	A	283	GLN
1	B	270	GLN
1	C	283	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LLP	D	82	1	23,24,25	2.28	4 (17%)	25,32,34	1.46	6 (24%)
1	LLP	A	82	1	23,24,25	2.46	5 (21%)	25,32,34	1.53	5 (20%)
1	LLP	B	82	1	23,24,25	2.39	5 (21%)	25,32,34	1.43	5 (20%)
1	LLP	C	82	1	23,24,25	2.29	4 (17%)	25,32,34	1.36	3 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	D	82	1	-	4/16/17/19	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	82	1	-	2/16/17/19	0/1/1/1
1	LLP	B	82	1	-	1/16/17/19	0/1/1/1
1	LLP	C	82	1	-	2/16/17/19	0/1/1/1

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	82	LLP	C3-C2	7.28	1.48	1.41
1	B	82	LLP	C3-C2	6.70	1.47	1.41
1	D	82	LLP	C3-C2	6.50	1.47	1.41
1	C	82	LLP	C3-C2	6.09	1.47	1.41
1	A	82	LLP	C4-C5	5.27	1.49	1.42
1	B	82	LLP	C4'-NZ	5.25	1.44	1.27
1	A	82	LLP	C4'-NZ	5.13	1.44	1.27
1	C	82	LLP	C4'-NZ	5.03	1.44	1.27
1	D	82	LLP	C4'-NZ	5.02	1.44	1.27
1	C	82	LLP	C4-C5	5.02	1.49	1.42
1	B	82	LLP	C4-C5	4.88	1.48	1.42
1	C	82	LLP	C4-C3	4.87	1.49	1.41
1	D	82	LLP	C4-C5	4.76	1.48	1.42
1	B	82	LLP	C4-C3	4.72	1.48	1.41
1	A	82	LLP	C4-C3	4.63	1.48	1.41
1	D	82	LLP	C4-C3	4.33	1.48	1.41
1	A	82	LLP	C4-C4'	2.16	1.51	1.46
1	B	82	LLP	C4-C4'	2.14	1.51	1.46

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	82	LLP	C4-C3-C2	-4.65	117.53	120.14
1	C	82	LLP	C4-C3-C2	-3.76	118.03	120.14
1	D	82	LLP	C4-C3-C2	-3.15	118.37	120.14
1	B	82	LLP	C4-C3-C2	-2.81	118.56	120.14
1	D	82	LLP	C6-N1-C2	2.78	124.23	119.20
1	D	82	LLP	OP4-C5'-C5	2.65	114.32	109.36
1	C	82	LLP	C4-C4'-NZ	-2.52	112.42	124.04
1	A	82	LLP	O3-C3-C2	2.37	122.50	117.58
1	C	82	LLP	C6-N1-C2	2.34	123.44	119.20
1	B	82	LLP	OP3-P-OP2	2.32	116.52	107.80
1	B	82	LLP	C4-C4'-NZ	-2.31	113.38	124.04
1	A	82	LLP	C4-C4'-NZ	-2.23	113.77	124.04
1	A	82	LLP	C6-N1-C2	2.22	123.22	119.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	82	LLP	C4-C4'-NZ	-2.16	114.09	124.04
1	B	82	LLP	OP3-P-OP4	-2.13	101.13	106.67
1	B	82	LLP	OP4-C5'-C5	2.10	113.30	109.36
1	D	82	LLP	O3-C3-C2	2.08	121.90	117.58
1	D	82	LLP	OP3-P-OP2	2.04	115.44	107.80
1	A	82	LLP	OP3-P-OP4	-2.03	101.39	106.67

There are no chirality outliers.

All (9) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	82	LLP	O-C-CA-CB
1	C	82	LLP	O-C-CA-CB
1	D	82	LLP	O-C-CA-CB
1	A	82	LLP	C4-C4'-NZ-CE
1	D	82	LLP	C4-C4'-NZ-CE
1	B	82	LLP	C4-C4'-NZ-CE
1	C	82	LLP	C4-C4'-NZ-CE
1	D	82	LLP	C-CA-CB-CG
1	D	82	LLP	CG-CD-CE-NZ

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PO4	B	402	-	4,4,4	1.05	0	6,6,6	0.65	0
3	PO4	C	402	-	4,4,4	0.99	0	6,6,6	0.45	0
3	PO4	D	402	-	4,4,4	0.92	0	6,6,6	0.51	0
3	PO4	A	402	-	4,4,4	1.03	0	6,6,6	0.42	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	383/396 (96%)	0.09	5 (1%) 74 77	13, 30, 46, 63	2 (0%)
1	B	381/396 (96%)	0.10	5 (1%) 74 77	13, 30, 47, 78	2 (0%)
1	C	383/396 (96%)	0.20	14 (3%) 45 48	18, 32, 52, 75	3 (0%)
1	D	384/396 (96%)	-0.01	4 (1%) 79 81	12, 28, 45, 66	3 (0%)
All	All	1531/1584 (96%)	0.10	28 (1%) 67 70	12, 30, 49, 78	10 (0%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	274[A]	PHE	5.0
1	D	385	SER	3.5
1	D	291	PRO	3.2
1	C	289	ILE	2.9
1	B	289	ILE	2.9
1	C	287	GLY	2.8
1	B	384	VAL	2.8
1	D	260	GLY	2.7
1	B	291	PRO	2.7
1	B	287	GLY	2.6
1	C	300[A]	ASP	2.6
1	C	291	PRO	2.6
1	A	384	VAL	2.5
1	B	155	ASN	2.4
1	C	281	PHE	2.4
1	C	284	ASP	2.4
1	D	169	LEU	2.4
1	C	162	LYS	2.3
1	A	289	ILE	2.3
1	C	301	TYR	2.3
1	C	384	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	260	GLY	2.2
1	C	288	GLN	2.1
1	C	270	GLN	2.1
1	A	300	ASP	2.1
1	A	267	ASN	2.1
1	C	107	ALA	2.0
1	A	288	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	LLP	C	82	24/25	0.97	0.07	22,24,26,27	0
1	LLP	B	82	24/25	0.98	0.05	20,22,23,23	0
1	LLP	A	82	24/25	0.98	0.05	20,23,23,24	0
1	LLP	D	82	24/25	0.98	0.05	18,20,21,21	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PO4	B	402	5/5	0.86	0.14	53,53,54,55	0
3	PO4	C	402	5/5	0.88	0.12	62,63,65,66	0
3	PO4	D	402	5/5	0.90	0.13	47,48,48,51	0
3	PO4	A	402	5/5	0.94	0.08	36,36,38,39	0
2	NA	A	401	1/1	0.95	0.10	29,29,29,29	0
2	NA	C	401	1/1	0.95	0.07	23,23,23,23	0
2	NA	D	401	1/1	0.97	0.07	19,19,19,19	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NA	B	401	1/1	0.98	0.08	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.