



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 19, 2025 – 09:59 AM EDT

PDB ID : 3DWG
Title : Crystal structure of a sulfur carrier protein complex found in the cysteine biosynthetic pathway of Mycobacterium tuberculosis
Authors : Jurgenson, C.T.; Burns, K.E.; Begley, T.P.; Ealick, S.E.
Deposited on : 2008-07-22
Resolution : 1.53 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.21
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.004 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.41.4


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3511 (1.56-1.52)
Clashscore	180529	3784 (1.56-1.52)
Ramachandran outliers	177936	3720 (1.56-1.52)
Sidechain outliers	177891	3717 (1.56-1.52)
RSRZ outliers	164620	3510 (1.56-1.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	325	<div> <div>4%</div> <div>92%</div> <div>7%</div> </div>
1	B	325	<div> <div>12%</div> <div>83%</div> <div>14%</div> <div>..</div> </div>
2	C	93	<div> <div>2%</div> <div>85%</div> <div>13%</div> <div>..</div> </div>

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6466 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cysteine synthase B.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	325	Total	C	N	O	S	0	1	0
			2436	1532	436	459	9			
1	B	321	Total	C	N	O	S	0	4	0
			2366	1489	415	451	11			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	ARG	-	expression tag	UNP P63873
A	0	HIS	-	expression tag	UNP P63873
B	-1	ARG	-	expression tag	UNP P63873
B	0	HIS	-	expression tag	UNP P63873

- Molecule 2 is a protein called 9.5 kDa culture filtrate antigen cfp10A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	92	Total	C	N	O	S	0	1	0
			649	404	110	134	1			

- Molecule 3 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C₈H₁₀NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
3	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	438	Total	O	0	0
			438	438		
4	B	409	Total	O	0	0
			409	409		
4	C	138	Total	O	0	0
			138	138		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Cysteine synthase B



• Molecule 1: Cysteine synthase B



• Molecule 2: 9.5 kDa culture filtrate antigen cfp10A



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.81Å 80.43Å 89.62Å 90.00° 105.74° 90.00°	Depositor
Resolution (Å)	27.13 – 1.53 27.13 – 1.53	Depositor EDS
% Data completeness (in resolution range)	96.2 (27.13-1.53) 96.3 (27.13-1.53)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.10 (at 1.53Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.176 , 0.217 0.177 , 0.216	Depositor DCC
R_{free} test set	5508 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	19.1	Xtriage
Anisotropy	0.070	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 56.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.020 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6466	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.72% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.04	0/2487	1.14	9/3384 (0.3%)
1	B	0.94	2/2420 (0.1%)	0.95	7/3296 (0.2%)
2	C	0.88	0/662	0.90	2/904 (0.2%)
All	All	0.98	2/5569 (0.0%)	1.03	18/7584 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	49	SER	CB-OG	7.06	1.51	1.42
1	B	226	PHE	CB-CG	-5.19	1.42	1.51

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	239	ARG	NE-CZ-NH1	19.90	130.25	120.30
1	A	239	ARG	NE-CZ-NH2	-18.91	110.85	120.30
1	A	239	ARG	CD-NE-CZ	10.52	138.33	123.60
1	A	239	ARG	CB-CG-CD	7.94	132.25	111.60
1	A	53	ARG	NE-CZ-NH2	-6.97	116.81	120.30
1	A	109	ARG	NE-CZ-NH2	-6.81	116.89	120.30
2	C	67	ASP	CB-CG-OD1	6.02	123.72	118.30
1	A	286	ARG	NE-CZ-NH2	-6.02	117.29	120.30
1	B	144	PRO	N-CA-CB	5.50	109.90	103.30
1	B	270[A]	LEU	CB-CG-CD2	5.40	120.19	111.00
1	B	270[B]	LEU	CB-CG-CD2	5.40	120.19	111.00
1	A	88	MET	CG-SD-CE	-5.33	91.67	100.20
1	B	232	ASP	CB-CG-OD1	5.27	123.04	118.30
1	B	239	ARG	NE-CZ-NH1	5.24	122.92	120.30
1	B	21	ARG	NE-CZ-NH1	-5.20	117.70	120.30
1	A	44	ARG	NE-CZ-NH2	-5.18	117.71	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	36	ARG	NE-CZ-NH2	-5.12	117.74	120.30
2	C	82	ASP	CB-CG-OD1	5.05	122.85	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2436	0	2415	30	0
1	B	2366	0	2308	55	0
2	C	649	0	628	15	0
3	A	15	0	7	0	0
3	B	15	0	7	0	0
4	A	438	0	0	13	0
4	B	409	0	0	19	0
4	C	138	0	0	3	0
All	All	6466	0	5365	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:190[A]:MET:HE3	4:B:1004:HOH:O	1.22	1.39
1:A:222:MET:CE	2:C:59:VAL:HG22	1.53	1.38
1:B:190[B]:MET:HG2	4:B:837:HOH:O	1.38	1.20
1:A:222:MET:HE2	2:C:59:VAL:HG22	1.17	1.14
1:A:209:GLU:OE2	1:A:239:ARG:HD3	1.46	1.11
1:B:99[B]:CYS:SG	1:B:101:MET:CE	2.42	1.07
1:A:222:MET:CE	2:C:59:VAL:CG2	2.38	1.01
1:B:190[A]:MET:HG2	4:B:837:HOH:O	1.64	0.96
1:A:222:MET:HE3	2:C:59:VAL:HG22	1.51	0.92
1:B:99[B]:CYS:SG	1:B:101:MET:HE2	2.08	0.91

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:198:GLU:HG3	4:B:1007:HOH:O	1.71	0.90
1:A:69:ARG:HD3	4:A:541:HOH:O	1.72	0.89
1:A:222:MET:HE3	2:C:59:VAL:CG2	2.02	0.86
1:A:222:MET:HE2	2:C:59:VAL:CG2	2.02	0.84
1:B:190[A]:MET:CE	4:B:1004:HOH:O	1.94	0.83
1:A:126:GLU:HG3	4:A:766:HOH:O	1.79	0.83
2:C:74:LEU:HG	4:C:227:HOH:O	1.80	0.81
1:B:110:ARG:HD2	1:B:122:PHE:HZ	1.46	0.78
1:B:110:ARG:CD	1:B:122:PHE:HZ	1.97	0.78
1:A:209:GLU:OE2	1:A:239:ARG:CD	2.32	0.76
1:A:115:LEU:O	1:B:256:HIS:HE1	1.68	0.75
1:B:250:ARG:HE	1:B:271:HIS:HD2	1.32	0.75
1:B:31:ASP:CB	4:B:998:HOH:O	2.36	0.73
1:A:34:HIS:HE1	1:A:288:ASP:OD2	1.73	0.72
1:B:190[A]:MET:SD	4:B:1004:HOH:O	2.38	0.72
1:B:91:ARG:HB3	4:B:1001:HOH:O	1.90	0.72
1:A:197:ARG:NH1	4:A:745:HOH:O	2.18	0.70
1:B:99[B]:CYS:SG	1:B:101:MET:HE3	2.32	0.69
1:B:22:LEU:HD21	1:B:270[A]:LEU:HD11	1.75	0.69
1:A:31:ASP:O	4:A:831:HOH:O	2.14	0.66
1:B:150:TYR:O	1:B:154:ASN:HB2	1.95	0.66
1:A:93:LYS:HE2	4:A:825:HOH:O	1.98	0.64
1:A:172:ASP:OD1	1:B:2:THR:HG23	1.98	0.64
1:A:172:ASP:OD1	1:B:2:THR:CG2	2.47	0.63
1:B:250:ARG:HE	1:B:271:HIS:CD2	2.16	0.62
1:B:150:TYR:CB	4:B:1008:HOH:O	2.47	0.60
1:A:34:HIS:HD2	4:A:787:HOH:O	1.84	0.60
1:A:311:ASP:O	1:A:312:ASP:CB	2.48	0.60
1:A:216:VAL:HG23	4:A:707:HOH:O	2.03	0.59
1:A:222:MET:HE3	2:C:59:VAL:HG23	1.82	0.58
1:B:110:ARG:HD2	1:B:122:PHE:CZ	2.36	0.58
1:A:115:LEU:O	1:B:256:HIS:CE1	2.54	0.58
1:B:76:GLU:HB3	1:B:78:THR:HG22	1.86	0.56
1:B:2:THR:HB	4:B:957:HOH:O	2.06	0.56
1:B:76:GLU:HG3	1:B:86:LEU:HD13	1.86	0.56
1:B:254:LEU:HD22	1:B:270[B]:LEU:HD11	1.87	0.56
1:B:198:GLU:CG	4:B:1007:HOH:O	2.43	0.56
1:B:76:GLU:HG3	1:B:86:LEU:CD1	2.35	0.55
1:B:213:GLY:HA2	1:B:320:GLN:O	2.07	0.55
2:C:28:LEU:HD22	2:C:68:VAL:HG11	1.89	0.54
1:B:149:LEU:O	1:B:154:ASN:ND2	2.40	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:110:ARG:CD	1:B:122:PHE:CZ	2.87	0.53
2:C:3:VAL:HG11	2:C:78:ILE:CG2	2.39	0.52
1:B:2:THR:CB	4:B:957:HOH:O	2.57	0.52
1:B:157:ASN:C	1:B:157:ASN:HD22	2.12	0.52
2:C:9[B]:THR:HG21	4:C:166:HOH:O	2.11	0.50
4:A:421:HOH:O	2:C:9[B]:THR:HG23	2.10	0.50
1:A:309:SER:O	1:A:311:ASP:O	2.30	0.50
1:B:76:GLU:HG2	4:B:842:HOH:O	2.13	0.49
1:B:227:VAL:HB	1:B:230:LEU:HG	1.95	0.49
1:A:-1:ARG:N	4:A:625:HOH:O	2.44	0.49
1:B:110:ARG:HD3	1:B:122:PHE:HZ	1.73	0.48
1:A:312:ASP:CB	4:A:602:HOH:O	2.60	0.48
1:B:110:ARG:HD3	1:B:122:PHE:CZ	2.49	0.47
1:B:226:PHE:CD1	4:B:997:HOH:O	2.68	0.47
1:B:134:ALA:O	1:B:138:GLU:HG2	2.15	0.47
1:B:21:ARG:HD2	4:B:874:HOH:O	2.15	0.46
1:B:103:GLU:HA	1:B:122:PHE:CD1	2.50	0.46
1:B:21:ARG:HB2	1:B:257[A]:THR:HG22	1.98	0.46
1:B:1:MET:HE3	4:B:925:HOH:O	2.17	0.45
1:A:93:LYS:HG3	4:A:825:HOH:O	2.15	0.45
2:C:2:ASN:N	4:C:202:HOH:O	2.49	0.44
1:B:244:ALA:O	1:B:248:VAL:HG23	2.18	0.44
1:A:172:ASP:OD1	1:B:2:THR:HG21	2.18	0.43
2:C:12:ARG:N	2:C:13:PRO:CD	2.81	0.43
1:B:215:GLY:N	4:B:1003:HOH:O	2.34	0.43
1:A:245:VAL:HG12	4:A:445:HOH:O	2.18	0.42
1:B:92:LEU:HD13	4:B:941:HOH:O	2.18	0.42
1:B:157:ASN:C	1:B:157:ASN:ND2	2.73	0.42
1:B:254:LEU:HD22	1:B:270[B]:LEU:HD21	2.02	0.42
1:A:172:ASP:O	1:B:3:ARG:HD3	2.20	0.42
2:C:3:VAL:HG11	2:C:78:ILE:HG21	2.03	0.41
1:A:128:GLY:HA2	4:A:434:HOH:O	2.20	0.41
1:B:138:GLU:HG2	1:B:138:GLU:H	1.61	0.41
1:B:256:HIS:HD2	4:B:821:HOH:O	2.03	0.41
1:B:99[B]:CYS:SG	1:B:101:MET:HE1	2.48	0.40
1:B:218:ALA:O	1:B:222:MET:HG3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	324/325 (100%)	315 (97%)	9 (3%)	0	100	100
1	B	321/325 (99%)	311 (97%)	7 (2%)	3 (1%)	14	3
2	C	91/93 (98%)	89 (98%)	2 (2%)	0	100	100
All	All	736/743 (99%)	715 (97%)	18 (2%)	3 (0%)	30	13

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	131	THR
1	B	150	TYR
1	B	141	ALA

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/245 (98%)	240 (100%)	1 (0%)	89	81
1	B	229/245 (94%)	221 (96%)	8 (4%)	31	6
2	C	69/74 (93%)	67 (97%)	2 (3%)	37	10
All	All	539/564 (96%)	528 (98%)	11 (2%)	52	20

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	ARG
1	B	67	LEU
1	B	137	LYS
1	B	138	GLU
1	B	148	MET
1	B	157	ASN
1	B	270[A]	LEU
1	B	270[B]	LEU
1	B	311	ASP
2	C	19	LYS
2	C	74	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	20	GLN
1	A	34	HIS
1	A	202	ASN
1	A	256	HIS
1	B	13	ASN
1	B	130	ASN
1	B	151	GLN
1	B	157	ASN
1	B	199	HIS
1	B	256	HIS
1	B	271	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PLP	A	401	1	15,15,16	1.09	2 (13%)	21,22,23	1.02	2 (9%)
3	PLP	B	401	1	15,15,16	1.11	2 (13%)	21,22,23	1.02	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PLP	A	401	1	-	0/6/6/8	0/1/1/1
3	PLP	B	401	1	-	0/6/6/8	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401	PLP	C2-N1	2.63	1.38	1.33
3	A	401	PLP	C2-N1	2.59	1.38	1.33
3	A	401	PLP	C6-N1	2.04	1.38	1.34
3	B	401	PLP	C6-N1	2.02	1.38	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401	PLP	C5-C6-N1	-2.30	120.09	123.83
3	A	401	PLP	C5-C6-N1	-2.27	120.14	123.83
3	B	401	PLP	C6-C5-C4	2.14	119.85	118.10
3	A	401	PLP	C6-C5-C4	2.09	119.81	118.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	325/325 (100%)	0.10	12 (3%) 45 54	10, 19, 35, 49	1 (0%)
1	B	321/325 (98%)	0.62	40 (12%) 9 11	9, 23, 50, 60	4 (1%)
2	C	92/93 (98%)	0.44	2 (2%) 62 70	14, 24, 35, 41	1 (1%)
All	All	738/743 (99%)	0.37	54 (7%) 22 27	9, 21, 41, 60	6 (0%)

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	141	ALA	5.9
1	B	133	VAL	5.0
1	B	145	SER	5.0
1	B	150	TYR	4.9
1	B	140	ALA	4.4
2	C	50	SER	4.3
1	B	78	THR	3.9
1	A	312	ASP	3.7
1	B	309	SER	3.7
1	B	146	TRP	3.6
1	A	29	GLY	3.4
1	B	319	GLY	3.4
1	B	136	ALA	3.3
1	B	147	VAL	3.3
1	A	64	ALA	3.2
1	B	120	ILE	3.2
1	B	310	LEU	3.2
1	A	283	ALA	3.1
1	B	134	ALA	3.1
1	B	152	TYR	3.1
1	B	131	THR	3.0
1	B	219	LEU	3.0
1	A	313	ALA	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	311	ASP	2.9
1	B	144	PRO	2.8
1	A	0	HIS	2.8
1	B	143	ASN	2.7
1	B	132	ALA	2.7
1	B	308	GLY	2.7
1	B	125	ALA	2.7
2	C	2	ASN	2.6
1	B	121	ILE	2.5
1	B	139	LEU	2.5
1	B	-1	ARG	2.5
1	B	98	ILE	2.5
1	B	64	ALA	2.4
1	A	1	MET	2.4
1	B	75	LEU	2.4
1	B	142	THR	2.4
1	B	122	PHE	2.4
1	A	221	ASN	2.3
1	B	100	VAL	2.3
1	B	0	HIS	2.3
1	B	216	VAL	2.3
1	A	315	THR	2.2
1	B	90	ALA	2.2
1	B	106	SER	2.2
1	B	227	VAL	2.2
1	B	307	ALA	2.1
1	B	117	GLY	2.1
1	B	137	LYS	2.1
1	B	66	GLY	2.1
1	A	70	PRO	2.1
1	A	27	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	PLP	A	401	15/16	0.97	0.07	13,14,20,24	0
3	PLP	B	401	15/16	0.97	0.06	15,16,18,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.