



wwPDB EM Validation Summary Report ⓘ

Dec 30, 2024 – 04:54 PM EST

PDB ID : 8DWO
EMDB ID : EMD-27757
Title : Cryo-EM Structure of Eastern Equine Encephalitis Virus in complex with SKE26 Fab
Authors : Pletnev, S.; Verardi, R.; Roedegeer, M.; Kwong, P.
Deposited on : 2022-08-01
Resolution : 3.50 Å(reported)
Based on initial model : 6XO4

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev113
MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.40


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY









The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	441	
1	F	441	
1	J	441	
2	B	420	
2	G	420	
2	K	420	
3	H	233	
3	S	233	

Continued on next page...

Mol	Chain	Length	Quality of chain
3	U	233	
4	L	219	
4	T	219	
4	V	219	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 22572 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Envelope glycoprotein E1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	392	Total 3000	C 1903	N 498	O 579	S 20	0	0
1	F	392	Total 3000	C 1903	N 498	O 579	S 20	0	0
1	J	392	Total 3000	C 1903	N 498	O 579	S 20	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	89	TYR	TRP	conflict	UNP Q88678
F	89	TYR	TRP	conflict	UNP Q88678
J	89	TYR	TRP	conflict	UNP Q88678

- Molecule 2 is a protein called Envelope glycoprotein E2.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	346	Total 2720	C 1706	N 503	O 495	S 16	0	0
2	G	346	Total 2720	C 1706	N 503	O 495	S 16	0	0
2	K	346	Total 2720	C 1706	N 503	O 495	S 16	0	0

- Molecule 3 is a protein called SKE26 Fab Heavy Chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	H	122	Total 972	C 624	N 164	O 181	S 3	0	0
3	S	122	Total 972	C 624	N 164	O 181	S 3	0	0
3	U	122	Total 972	C 624	N 164	O 181	S 3	0	0


- Molecule 4 is a protein called SKE26 Fab Light Chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	L	112	Total	C	N	O	S	0	0
			832	521	139	168	4		
4	T	112	Total	C	N	O	S	0	0
			832	521	139	168	4		
4	V	112	Total	C	N	O	S	0	0
			832	521	139	168	4		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Envelope glycoprotein E1




• Molecule 1: Envelope glycoprotein E1




• Molecule 1: Envelope glycoprotein E1




- Molecule 2: Envelope glycoprotein E2




- Molecule 2: Envelope glycoprotein E2




- Molecule 2: Envelope glycoprotein E2




- Molecule 3: SKE26 Fab Heavy Chain



- Molecule 3: SKE26 Fab Heavy Chain



- Molecule 3: SKE26 Fab Heavy Chain



- Molecule 4: SKE26 Fab Light Chain

[illegible]

Position	Residue	Score
D1	SER	0.98
E17	VAL	0.96
S26	CYS	0.95
Q27	LEU	0.94
L27C	ASN	0.93
T31	PHE	0.92
Y32	PRO	0.91
L33	ARG	0.90
H34	GLU	0.89
K39	ALA	0.88
L46	SER	0.87
L47	LYS	0.86
I48	TRP	0.85
Y49	LYS	0.84
N53	ASP	0.83
R61	GLY	0.82
D70	ALA	0.81
A80	LEU	0.80
G81	LYS	0.79
Q90	THR	0.78
V91	GLY	0.77
I92	THR	0.76
L96	GLN	0.75
K107	ASP	0.74
THR	SER	0.73
VAL	LYS	0.72
ALA	ASN	0.71
ALA	ASP	0.70
PRO	THR	0.69
SER	TTR	0.68
VAL	SER	0.67
PHE	LEU	0.66
ILE	SER	0.65
PHE	THR	0.64
PRO	LEU	0.63
PRO	THR	0.62
SER	LEU	0.61
GLU	SER	0.60
ASP	THR	0.59
GLN	THR	0.58
VAL	GLU	0.57
LYS	TTR	0.56
THR	GLN	0.55
GLY	THR	0.54
ASN	LYS	0.53
SER	ASN	0.52
GLN	SER	0.51
GLU	GLN	0.50
VAL	GLY	0.49
THR	GLU	0.48
THR	GLY	0.47
LYS	ALA	0.46
LEU	ALA	0.45
LYS	LEU	0.44
THR	LYS	0.43
SER	THR	0.42
PHE	SER	0.41
ASN	PHE	0.40
ARG	ASN	0.39
GLY	ARG	0.38
GLY	GLY	0.37
GLU	GLU	0.36
CYS	CYS	0.35
VAL	VAL	0.34
THR	THR	0.33
HIS	THR	0.32
GLN	HIS	0.31
ASN	GLN	0.30
PHE	ASN	0.29
THR	PHE	0.28
PRO	THR	0.27
PRO	PRO	0.26
SER	PRO	0.25
THR	SER	0.24
LYS	THR	0.23
LYS	LYS	0.22
SER	LYS	0.21
PHE	SER	0.20
THR	PHE	0.19
THR	THR	0.18
THR	THR	0.17
THR	THR	0.16
THR	THR	0.15
THR	THR	0.14
THR	THR	0.13
THR	THR	0.12
THR	THR	0.11
THR	THR	0.10
THR	THR	0.09
THR	THR	0.08
THR	THR	0.07
THR	THR	0.06
THR	THR	0.05
THR	THR	0.04
THR	THR	0.03
THR	THR	0.02
THR	THR	0.01

SER	GLY	D1
HIS	THR	
LYS	VAL	Q6
VAL	SER	
TYR	VAL	E17
ALA	VAL	
CYS	CYS	T21
GLU	LEU	
VAL	LEU	S26
THR	ASN	Q27
HIS	ASN	S27A
GLN	PHE	L27B
GLY	TYR	
LEU	PRO	T31
SER	ARG	Y32
SER	GLU	L33
PRO	ALA	
VAL	SER	Q38
THR	VAL	
LYS	LYS	G50
SER	TRP	
PHE	LYS	S65
ASN	VAL	G66
ARG	ASP	
GLY	GLY	F71
GLU	ALA	I72
CYS	LEU	L73
	LYS	I74
	THR	
	GLY	C88
	ASN	P89
	SER	Q90
	GLN	V91
	GLU	
	SER	L96
	VAL	
	THR	G101
	GLU	I102
	GLN	
	ASP	K107
	SER	
	LYS	THR
	ASP	VAL
	ASN	ALA
	THR	ALA
	TYR	PRO
	SER	SER
	LEU	VAL
	SER	PHE
	SER	ILE
	THR	PHE
	LEU	PRO
	THR	PRO
	LEU	SER
	SER	GLU
	SER	ASP
	THR	GLN
	GLU	VAL
	TYR	LYS
	GLN	SER

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	3071400	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	58.06	Depositor
Minimum defocus (nm)	700	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	5.102	Depositor
Minimum map value	-3.035	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.061	Depositor
Recommended contour level	0.4	Depositor
Map size (Å)	1162.0, 1162.0, 1162.0	wwPDB
Map dimensions	700, 700, 700	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.66, 1.66, 1.66	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/3080	0.58	1/4197 (0.0%)
1	F	0.29	0/3080	0.58	1/4197 (0.0%)
1	J	0.29	0/3080	0.58	2/4197 (0.0%)
2	B	0.26	0/2799	0.60	0/3808
2	G	0.26	0/2799	0.60	0/3808
2	K	0.27	0/2799	0.60	0/3808
3	H	0.27	0/1001	0.61	1/1363 (0.1%)
3	S	0.27	0/1001	0.61	1/1363 (0.1%)
3	U	0.28	0/1001	0.58	0/1363
4	L	0.29	0/850	0.59	0/1159
4	T	0.30	0/850	0.62	1/1159 (0.1%)
4	V	0.29	0/850	0.60	0/1159
All	All	0.28	0/23190	0.59	7/31581 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	J	146	ASP	CB-CG-OD1	7.29	124.86	118.30
1	F	117	ASP	CB-CG-OD2	6.44	124.10	118.30
1	J	154	PRO	CA-N-CD	-5.78	103.40	111.50
1	A	117	ASP	CB-CG-OD2	5.71	123.44	118.30
3	H	31	ASP	CB-CG-OD1	5.68	123.41	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	62	CYS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3000	0	2900	34	0
1	F	3000	0	2900	42	0
1	J	3000	0	2902	29	0
2	B	2720	0	2655	27	0
2	G	2720	0	2655	45	0
2	K	2720	0	2655	44	0
3	H	972	0	936	14	0
3	S	972	0	936	14	0
3	U	972	0	936	18	0
4	L	832	0	811	11	0
4	T	832	0	811	11	0
4	V	832	0	811	14	0
All	All	22572	0	21908	286	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 286 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:J:135:ILE:HD12	1:J:157:ILE:HD13	1.59	0.84
3:U:51:ILE:HD13	3:U:57:ILE:HD13	1.65	0.78
3:S:56:ASP:OD2	3:S:58:ARG:NH2	2.20	0.75
1:J:92:ALA:O	2:K:224:ARG:NH1	2.18	0.74
1:J:75:ASP:OD2	1:J:218:THR:OG1	2.05	0.74

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	388/441 (88%)	359 (92%)	29 (8%)	0	100	100
1	F	388/441 (88%)	359 (92%)	29 (8%)	0	100	100
1	J	388/441 (88%)	355 (92%)	33 (8%)	0	100	100
2	B	344/420 (82%)	308 (90%)	36 (10%)	0	100	100
2	G	344/420 (82%)	310 (90%)	34 (10%)	0	100	100
2	K	344/420 (82%)	306 (89%)	38 (11%)	0	100	100
3	H	120/233 (52%)	116 (97%)	4 (3%)	0	100	100
3	S	120/233 (52%)	117 (98%)	3 (2%)	0	100	100
3	U	120/233 (52%)	118 (98%)	2 (2%)	0	100	100
4	L	110/219 (50%)	107 (97%)	3 (3%)	0	100	100
4	T	110/219 (50%)	106 (96%)	4 (4%)	0	100	100
4	V	110/219 (50%)	107 (97%)	3 (3%)	0	100	100
All	All	2886/3939 (73%)	2668 (92%)	218 (8%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/371 (90%)	326 (98%)	7 (2%)	48	71
1	F	333/371 (90%)	322 (97%)	11 (3%)	33	61

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	J	333/371 (90%)	319 (96%)	14 (4%)	25	54
2	B	300/367 (82%)	286 (95%)	14 (5%)	22	51
2	G	300/367 (82%)	291 (97%)	9 (3%)	36	63
2	K	300/367 (82%)	285 (95%)	15 (5%)	20	49
3	H	106/204 (52%)	104 (98%)	2 (2%)	52	73
3	S	106/204 (52%)	103 (97%)	3 (3%)	38	65
3	U	106/204 (52%)	102 (96%)	4 (4%)	28	57
4	L	95/192 (50%)	91 (96%)	4 (4%)	25	54
4	T	95/192 (50%)	91 (96%)	4 (4%)	25	54
4	V	95/192 (50%)	93 (98%)	2 (2%)	48	71
All	All	2502/3402 (74%)	2413 (96%)	89 (4%)	32	59

5 of 89 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	K	9	TYR
2	K	334	ARG
2	K	26	ARG
2	K	205	ARG
4	L	70	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	5	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-27757. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 350




Y Index: 350




Z Index: 350


6.2.2 Raw map



X Index: 350



Y Index: 350




Z Index: 350


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 336




Y Index: 489




Z Index: 473


6.3.2 Raw map



X Index: 336



Y Index: 472




Z Index: 473


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.4. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_27757_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 818 nm³; this corresponds to an approximate mass of 739 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.286 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.286 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.50	-	-
Author-provided FSC curve	3.36	3.65	3.37
Unmasked-calculated*	4.97	8.06	5.34

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.97 differs from the reported value 3.5 by more than 10 %

9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-27757 and PDB model 8DWO. Per-residue inclusion information can be found in [section 3](#) on [page 6](#).

9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.4 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.4).

9.4 Atom inclusion [i](#)



At the recommended contour level, 93% of all backbone atoms, 88% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.4) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	<div><div></div></div> 0.8850	<div><div></div></div> 0.4830
A	<div><div></div></div> 0.8920	<div><div></div></div> 0.4950
B	<div><div></div></div> 0.8990	<div><div></div></div> 0.4990
F	<div><div></div></div> 0.8930	<div><div></div></div> 0.4980
G	<div><div></div></div> 0.9000	<div><div></div></div> 0.4990
H	<div><div></div></div> 0.8620	<div><div></div></div> 0.4320
J	<div><div></div></div> 0.8630	<div><div></div></div> 0.4740
K	<div><div></div></div> 0.9000	<div><div></div></div> 0.4970
L	<div><div></div></div> 0.8500	<div><div></div></div> 0.4540
S	<div><div></div></div> 0.8910	<div><div></div></div> 0.4500
T	<div><div></div></div> 0.8400	<div><div></div></div> 0.4610
U	<div><div></div></div> 0.8810	<div><div></div></div> 0.4480
V	<div><div></div></div> 0.8540	<div><div></div></div> 0.4480

1.0

0.0

<0.0