



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 23, 2024 – 10:40 AM EDT

PDB ID : 5DXV  
Title : Crystal structure of Rethreaded DHFR  
Authors : Faham, S.  
Deposited on : 2015-09-24  
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1483 (1.56-1.56)
Clashscore	141614	1529 (1.56-1.56)
Ramachandran outliers	138981	1498 (1.56-1.56)
Sidechain outliers	138945	1495 (1.56-1.56)
RSRZ outliers	127900	1465 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	156	<div> <div>2%</div> <div> <div></div> <div>87%</div> <div>8%</div> <div>••</div> </div> </div>
1	B	156	<div> <div>7%</div> <div> <div></div> <div>83%</div> <div>7%</div> <div>• 7%</div> </div> </div>

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 2543 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called rethreaded DHFR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	149	Total	C	N	O	S	0	0	0
			1166	743	198	220	5			
1	B	145	Total	C	N	O	S	0	1	0
			1152	735	199	213	5			

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula:  $C_{21}H_{28}N_7O_{17}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula:  $C_4H_{10}O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total 1	Cl 1	0	0
5	B	1	Total 1	Cl 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	87	Total 87	O 87	0	0
6	B	66	Total 66	O 66	0	0



- Molecule 1: rethreaded DHFR.



MET	A1	L8	R12	V20	D21	E22	E23	A39	Q40	W47	R61	R69	P83	G84	T88	D133	H137	C140	F141	I142	I143	L144	F145	R146	E149	H1S	H1S	H1S	H1S	H1S
-----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----

## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	44.82Å 120.05Å 111.19Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 1.55 31.54 – 1.55	Depositor EDS
% Data completeness (in resolution range)	91.4 (30.00-1.55) 91.4 (31.54-1.55)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.55 (at 1.55Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.185 , 0.214 0.194 , 0.224	Depositor DCC
$R_{free}$ test set	2025 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.6	Xtriage
Anisotropy	0.775	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 43.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2543	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.06% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, EDO, NAP, CME, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.23	3/1186 (0.3%)	1.22	8/1616 (0.5%)
1	B	1.10	1/1174 (0.1%)	1.09	4/1595 (0.3%)
All	All	1.17	4/2360 (0.2%)	1.16	12/3211 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	32	SER	CA-CB	5.65	1.61	1.52
1	B	47	TRP	CZ3-CH2	5.10	1.48	1.40
1	A	54	ASP	CB-CG	5.06	1.62	1.51
1	A	91	TRP	CE3-CZ3	-5.01	1.29	1.38

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	133	ASP	CB-CG-OD1	7.91	125.42	118.30
1	A	146	ARG	NE-CZ-NH1	7.78	124.19	120.30
1	B	61	ARG	NE-CZ-NH2	-7.56	116.52	120.30
1	A	146	ARG	NE-CZ-NH2	-7.33	116.63	120.30
1	A	74	ARG	NE-CZ-NH1	7.26	123.93	120.30
1	A	16	ASP	CB-CG-OD1	-7.20	111.82	118.30
1	B	61	ARG	NE-CZ-NH1	6.23	123.41	120.30

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	146	ARG	NE-CZ-NH1	5.91	123.25	120.30
1	A	74	ARG	NE-CZ-NH2	-5.77	117.41	120.30
1	A	133	ASP	CB-CG-OD1	5.28	123.05	118.30
1	A	16	ASP	CB-CG-OD2	5.17	122.96	118.30
1	A	133	ASP	CB-CG-OD2	-5.11	113.70	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	84	GLY	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1166	0	1112	4	0
1	B	1152	0	1112	8	0
2	A	48	0	25	1	0
3	A	7	0	10	0	0
3	B	7	0	10	0	0
4	A	4	0	6	0	1
4	B	4	0	6	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	87	0	0	0	1
6	B	66	0	0	0	1
All	All	2543	0	2281	12	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:61:ARG:HH22	1:B:84:GLY:HA2	1.11	1.08

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:GLN:HE22	2:A:201:NAP:H61A	1.27	0.81
1:B:61:ARG:NH2	1:B:84:GLY:HA2	1.95	0.75
1:A:140:CME:SD	1:A:142:GLU:HB2	2.39	0.62
1:B:140:CME:SD	1:B:142:GLU:HB2	2.45	0.56
1:B:39:ALA:HA	1:B:137:HIS:CD2	2.44	0.52
1:A:15:GLY:H	1:A:18:HIS:CD2	2.29	0.51
1:B:22:TYR:O	1:B:24:PRO:HD3	2.16	0.46
1:B:8:LEU:HD11	1:B:19:PHE:CG	2.51	0.45
1:A:23:GLU:OE2	1:A:24:PRO:HD2	2.19	0.42
1:B:29:SER:HA	1:B:144:LEU:HD23	2.03	0.41
1:B:61:ARG:HH22	1:B:84:GLY:CA	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:203:EDO:O1	4:A:203:EDO:O1[3_654]	1.92	0.28
6:A:366:HOH:O	6:B:401:HOH:O[5_545]	2.10	0.10

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	146/156 (94%)	144 (99%)	2 (1%)	0	100	100
1	B	141/156 (90%)	137 (97%)	4 (3%)	0	100	100
All	All	287/312 (92%)	281 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	120/132 (91%)	120 (100%)	0	100	100
1	B	119/132 (90%)	118 (99%)	1 (1%)	81	66
All	All	239/264 (90%)	238 (100%)	1 (0%)	91	82

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	8	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	HIS
1	A	119	GLN
1	A	137	HIS
1	B	51	ASN
1	B	137	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CME	B	140	1	8,9,10	0.93	0	5,9,11	1.64	1 (20%)
1	CME	A	140	1	8,9,10	0.76	0	5,9,11	2.17	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CME	B	140	1	-	0/5/8/10	-
1	CME	A	140	1	-	0/5/8/10	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	140	CME	CB-SG-SD	3.92	113.97	103.82
1	B	140	CME	CB-SG-SD	3.17	112.04	103.82
1	A	140	CME	OH-CZ-CE	-2.73	100.05	110.83

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	140	CME	1	0
1	A	140	CME	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAP	A	201	-	45,52,52	1.21	3 (6%)	56,80,80	1.54	12 (21%)
3	PEG	A	202	-	6,6,6	0.38	0	5,5,5	0.40	0
4	EDO	A	203	-	3,3,3	0.95	0	2,2,2	0.67	0
4	EDO	B	302	-	3,3,3	0.52	0	2,2,2	0.04	0
3	PEG	B	301	-	6,6,6	0.40	0	5,5,5	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	201	-	-	3/31/67/67	0/5/5/5
3	PEG	A	202	-	-	3/4/4/4	-
4	EDO	A	203	-	-	0/1/1/1	-
4	EDO	B	302	-	-	1/1/1/1	-
3	PEG	B	301	-	-	3/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	201	NAP	P2B-O2B	5.45	1.69	1.59
2	A	201	NAP	C2A-N1A	-2.40	1.29	1.33
2	A	201	NAP	C5A-C4A	2.00	1.46	1.40

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	201	NAP	O2A-PA-O1A	3.60	130.02	112.24
2	A	201	NAP	C3N-C7N-N7N	3.45	121.89	117.75
2	A	201	NAP	C3D-C2D-C1D	3.08	105.61	100.98
2	A	201	NAP	O2B-P2B-O1X	-3.02	97.75	109.39
2	A	201	NAP	O7N-C7N-N7N	-2.66	118.79	122.58
2	A	201	NAP	C3N-C2N-N1N	-2.66	117.83	120.43
2	A	201	NAP	C5A-C6A-N1A	-2.64	114.38	120.35

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	201	NAP	N6A-C6A-N1A	2.60	123.97	118.57
2	A	201	NAP	C1B-N9A-C4A	-2.37	122.47	126.64
2	A	201	NAP	C4A-C5A-N7A	-2.20	107.11	109.40
2	A	201	NAP	C6N-N1N-C2N	2.18	123.96	121.97
2	A	201	NAP	O3X-P2B-O1X	2.05	118.70	110.68

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	201	NAP	C2B-O2B-P2B-O3X
2	A	201	NAP	O4D-C1D-N1N-C6N
3	B	301	PEG	C4-C3-O2-C2
3	B	301	PEG	O1-C1-C2-O2
4	B	302	EDO	O1-C1-C2-O2
3	A	202	PEG	O2-C3-C4-O4
2	A	201	NAP	C2B-O2B-P2B-O1X
3	A	202	PEG	C4-C3-O2-C2
3	A	202	PEG	C1-C2-O2-C3
3	B	301	PEG	O2-C3-C4-O4

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	201	NAP	1	0
4	A	203	EDO	0	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	148/156 (94%)	0.06	3 (2%) 65 71	18, 28, 48, 63	0
1	B	144/156 (92%)	0.25	11 (7%) 13 16	20, 30, 59, 68	0
All	All	292/312 (93%)	0.15	14 (4%) 30 35	18, 29, 57, 68	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	85	THR	6.2
1	B	17	THR	5.8
1	B	83	PRO	4.1
1	A	10	VAL	3.1
1	B	149	GLU	3.0
1	B	25	ASP	2.7
1	B	37	ALA	2.5
1	B	21	ASP	2.4
1	B	39	ALA	2.4
1	B	40	GLN	2.4
1	B	69	ARG	2.3
1	A	134	ALA	2.2
1	A	68	GLY	2.2
1	B	18	HIS	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
1	CME	A	140	10/11	0.92	0.09	23,26,47,50	0
1	CME	B	140	10/11	0.95	0.08	26,29,60,65	0

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PEG	A	202	7/7	0.80	0.12	46,55,60,63	1
3	PEG	B	301	7/7	0.81	0.13	46,49,56,59	1
4	EDO	B	302	4/4	0.86	0.19	24,39,42,46	1
4	EDO	A	203	4/4	0.92	0.13	33,43,44,44	1
2	NAP	A	201	48/48	0.96	0.08	21,28,36,38	0
5	CL	B	303	1/1	0.99	0.03	28,28,28,28	0
5	CL	A	204	1/1	1.00	0.06	34,34,34,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.5 Other polymers [i](#)

There are no such residues in this entry.