



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 27, 2024 – 02:57 AM EDT

PDB ID : 5E5R
Title : Crystal structure of the complex between Carbonic anhydrase-like domain of PTPRG and Immunoglobulin domains 2-3 of CNTN3
Authors : Nikolaienko, R.M.; Bouyain, S.
Deposited on : 2015-10-09
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3775 (2.60-2.60)
Clashscore	180529	4181 (2.60-2.60)
Ramachandran outliers	177936	4129 (2.60-2.60)
Sidechain outliers	177891	4129 (2.60-2.60)
RSRZ outliers	164620	3775 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	265	
1	C	265	
2	B	193	
2	D	193	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MLI	D	401	-	-	X	-
4	FMT	C	402	-	X	-	-
4	FMT	C	403	-	X	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7291 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Receptor-type tyrosine-protein phosphatase gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	258	Total	C	N	O	S	0	0	0
			2081	1328	360	388	5			
1	C	264	Total	C	N	O	S	0	0	0
			2138	1364	368	401	5			

- Molecule 2 is a protein called Contactin-3.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	187	Total	C	N	O	S	0	0	0
			1476	930	258	280	8			
2	D	192	Total	C	N	O	S	0	0	0
			1510	952	263	287	8			

- Molecule 3 is MALONATE ION (three-letter code: MLI) (formula: $C_3H_2O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	3	4		
3	C	1	Total	C	O	0	0
			7	3	4		
3	D	1	Total	C	O	0	0
			7	3	4		

- Molecule 4 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			3	1	2		
4	C	1	Total	C	O	0	0
			3	1	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total	O	0	0
			3	3		
5	B	3	Total	O	0	0
			3	3		
5	C	24	Total	O	0	0
			24	24		
5	D	29	Total	O	0	0
			29	29		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Receptor-type tyrosine-protein phosphatase gamma




- Molecule 1: Receptor-type tyrosine-protein phosphatase gamma




- Molecule 2: Contactin-3



- Molecule 2: Contactin-3





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	74.14Å 90.53Å 147.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.15 – 2.60 49.15 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.2 (49.15-2.60) 95.1 (49.15-2.60)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 2.61Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.187 , 0.249 0.194 , 0.254	Depositor DCC
R_{free} test set	1557 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	54.3	Xtriage
Anisotropy	0.192	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 49.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7291	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMT, MLI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/2143	0.49	0/2906
1	C	0.45	0/2203	0.56	0/2991
2	B	0.44	0/1507	0.57	0/2041
2	D	0.50	0/1544	0.61	0/2093
All	All	0.44	0/7397	0.55	0/10031

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2081	0	1966	20	1
1	C	2138	0	2021	24	1
2	B	1476	0	1450	17	0
2	D	1510	0	1480	25	0
3	A	7	0	2	0	0
3	C	7	0	2	0	0
3	D	7	0	2	3	0
4	C	6	0	4	0	0
5	A	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	3	0	0	1	0
5	C	24	0	0	0	0
5	D	29	0	0	2	0
All	All	7291	0	6927	80	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94:ARG:HB2	1:A:319:LYS:HB3	1.68	0.74
1:C:184:PHE:CZ	1:C:195:ARG:HG3	2.31	0.66
1:C:85:PRO:HG3	1:C:162:GLU:HB3	1.80	0.61
1:A:183:ASP:HB3	1:C:272:ARG:HH12	1.66	0.61
2:B:203:THR:OG1	2:B:204:ASN:N	2.28	0.61
2:B:227:PRO:HG2	2:B:300:GLU:HA	1.82	0.61
2:D:296:GLU:HG2	2:D:309:ARG:HG2	1.84	0.59
1:C:143:PHE:HB3	1:C:178:PHE:HB3	1.84	0.59
2:B:158:VAL:HG23	2:B:195:THR:HG23	1.84	0.59
1:A:183:ASP:CB	1:C:272:ARG:HH12	2.15	0.59
1:C:226:HIS:HB3	2:D:228:LYS:HD2	1.84	0.59
2:D:270:PRO:HD2	2:D:273:ILE:HD12	1.87	0.57
1:C:226:HIS:CD2	2:D:228:LYS:HB2	2.40	0.56
1:A:82:HIS:HA	1:A:315:ARG:CZ	2.36	0.55
2:D:137:GLY:HA2	2:D:185:LYS:HG3	1.86	0.55
1:C:181:PRO:C	1:C:183:ASP:H	2.10	0.55
1:C:213:ALA:HB1	1:C:240:ASP:HB3	1.88	0.55
1:A:85:PRO:HG3	1:A:162:GLU:HB3	1.89	0.54
2:D:197:VAL:HG22	2:D:208:LEU:HD22	1.90	0.54
1:C:296:VAL:HG22	2:D:142:LEU:HD23	1.90	0.54
1:C:148:VAL:HG22	1:C:176:ILE:HG12	1.90	0.53
2:D:248:GLU:HG3	3:D:401:MLI:H12	1.90	0.53
2:B:197:VAL:HB	2:B:208:LEU:HD22	1.90	0.53
2:B:272:LYS:NZ	2:B:286:ASN:O	2.40	0.53
1:A:276:ILE:HD13	1:A:281:LEU:HG	1.91	0.52
2:D:227:PRO:HG2	2:D:300:GLU:HA	1.93	0.51
1:C:105:ASP:HB3	1:C:133:PHE:HB2	1.93	0.51
1:A:196:ILE:HG13	1:C:208:PRO:HD3	1.92	0.51
2:D:246:ARG:O	3:D:401:MLI:H11	2.12	0.50
2:D:125:LYS:HD3	2:D:207:VAL:HG13	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:246:ARG:HB3	2:D:281:MET:SD	2.52	0.49
2:B:300:GLU:HB3	2:B:305:LYS:HB3	1.94	0.49
2:B:171:ARG:NH1	2:B:190:ASP:OD2	2.44	0.48
2:D:129:ARG:NH2	5:D:503:HOH:O	2.46	0.48
2:D:235:GLU:HG3	2:D:236:THR:HG23	1.93	0.48
2:B:188:PRO:HA	2:B:216:LEU:HD11	1.95	0.47
1:C:145:ALA:HB2	1:C:178:PHE:CE2	2.50	0.47
2:D:214:LEU:HD12	2:D:214:LEU:HA	1.64	0.47
2:B:245:VAL:HG22	2:B:284:ILE:HB	1.96	0.47
1:C:127:LEU:HG	1:C:129:LYS:HE2	1.97	0.47
2:D:273:ILE:HG12	2:D:284:ILE:HG12	1.97	0.47
1:A:89:LEU:HD13	1:A:92:TYR:HB2	1.97	0.46
1:A:143:PHE:HB3	1:A:178:PHE:HB3	1.97	0.46
2:B:263:ARG:HG3	2:B:269:PHE:CE1	2.51	0.46
2:D:142:LEU:HD12	2:D:181:LEU:HD23	1.97	0.46
2:D:158:VAL:HG22	2:D:195:THR:HB	1.98	0.45
1:A:243:PRO:HG2	1:A:270:PHE:HB3	1.97	0.45
1:A:227:HIS:O	1:A:228:GLU:HB2	2.17	0.45
1:C:159:ALA:HB1	1:C:169:ARG:HD3	1.99	0.45
2:D:262:ARG:HD2	5:D:528:HOH:O	2.16	0.45
1:A:105:ASP:HB2	1:A:133:PHE:HB2	1.99	0.44
1:A:184:PHE:CZ	1:A:195:ARG:HG3	2.51	0.44
1:C:86:ILE:HG12	1:C:252:TYR:CZ	2.52	0.44
2:D:271:ASN:N	2:D:271:ASN:OD1	2.49	0.44
1:C:175:GLN:HA	1:C:200:MET:O	2.18	0.44
1:C:289:THR:OG1	1:C:290:THR:N	2.49	0.44
2:B:168:GLU:OE1	2:B:277:LYS:HE3	2.18	0.43
1:C:84:SER:O	1:C:307:ARG:NH1	2.51	0.43
1:A:133:PHE:CE1	1:A:144:LYS:HD2	2.54	0.43
2:D:246:ARG:HD3	2:D:281:MET:SD	2.58	0.43
1:A:166:ASN:O	1:A:168:ARG:N	2.37	0.43
1:A:172:VAL:HB	1:A:204:PHE:HB2	2.01	0.42
2:B:239:ALA:HB3	2:B:287:PHE:CE2	2.54	0.42
1:A:215:ASP:HB2	1:A:216:PRO:HD3	2.00	0.42
2:D:262:ARG:HE	2:D:298:ILE:HD11	1.85	0.42
1:A:82:HIS:HA	1:A:315:ARG:NE	2.35	0.42
2:B:158:VAL:HG11	2:B:163:PRO:HA	2.02	0.42
1:C:161:SER:O	1:C:169:ARG:NH1	2.53	0.42
2:D:233:PHE:HA	3:D:401:MLI:O7	2.20	0.42
2:B:256:VAL:HA	2:B:257:PRO:HD3	1.86	0.42
2:D:225:TYR:CE2	2:D:253:GLY:HA2	2.55	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:226:GLU:HG3	5:B:403:HOH:O	2.20	0.41
1:C:94:ARG:HB3	1:C:318:SER:HA	2.03	0.41
2:B:278:PHE:O	2:B:279:ASN:HB2	2.21	0.41
1:C:272:ARG:HA	1:C:272:ARG:HD2	1.88	0.41
1:A:154:HIS:H	1:A:158:SER:HB3	1.86	0.40
1:C:90:ASP:OD2	1:C:250:TYR:OH	2.39	0.40
2:B:135:ARG:O	2:B:138:GLN:HB3	2.21	0.40
1:A:204:PHE:CD1	1:A:274:VAL:HB	2.55	0.40
2:D:158:VAL:CG2	2:D:195:THR:HB	2.52	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:278:TYR:OH	1:C:100:GLN:OE1[1_455]	2.14	0.06

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/265 (95%)	240 (95%)	12 (5%)	0	100	100
1	C	262/265 (99%)	244 (93%)	18 (7%)	0	100	100
2	B	181/193 (94%)	172 (95%)	9 (5%)	0	100	100
2	D	188/193 (97%)	178 (95%)	10 (5%)	0	100	100
All	All	883/916 (96%)	834 (94%)	49 (6%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/232 (97%)	220 (98%)	5 (2%)	47	72
1	C	232/232 (100%)	229 (99%)	3 (1%)	65	84
2	B	164/169 (97%)	157 (96%)	7 (4%)	25	49
2	D	168/169 (99%)	161 (96%)	7 (4%)	25	50
All	All	789/802 (98%)	767 (97%)	22 (3%)	38	65

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	89	LEU
1	A	127	LEU
1	A	212	SER
1	A	276	ILE
1	A	281	LEU
2	B	131	THR
2	B	153	LEU
2	B	195	THR
2	B	198	VAL
2	B	203	THR
2	B	245	VAL
2	B	274	LYS
1	C	217	ILE
1	C	258	THR
1	C	271	ARG
2	D	129	ARG
2	D	138	GLN
2	D	204	ASN
2	D	210	SER
2	D	231	VAL
2	D	245	VAL
2	D	271	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	309	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	FMT	C	402	-	2,2,2	2.66	2 (100%)	1,1,1	1.33	0
3	MLI	C	401	-	6,6,6	1.09	0	7,7,7	1.89	3 (42%)
4	FMT	C	403	-	2,2,2	2.69	2 (100%)	1,1,1	1.29	0
3	MLI	D	401	-	6,6,6	1.27	0	7,7,7	1.56	1 (14%)
3	MLI	A	401	-	6,6,6	1.24	0	7,7,7	1.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MLI	D	401	-	-	0/4/4/4	-
3	MLI	A	401	-	-	2/4/4/4	-
3	MLI	C	401	-	-	2/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	402	FMT	O1-C	3.16	1.39	1.22
4	C	403	FMT	O1-C	3.09	1.39	1.22
4	C	403	FMT	O2-C	2.22	1.40	1.28
4	C	402	FMT	O2-C	2.05	1.39	1.28

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	401	MLI	C3-C1-C2	2.80	122.83	112.95
3	C	401	MLI	O9-C3-C1	2.53	122.33	114.51
3	C	401	MLI	O7-C2-C1	2.33	121.72	114.51
3	C	401	MLI	C3-C1-C2	2.08	120.28	112.95

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	401	MLI	C3-C1-C2-O6
3	C	401	MLI	C3-C1-C2-O7
3	A	401	MLI	C3-C1-C2-O7
3	A	401	MLI	C3-C1-C2-O6

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	401	MLI	3	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	258/265 (97%)	-0.11	0 100 100	39, 79, 125, 180	0
1	C	264/265 (99%)	-0.44	0 100 100	32, 63, 107, 145	0
2	B	187/193 (96%)	-0.22	4 (2%) 63 58	42, 72, 112, 130	0
2	D	192/193 (99%)	-0.55	0 100 100	32, 54, 89, 114	0
All	All	901/916 (98%)	-0.32	4 (0%) 89 86	32, 67, 114, 180	0

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	147	PRO	2.6
2	B	152	GLU	2.4
2	B	204	ASN	2.2
2	B	126	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	MLI	D	401	7/7	0.68	0.20	121,123,130,133	0
3	MLI	C	401	7/7	0.86	0.15	80,83,99,117	0
4	FMT	C	402	3/3	0.86	0.16	73,73,76,76	0
4	FMT	C	403	3/3	0.88	0.16	48,48,55,62	0
3	MLI	A	401	7/7	0.89	0.12	93,99,108,114	0

6.5 Other polymers [i](#)

There are no such residues in this entry.