



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 28, 2025 – 03:12 PM EDT

PDB ID : 3ECV / pdb_00003ecv
Title : Crystal structure of the ALS-related pathological mutant I113T of human apo Cu,Zn Superoxide Dismutase (SOD1)
Authors : Calderone, V.
Deposited on : 2008-09-02
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.006 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.43.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	7293 (1.90-1.90)
Clashscore	180529	8090 (1.90-1.90)
Ramachandran outliers	177936	8022 (1.90-1.90)
Sidechain outliers	177891	8022 (1.90-1.90)
RSRZ outliers	164620	7292 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	153	
1	B	153	
1	C	153	
1	D	153	

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 4291 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Superoxide dismutase [Cu-Zn].

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	153	Total	C	N	O	S	0	0	0
			1109	677	203	225	4			
1	B	153	Total	C	N	O	S	0	0	0
			1109	677	203	225	4			
1	C	126	Total	C	N	O	S	0	0	0
			914	565	165	180	4			
1	D	125	Total	C	N	O	S	0	0	0
			910	563	164	179	4			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	113	THR	ILE	conflict	UNP P00441
B	113	THR	ILE	conflict	UNP P00441
C	113	THR	ILE	conflict	UNP P00441
D	113	THR	ILE	conflict	UNP P00441


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	71	Total	O	0	0
			71	71		
2	B	78	Total	O	0	0
			78	78		
2	C	49	Total	O	0	0
			49	49		
2	D	51	Total	O	0	0
			51	51		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Superoxide dismutase [Cu-Zn]




• Molecule 1: Superoxide dismutase [Cu-Zn]



• Molecule 1: Superoxide dismutase [Cu-Zn]



• Molecule 1: Superoxide dismutase [Cu-Zn]



G114	L117	V118	V119	H120	E121	D124	ASP	LEU	GLY	LYS	GLY	GLY	GLY	ASN	GLU	GLU	SER	THR	LYS	THR	GLY	ASN	ALA	GLY	S142	R143	L144	A145	C146	I149	Q153
------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	155.90Å 34.40Å 115.00Å 90.00° 112.10° 90.00°	Depositor
Resolution (Å)	38.95 – 1.90 38.95 – 1.90	Depositor EDS
% Data completeness (in resolution range)	100.0 (38.95-1.90) 98.3 (38.95-1.90)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.77 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.238 , 0.277 0.238 , 0.275	Depositor DCC
R_{free} test set	4053 reflections (9.07%)	wwPDB-VP
Wilson B-factor (Å ²)	12.4	Xtriage
Anisotropy	1.382	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4291	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.47	8/1127 (0.7%)	1.47	11/1519 (0.7%)
1	B	1.50	6/1127 (0.5%)	1.44	11/1519 (0.7%)
1	C	1.46	6/928 (0.6%)	1.31	4/1252 (0.3%)
1	D	1.39	3/924 (0.3%)	1.42	10/1247 (0.8%)
All	All	1.46	23/4106 (0.6%)	1.42	36/5537 (0.7%)

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	148	VAL	CA-CB	7.74	1.63	1.54
1	B	5	VAL	CA-CB	7.69	1.63	1.55
1	C	104	ILE	CA-CB	7.53	1.61	1.54
1	B	47	VAL	CA-CB	7.13	1.62	1.54
1	D	104	ILE	CA-CB	6.95	1.62	1.54
1	A	13	PRO	C-O	6.84	1.31	1.23
1	B	149	ILE	N-CA	6.52	1.54	1.46
1	D	7	VAL	C-O	-6.01	1.17	1.24
1	B	27	GLY	C-O	6.01	1.32	1.24
1	C	50	PHE	CA-C	-5.98	1.45	1.52
1	A	94	VAL	CA-CB	5.97	1.61	1.54
1	A	5	VAL	CA-CB	5.90	1.63	1.55
1	A	2	THR	CA-C	5.88	1.60	1.52
1	C	147	GLY	N-CA	5.79	1.50	1.45
1	A	118	VAL	N-CA	5.78	1.53	1.46
1	B	145	ALA	CA-CB	5.73	1.63	1.53
1	C	18	ILE	CA-CB	5.62	1.61	1.54
1	A	148	VAL	CA-C	5.36	1.58	1.52
1	D	149	ILE	CA-CB	5.33	1.60	1.53
1	A	104	ILE	CA-CB	5.31	1.60	1.54
1	C	116	THR	CA-CB	5.31	1.63	1.53
1	C	113	THR	CA-CB	5.08	1.61	1.53
1	B	18	ILE	CA-CB	5.04	1.60	1.54

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2	THR	CB-CA-C	8.86	123.72	109.90
1	B	12	GLY	CA-C-N	-8.63	110.99	121.00
1	B	12	GLY	C-N-CA	-8.63	110.99	121.00
1	B	65	ASN	CA-C-N	-8.54	109.16	119.84
1	B	65	ASN	C-N-CA	-8.54	109.16	119.84
1	A	82	GLY	N-CA-C	8.05	123.33	114.40
1	D	65	ASN	CA-C-N	7.63	127.27	119.19
1	D	65	ASN	C-N-CA	7.63	127.27	119.19
1	C	53	ASN	CB-CA-C	7.24	122.79	111.13
1	A	120	HIS	CA-C-N	6.86	130.33	120.79
1	A	120	HIS	C-N-CA	6.86	130.33	120.79
1	C	151	ILE	N-CA-C	-6.62	100.21	109.21
1	C	49	GLU	N-CA-C	6.61	118.48	111.28
1	D	2	THR	N-CA-C	-6.61	105.38	113.50
1	B	119	VAL	CB-CA-C	6.28	119.38	110.84
1	B	109	ASP	N-CA-C	6.12	118.75	111.71
1	B	151	ILE	N-CA-C	-6.06	101.73	109.30
1	A	50	PHE	N-CA-C	6.05	119.91	110.17
1	D	16	GLY	N-CA-C	5.91	119.22	110.42
1	D	149	ILE	N-CA-C	5.85	115.96	108.12
1	B	121	GLU	N-CA-C	5.80	118.07	111.11
1	A	59	SER	N-CA-C	5.72	119.78	112.34
1	A	18	ILE	N-CA-C	5.56	115.95	108.17
1	A	5	VAL	CB-CA-C	5.45	120.99	111.18
1	A	57	CYS	CA-C-N	5.26	127.77	120.29
1	A	57	CYS	C-N-CA	5.26	127.77	120.29
1	D	114	GLY	N-CA-C	-5.25	108.03	115.43
1	B	57	CYS	N-CA-C	5.24	118.77	112.38
1	B	73	GLY	CA-C-N	-5.24	114.42	119.87
1	B	73	GLY	C-N-CA	-5.24	114.42	119.87
1	A	122	LYS	N-CA-C	5.21	118.17	110.46
1	D	49	GLU	N-CA-C	5.14	116.96	111.36
1	D	12	GLY	CA-C-N	5.13	125.95	120.47
1	D	12	GLY	C-N-CA	5.13	125.95	120.47
1	C	82	GLY	N-CA-C	5.06	121.58	115.31
1	D	119	VAL	CB-CA-C	5.02	117.67	110.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1109	0	1073	18	0
1	B	1109	0	1073	28	0
1	C	914	0	890	20	0
1	D	910	0	887	20	0
2	A	71	0	0	3	0
2	B	78	0	0	8	0
2	C	49	0	0	1	0
2	D	51	0	0	3	0
All	All	4291	0	3923	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:229:HOH:O	1:C:24:GLU:HG2	1.39	1.20
1:D:46:HIS:HD2	1:D:80:HIS:CE1	1.79	1.01
1:D:46:HIS:CD2	1:D:80:HIS:HE1	1.79	0.99
1:B:77:GLU:HB2	2:B:238:HOH:O	1.64	0.96
1:D:64:PHE:O	1:D:80:HIS:HD2	1.65	0.79
1:D:63:HIS:HB3	1:D:80:HIS:CD2	2.19	0.78
1:C:79:ARG:NH2	1:C:101:ASP:OD1	2.17	0.78
1:C:38:LEU:O	1:C:93:GLY:HA2	1.88	0.74
1:D:27:GLY:HA3	2:D:165:HOH:O	1.88	0.73
1:B:6:CYS:SG	1:B:147:GLY:O	2.49	0.70
1:A:69:ARG:NH1	1:A:77:GLU:O	2.24	0.70
1:D:110:HIS:HB2	2:D:185:HOH:O	1.92	0.69
1:C:40:GLU:OE1	1:C:91:LYS:HB3	1.94	0.66
2:A:168:HOH:O	1:B:9:LYS:HD3	1.96	0.66
1:A:121:GLU:HB2	1:A:142:SER:HB3	1.79	0.65
1:C:49:GLU:O	1:C:115:ARG:HG2	1.97	0.63
1:B:42:LEU:HB3	1:B:123:ALA:HB2	1.82	0.62
1:B:52:ASP:OD1	1:B:54:THR:HG23	1.99	0.61
1:C:100:GLU:HG2	2:C:176:HOH:O	2.00	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:80:HIS:CD2	1:D:82:GLY:H	2.19	0.60
1:D:46:HIS:CD2	1:D:80:HIS:CE1	2.68	0.58
1:D:90:ASP:OD1	1:D:90:ASP:C	2.47	0.57
1:B:26:ASN:HA	1:C:26:ASN:HA	1.86	0.57
1:A:125:ASP:C	1:A:125:ASP:OD1	2.49	0.55
1:B:9:LYS:HG3	2:B:183:HOH:O	2.07	0.55
1:C:79:ARG:HH22	1:C:101:ASP:CG	2.13	0.55
1:B:126:LEU:H	1:B:126:LEU:HD23	1.72	0.55
1:A:39:THR:HG23	2:A:160:HOH:O	2.07	0.54
1:D:80:HIS:HB3	2:D:205:HOH:O	2.08	0.54
1:B:52:ASP:OD1	1:B:52:ASP:C	2.51	0.52
1:D:64:PHE:O	1:D:80:HIS:CD2	2.56	0.52
1:B:79:ARG:HH22	1:B:101:ASP:CG	2.18	0.52
1:B:126:LEU:HD23	1:B:126:LEU:N	2.26	0.51
1:A:52:ASP:O	1:A:59:SER:HB2	2.12	0.50
1:A:66:PRO:HD2	1:A:81:VAL:CG2	2.41	0.50
1:C:38:LEU:O	1:C:93:GLY:CA	2.59	0.50
1:D:14:VAL:HG13	1:D:38:LEU:HD13	1.95	0.49
1:A:5:VAL:HG12	1:A:152:ALA:HB2	1.94	0.49
1:A:66:PRO:HD2	1:A:81:VAL:HG23	1.95	0.49
1:B:79:ARG:CD	1:B:80:HIS:O	2.61	0.49
1:B:79:ARG:NH2	1:B:101:ASP:OD1	2.40	0.48
1:B:42:LEU:HB3	1:B:123:ALA:CB	2.42	0.48
1:D:63:HIS:HB3	1:D:80:HIS:NE2	2.28	0.48
1:C:66:PRO:HD2	1:C:81:VAL:CG2	2.43	0.48
1:D:14:VAL:CG1	1:D:38:LEU:HD13	2.44	0.48
2:B:229:HOH:O	1:C:24:GLU:CG	2.22	0.47
1:C:80:HIS:N	1:C:80:HIS:CD2	2.83	0.47
1:C:121:GLU:HB3	1:C:141:GLY:HA3	1.97	0.47
1:D:117:LEU:O	1:D:146:CYS:HA	2.15	0.47
1:A:121:GLU:HB2	1:A:142:SER:CB	2.44	0.47
1:A:52:ASP:O	1:A:59:SER:CB	2.62	0.46
1:D:45:PHE:O	1:D:84:LEU:HB2	2.16	0.46
1:B:15:GLN:NE2	1:B:36:LYS:HE2	2.29	0.46
1:C:14:VAL:HG21	1:C:144:LEU:HB3	1.97	0.46
1:B:105:SER:O	1:B:111:CYS:HA	2.16	0.46
1:B:46:HIS:CE1	2:B:193:HOH:O	2.68	0.46
1:C:141:GLY:C	1:C:142:SER:O	2.58	0.46
1:B:100:GLU:OE1	2:B:239:HOH:O	2.21	0.45
1:A:38:LEU:O	1:A:93:GLY:HA2	2.17	0.45
1:A:41:GLY:O	1:A:89:ALA:N	2.43	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:117:LEU:O	1:A:146:CYS:HA	2.17	0.44
1:B:79:ARG:HD3	1:B:80:HIS:O	2.17	0.44
1:C:101:ASP:HB3	1:C:104:ILE:HG12	1.98	0.44
1:A:84:LEU:O	2:A:204:HOH:O	2.21	0.44
1:A:43:HIS:HB3	1:A:120:HIS:O	2.17	0.44
1:B:6:CYS:SG	1:B:147:GLY:C	3.01	0.44
1:B:20:PHE:CD1	1:B:20:PHE:N	2.85	0.44
1:D:63:HIS:HB3	1:D:80:HIS:CG	2.52	0.43
1:A:14:VAL:HG21	1:A:144:LEU:HB3	1.99	0.43
1:D:121:GLU:HA	1:D:144:LEU:HD11	2.01	0.43
1:C:46:HIS:CD2	1:C:83:ASP:HA	2.53	0.43
1:A:116:THR:CG2	1:A:146:CYS:HB2	2.48	0.43
1:B:31:VAL:HB	1:B:99:ILE:HB	1.99	0.43
1:B:74:PRO:HG2	2:B:216:HOH:O	2.18	0.43
1:B:118:VAL:HG11	1:B:143:ARG:HG2	2.01	0.43
1:B:40:GLU:HG3	1:B:90:ASP:O	2.18	0.42
1:A:71:HIS:NE2	1:A:124:ASP:OD1	2.43	0.42
1:D:14:VAL:HG21	1:D:144:LEU:HB3	2.02	0.42
1:C:31:VAL:HB	1:C:99:ILE:HB	2.03	0.41
1:C:28:PRO:HB3	1:C:102:SER:OG	2.20	0.41
1:B:125:ASP:OD1	1:B:134:SER:OG	2.38	0.41
1:D:65:ASN:OD1	1:D:80:HIS:HA	2.20	0.41
1:C:35:ILE:CG2	1:C:38:LEU:HD21	2.50	0.41
1:B:85:GLY:N	2:B:162:HOH:O	2.54	0.40
1:B:79:ARG:HD2	1:B:80:HIS:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	151/153 (99%)	147 (97%)	4 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	151/153 (99%)	143 (95%)	8 (5%)	0	100	100
1	C	120/153 (78%)	112 (93%)	8 (7%)	0	100	100
1	D	119/153 (78%)	117 (98%)	2 (2%)	0	100	100
All	All	541/612 (88%)	519 (96%)	22 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	118/118 (100%)	107 (91%)	11 (9%)	7	3
1	B	118/118 (100%)	108 (92%)	10 (8%)	8	3
1	C	98/118 (83%)	92 (94%)	6 (6%)	15	8
1	D	98/118 (83%)	89 (91%)	9 (9%)	7	3
All	All	432/472 (92%)	396 (92%)	36 (8%)	9	4

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	THR
1	A	8	LEU
1	A	24	GLU
1	A	25	SER
1	A	31	VAL
1	A	38	LEU
1	A	98	SER
1	A	107	SER
1	A	128	LYS
1	A	135	THR
1	A	142	SER
1	B	8	LEU
1	B	9	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	34	SER
1	B	38	LEU
1	B	59	SER
1	B	75	LYS
1	B	79	ARG
1	B	91	LYS
1	B	98	SER
1	B	122	LYS
1	C	36	LYS
1	C	53	ASN
1	C	79	ARG
1	C	91	LYS
1	C	122	LYS
1	C	142	SER
1	D	24	GLU
1	D	31	VAL
1	D	38	LEU
1	D	42	LEU
1	D	62	PRO
1	D	67	LEU
1	D	80	HIS
1	D	91	LYS
1	D	96	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	15	GLN
1	C	19	ASN
1	C	46	HIS
1	C	80	HIS
1	D	26	ASN
1	D	46	HIS
1	D	53	ASN
1	D	80	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	153/153 (100%)	0.06	2 (1%) 74 76	4, 15, 27, 39	0
1	B	153/153 (100%)	0.08	0 100 100	3, 14, 26, 34	0
1	C	126/153 (82%)	0.27	7 (5%) 31 32	3, 16, 32, 40	2 (1%)
1	D	125/153 (81%)	0.27	5 (4%) 43 44	6, 17, 31, 43	0
All	All	557/612 (91%)	0.16	14 (2%) 58 60	3, 16, 30, 43	2 (0%)

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	67	LEU	5.1
1	C	141	GLY	5.0
1	C	67	LEU	4.2
1	A	1	ALA	3.4
1	D	1	ALA	2.9
1	C	79	ARG	2.7
1	C	124	ASP	2.5
1	D	25	SER	2.3
1	D	66	PRO	2.3
1	C	109	ASP	2.3
1	D	80	HIS	2.2
1	C	96	ASP	2.0
1	C	80	HIS	2.0
1	A	26	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.