



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2024 – 10:52 AM EDT

PDB ID : 6ECV  
Title : StiD O-MT residues 976-1266  
Authors : Skiba, M.A.; Bivins, M.M.; Smith, J.L.  
Deposited on : 2018-08-08  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	<div> <div>5%</div> <div> <div></div> <div>82%</div> <div>8%</div> <div>9%</div> </div> </div>
1	B	316	<div> <div>6%</div> <div> <div></div> <div>79%</div> <div>9%</div> <div>12%</div> </div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4995 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called StiD protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	278	Total	C	N	O	S	0	11	0
			2306	1456	414	431	5			
1	A	286	Total	C	N	O	S	0	8	0
			2322	1464	414	438	6			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	952	MET	-	initiating methionine	UNP Q8RJY3
B	953	HIS	-	expression tag	UNP Q8RJY3
B	954	HIS	-	expression tag	UNP Q8RJY3
B	955	HIS	-	expression tag	UNP Q8RJY3
B	956	HIS	-	expression tag	UNP Q8RJY3
B	957	HIS	-	expression tag	UNP Q8RJY3
B	958	HIS	-	expression tag	UNP Q8RJY3
B	959	SER	-	expression tag	UNP Q8RJY3
B	960	SER	-	expression tag	UNP Q8RJY3
B	961	GLY	-	expression tag	UNP Q8RJY3
B	962	VAL	-	expression tag	UNP Q8RJY3
B	963	ASP	-	expression tag	UNP Q8RJY3
B	964	LEU	-	expression tag	UNP Q8RJY3
B	965	GLY	-	expression tag	UNP Q8RJY3
B	966	THR	-	expression tag	UNP Q8RJY3
B	967	GLU	-	expression tag	UNP Q8RJY3
B	968	ASN	-	expression tag	UNP Q8RJY3
B	969	LEU	-	expression tag	UNP Q8RJY3
B	970	TYR	-	expression tag	UNP Q8RJY3
B	971	PHE	-	expression tag	UNP Q8RJY3
B	972	GLN	-	expression tag	UNP Q8RJY3
B	973	SER	-	expression tag	UNP Q8RJY3
B	974	ASN	-	expression tag	UNP Q8RJY3
B	975	ALA	-	expression tag	UNP Q8RJY3
A	952	MET	-	initiating methionine	UNP Q8RJY3

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	953	HIS	-	expression tag	UNP Q8RJY3
A	954	HIS	-	expression tag	UNP Q8RJY3
A	955	HIS	-	expression tag	UNP Q8RJY3
A	956	HIS	-	expression tag	UNP Q8RJY3
A	957	HIS	-	expression tag	UNP Q8RJY3
A	958	HIS	-	expression tag	UNP Q8RJY3
A	959	SER	-	expression tag	UNP Q8RJY3
A	960	SER	-	expression tag	UNP Q8RJY3
A	961	GLY	-	expression tag	UNP Q8RJY3
A	962	VAL	-	expression tag	UNP Q8RJY3
A	963	ASP	-	expression tag	UNP Q8RJY3
A	964	LEU	-	expression tag	UNP Q8RJY3
A	965	GLY	-	expression tag	UNP Q8RJY3
A	966	THR	-	expression tag	UNP Q8RJY3
A	967	GLU	-	expression tag	UNP Q8RJY3
A	968	ASN	-	expression tag	UNP Q8RJY3
A	969	LEU	-	expression tag	UNP Q8RJY3
A	970	TYR	-	expression tag	UNP Q8RJY3
A	971	PHE	-	expression tag	UNP Q8RJY3
A	972	GLN	-	expression tag	UNP Q8RJY3
A	973	SER	-	expression tag	UNP Q8RJY3
A	974	ASN	-	expression tag	UNP Q8RJY3
A	975	ALA	-	expression tag	UNP Q8RJY3

- Molecule 2 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula:  $C_{14}H_{20}N_6O_5S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
2	A	1	Total	C	N	O	S	0	0
			26	14	6	5	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	157	Total	O	0	0
			157	157		
4	A	157	Total	O	0	0
			157	157		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: StiD protein



#### • Molecule 1: StiD protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	40.18Å 56.17Å 72.25Å 86.31° 84.89° 75.81°	Depositor
Resolution (Å)	42.49 – 1.80 44.33 – 1.80	Depositor EDS
% Data completeness (in resolution range)	95.1 (42.49-1.80) 90.0 (44.33-1.80)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.70 (at 1.79Å)	Xtriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.167 , 0.194 0.170 , 0.196	Depositor DCC
$R_{free}$ test set	1964 reflections (3.64%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.9	Xtriage
Anisotropy	0.289	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 48.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4995	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.35	0/2367	0.52	0/3198
1	B	0.37	0/2351	0.53	0/3174
All	All	0.36	0/4718	0.52	0/6372

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2322	0	2284	15	0
1	B	2306	0	2273	18	0
2	A	26	0	19	1	0
2	B	26	0	19	3	0
3	A	1	0	0	1	0
4	A	157	0	0	1	0
4	B	157	0	0	1	0
All	All	4995	0	4595	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.



All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:1301:SAH:N7	3:A:1302:CL:CL	2.66	0.66
1:B:1242[A]:VAL:HG22	1:A:1134:PHE:HZ	1.62	0.64
1:A:1066:SER:O	4:A:1401:HOH:O	2.14	0.64
1:B:1209:ILE:HD13	1:B:1212:ARG:HH12	1.66	0.60
1:B:1109:PHE:O	2:B:1300:SAH:N	2.36	0.59
1:A:1011:MET:HE3	1:A:1015:LEU:HD11	1.87	0.55
1:B:1142:ALA:HB2	1:B:1157:ASN:HB3	1.90	0.54
1:A:1089:LEU:HD11	1:A:1099:LEU:HG	1.94	0.49
1:B:1184:VAL:HG21	1:B:1231[B]:TYR:CD2	2.49	0.48
1:A:1002:TYR:CD2	1:A:1200:LEU:HD11	2.49	0.48
1:B:1242[B]:VAL:HG12	1:A:1134:PHE:HZ	1.79	0.47
1:B:1112:ALA:O	1:B:1118:LYS:HE3	2.15	0.46
1:B:1113:THR:HG21	1:B:1140:PHE:HD2	1.79	0.46
1:A:1194:ASP:O	1:A:1198:THR:HG23	2.15	0.46
1:A:1074:LEU:HD21	1:A:1088:VAL:H	1.81	0.45
1:A:990:GLU:N	1:A:990:GLU:OE1	2.50	0.45
1:B:1071:GLU:O	1:B:1075:ARG:HG3	2.17	0.45
1:B:1220:PHE:CE2	1:B:1224:LEU:HD11	2.51	0.45
1:B:1243[A]:ARG:NH2	4:B:1411:HOH:O	2.49	0.45
1:B:982:TYR:CE1	1:B:1015:LEU:HD21	2.52	0.44
1:A:1261:TYR:O	1:A:1264:ARG:HB2	2.17	0.44
1:B:1242[A]:VAL:HG22	1:A:1134:PHE:CZ	2.49	0.43
2:B:1300:SAH:H5'1	2:B:1300:SAH:HB1	1.31	0.43
1:B:1239:ASP:CG	1:B:1242[A]:VAL:HG23	2.39	0.42
1:A:1005:ASP:OD2	1:A:1010:ARG:NH1	2.51	0.41
1:A:991:LYS:HE3	1:A:1189:PHE:CG	2.55	0.41
1:A:1106:ILE:HG13	1:A:1129:LEU:HD13	2.03	0.41
1:B:1134:PHE:HZ	1:A:1242[B]:VAL:HG22	1.85	0.41
2:B:1300:SAH:HN1	2:B:1300:SAH:HG2	1.72	0.41
1:B:1185:ALA:HB1	1:B:1225:GLU:HG2	2.03	0.41
1:B:1015:LEU:O	1:B:1019[B]:ARG:HG3	2.21	0.41
1:B:1044:TYR:HE1	1:B:1072:LEU:HD12	1.87	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	292/316 (92%)	286 (98%)	6 (2%)	0	100	100
1	B	285/316 (90%)	278 (98%)	7 (2%)	0	100	100
All	All	577/632 (91%)	564 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	250/268 (93%)	245 (98%)	5 (2%)	55	44
1	B	248/268 (92%)	244 (98%)	4 (2%)	62	54
All	All	498/536 (93%)	489 (98%)	9 (2%)	57	48

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	1090	ARG
1	B	1110	GLU
1	B	1113	THR
1	B	1199	GLN
1	A	1022	ARG
1	A	1110	GLU
1	A	1139	ASP
1	A	1231	TYR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	1264	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	SAH	B	1300	-	23,28,28	1.24	3 (13%)	22,40,40	2.15	6 (27%)
2	SAH	A	1301	-	23,28,28	1.25	3 (13%)	22,40,40	2.37	6 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAH	B	1300	-	-	5/11/31/31	0/3/3/3
2	SAH	A	1301	-	-	4/11/31/31	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1300	SAH	C2-N3	4.10	1.38	1.32
2	A	1301	SAH	C2-N3	3.94	1.38	1.32
2	A	1301	SAH	C2-N1	2.49	1.38	1.33
2	B	1300	SAH	C2-N1	2.46	1.38	1.33
2	B	1300	SAH	OXT-C	-2.22	1.23	1.30
2	A	1301	SAH	OXT-C	-2.14	1.23	1.30

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1300	SAH	N3-C2-N1	-6.87	119.34	128.67
2	A	1301	SAH	N3-C2-N1	-6.41	119.98	128.67
2	A	1301	SAH	C5'-SD-CG	-5.27	86.63	102.26
2	A	1301	SAH	O4'-C1'-N9	4.82	115.13	108.75
2	B	1300	SAH	O4'-C1'-N9	3.68	113.63	108.75
2	A	1301	SAH	C5'-C4'-C3'	-3.05	107.42	115.06
2	B	1300	SAH	C5'-SD-CG	-2.96	93.49	102.26
2	B	1300	SAH	OXT-C-O	-2.68	118.01	124.08
2	A	1301	SAH	OXT-C-O	-2.63	118.11	124.08
2	B	1300	SAH	C1'-N9-C4	-2.38	122.46	126.64
2	B	1300	SAH	C4'-O4'-C1'	-2.17	107.94	109.92
2	A	1301	SAH	C4'-O4'-C1'	-2.11	107.99	109.92

There are no chirality outliers.

All (9) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	1300	SAH	N-CA-CB-CG
2	B	1300	SAH	O4'-C4'-C5'-SD
2	A	1301	SAH	CA-CB-CG-SD
2	B	1300	SAH	CB-CG-SD-C5'
2	B	1300	SAH	CA-CB-CG-SD
2	A	1301	SAH	C-CA-CB-CG
2	B	1300	SAH	C3'-C4'-C5'-SD
2	A	1301	SAH	N-CA-CB-CG
2	A	1301	SAH	CB-CG-SD-C5'


There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1300	SAH	3	0
2	A	1301	SAH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	286/316 (90%)	0.10	16 (5%) 24 19	17, 39, 75, 123	0
1	B	278/316 (87%)	0.13	20 (7%) 15 12	17, 35, 77, 109	0
All	All	564/632 (89%)	0.11	36 (6%) 19 15	17, 37, 77, 123	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	982	TYR	7.2
1	A	1072	LEU	4.5
1	B	1074	LEU	4.2
1	B	1204	VAL	3.8
1	A	1265	GLU	3.8
1	A	1205	GLY	3.4
1	B	1072	LEU	3.3
1	A	981	SER	3.3
1	B	1212	ARG	3.1
1	B	1144	SER	3.1
1	A	980	GLU	3.0
1	B	1206	ILE	3.0
1	A	1068	GLU	2.9
1	B	1203	SER	2.7
1	A	1067	SER	2.7
1	B	1071	GLU	2.7
1	B	1265	GLU	2.6
1	B	982	TYR	2.6
1	A	1079	ALA	2.6
1	B	1207	SER	2.6
1	B	1202	THR	2.6
1	B	1155	SER	2.5
1	A	1078	GLU	2.5
1	B	1078	GLU	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	1209	ILE	2.4
1	A	1044	TYR	2.3
1	A	1071	GLU	2.3
1	A	1075	ARG	2.3
1	B	1205	GLY	2.3
1	A	1080	ARG	2.2
1	B	1068	GLU	2.2
1	B	1209	ILE	2.2
1	B	1075	ARG	2.1
1	A	1074	LEU	2.1
1	B	1156	TYR	2.1
1	B	978	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CL	A	1302	1/1	0.85	0.10	66,66,66,66	0
2	SAH	A	1301	26/26	0.95	0.15	23,42,79,82	0
2	SAH	B	1300	26/26	0.96	0.16	24,41,107,110	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






**Electron density around SAH A 1301:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around SAH B 1300:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.