



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 17, 2024 – 01:27 AM EDT

PDB ID : 3EEG  
Title : Crystal structure of a 2-isopropylmalate synthase from *Cytophaga hutchinsonii*  
Authors : Sugadev, R.; Burley, S.K.; Swaminathan, S.; New York SGX Research Center  
for Structural Genomics (NYSGXRC)  
Deposited on : 2008-09-04  
Resolution : 2.78 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.78 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4107 (2.80-2.76)
Clashscore	141614	4575 (2.80-2.76)
Ramachandran outliers	138981	4487 (2.80-2.76)
Sidechain outliers	138945	4489 (2.80-2.76)
RSRZ outliers	127900	4027 (2.80-2.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	325	
1	B	325	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4170 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-isopropylmalate synthase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	Se	0	0	0
			2029	1267	358	391	5	8			
1	B	273	Total	C	N	O	S	Se	0	0	0
			2085	1301	371	400	5	8			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MSE	-	expression tag	UNP Q11NN9
A	0	SER	-	expression tag	UNP Q11NN9
A	1	LEU	-	expression tag	UNP Q11NN9
A	316	GLU	-	expression tag	UNP Q11NN9
A	317	GLY	-	expression tag	UNP Q11NN9
A	318	HIS	-	expression tag	UNP Q11NN9
A	319	HIS	-	expression tag	UNP Q11NN9
A	320	HIS	-	expression tag	UNP Q11NN9
A	321	HIS	-	expression tag	UNP Q11NN9
A	322	HIS	-	expression tag	UNP Q11NN9
A	323	HIS	-	expression tag	UNP Q11NN9
B	-1	MSE	-	expression tag	UNP Q11NN9
B	0	SER	-	expression tag	UNP Q11NN9
B	1	LEU	-	expression tag	UNP Q11NN9
B	316	GLU	-	expression tag	UNP Q11NN9
B	317	GLY	-	expression tag	UNP Q11NN9
B	318	HIS	-	expression tag	UNP Q11NN9
B	319	HIS	-	expression tag	UNP Q11NN9
B	320	HIS	-	expression tag	UNP Q11NN9
B	321	HIS	-	expression tag	UNP Q11NN9
B	322	HIS	-	expression tag	UNP Q11NN9
B	323	HIS	-	expression tag	UNP Q11NN9

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	39	Total 39	O 39	0	0
2	B	17	Total 17	O 17	0	0

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Chain A:

47% 31% 18%

1 MSE SER LEU G2 K3 R4 I5 F8 D9 T10 T11 G15 GLU GLN VAL PRO GLY CYS GLN L23 N24 T25 E26 E27 I30 A34 L38 G39 V40 P41 V42 I43 P48 V49 N56 S57 V58 V59 E60 I61 R67 P68 T69 I70 C71 T74 R75 D80 A81 A91 R92 R93 S94 R95 I96 H97 T98 G99 I100 S103 D104 I105 H106 I107 F108 HIS LYS LEU ARG S113 T114 R115 E116 N117 I118 L119 E120 M121 A122 V126 K131 V132 V133 H134 E137 F138 F139 C140 A143 G144 R145 Q148 E156 A157 V158 I159 E160 A163 V164 V165 N166 N167 D170 T171 T172 M175 L176 T177 P177 Y180 R183 I188 M188 V191 S192 N193 I194 D195 K196 A197 I198 L207 G208 L209 A210 T211 A212 N213 S214 H1215 S216 A217 L218 H1219 G221 A222 R223 P224 V225 L226 C227 H1228 T229 N230 E242 E243 V244 V245 M248 E249 H1252 K253 H254

[illegible]

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	118.67Å 118.67Å 154.91Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.95 – 2.78 38.95 – 2.78	Depositor EDS
% Data completeness (in resolution range)	96.2 (38.95-2.78) 96.3 (38.95-2.78)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.59 (at 2.77Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.245 , 0.298 0.243 , 0.291	Depositor DCC
$R_{free}$ test set	655 reflections (3.90%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.7	Xtriage
Anisotropy	0.124	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 52.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	4170	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	0/2045	0.73	0/2748
1	B	0.42	0/2103	0.67	1/2826 (0.0%)
All	All	0.45	0/4148	0.70	1/5574 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	121	MSE	N-CA-C	-5.61	95.86	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2029	0	2048	130	0
1	B	2085	0	2107	152	0
2	A	39	0	0	3	0
2	B	17	0	0	2	0
All	All	4170	0	4155	281	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 34.

The worst 5 of 281 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:103:SER:HB3	1:B:105:ILE:HG12	1.35	1.08
1:B:114:THR:HG23	1:B:118:ILE:HB	1.46	0.97
1:B:94:SER:H	1:B:134:HIS:HD2	1.13	0.96
1:B:170:ASP:HB2	1:B:175:MSE:HE3	1.45	0.96
1:A:9:ASP:OD1	1:A:11:THR:HG22	1.70	0.92

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/325 (80%)	236 (90%)	22 (8%)	3 (1%)	14	38
1	B	269/325 (83%)	235 (87%)	27 (10%)	7 (3%)	5	16
All	All	530/650 (82%)	471 (89%)	49 (9%)	10 (2%)	8	23

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	112	ARG
1	B	192	SER
1	A	120	GLU
1	B	16	GLU
1	B	116	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/258 (84%)	204 (94%)	13 (6%)	19	45
1	B	223/258 (86%)	196 (88%)	27 (12%)	5	13
All	All	440/516 (85%)	400 (91%)	40 (9%)	9	25

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	194	ILE
1	B	258	GLU
1	B	195	ASP
1	B	252	LYS
1	B	267	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	263	HIS
1	B	134	HIS
1	B	230	ASN
1	B	213	ASN
1	B	224	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	259/325 (79%)	-0.25	1 (0%) 92 92	9, 24, 44, 50	0
1	B	265/325 (81%)	0.06	8 (3%) 50 45	16, 36, 56, 66	0
All	All	524/650 (80%)	-0.09	9 (1%) 70 67	9, 30, 52, 66	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	194	ILE	3.0
1	B	113	SER	2.6
1	B	112	ARG	2.3
1	B	148	GLN	2.3
1	A	253	GLU	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.