



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2024 – 03:13 PM EDT

PDB ID : 6EK7
Title : YaxA from Yersinia enterocolitica
Authors : Braeuning, B.; Groll, M.
Deposited on : 2017-09-25
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	410	<div> <div>13%</div> <div>78%</div> <div>11%</div> <div>11%</div> </div>

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3133 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called YaxA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	364	2898	1828	495	567	8	0	3	0

- Molecule 2 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	8	6	2	0	0
2	A	1	8	6	2	0	0
2	A	1	8	6	2	0	0
2	A	1	8	6	2	0	0
2	A	1	8	6	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			8	6	2		
2	A	1	Total	C	O	0	0
			8	6	2		
2	A	1	Total	C	O	0	0
			8	6	2		
2	A	1	Total	C	O	0	0
			8	6	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	163	Total	O	0	0
			163	163		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: YaxA



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	203.76Å 24.05Å 109.42Å 90.00° 113.92° 90.00°	Depositor
Resolution (Å)	14.95 – 1.80 44.16 – 1.80	Depositor EDS
% Data completeness (in resolution range)	97.7 (14.95-1.80) 97.9 (44.16-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.07 (at 1.79Å)	Xtriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, R_{free}	0.190 , 0.228 0.195 , 0.195	Depositor DCC
R_{free} test set	2276 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	27.7	Xtriage
Anisotropy	0.932	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 53.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3133	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/2937	0.55	1/3953 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	255	LEU	CA-CB-CG	5.41	127.75	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	255	LEU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2898	0	2965	38	0
2	A	72	0	126	8	0
3	A	163	0	0	3	0
All	All	3133	0	3091	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:296:ASN:HA	1:A:299:ILE:HG12	1.65	0.78
1:A:360:GLN:HA	1:A:363:MET:HE2	1.71	0.72
1:A:116:ASP:HB3	2:A:507:MPD:H11	1.72	0.70
1:A:245:ILE:HD11	1:A:305:LEU:HB3	1.74	0.68
1:A:128:GLY:HA2	1:A:377:PHE:HE2	1.57	0.68
1:A:67:GLN:NE2	3:A:601:HOH:O	2.29	0.64
1:A:58:ARG:HH22	2:A:505:MPD:H4	1.67	0.59
1:A:187:GLN:OE1	1:A:358:GLN:NE2	2.35	0.58
1:A:179:MET:HG3	2:A:506:MPD:H53	1.86	0.58
1:A:236:LYS:O	1:A:240:GLU:HG2	2.06	0.56
1:A:360:GLN:HA	1:A:363:MET:CE	2.36	0.56
1:A:255:LEU:HA	1:A:258:ASP:H	1.74	0.53
1:A:234:MET:CG	1:A:315:LEU:HD23	2.39	0.52
1:A:245:ILE:HD11	1:A:305:LEU:CB	2.39	0.51
2:A:504:MPD:O4	2:A:504:MPD:O2	2.23	0.51
1:A:237:SER:O	1:A:241:LEU:HD12	2.11	0.51
1:A:245:ILE:HG23	1:A:249:ARG:HD2	1.94	0.49
2:A:501:MPD:H52	2:A:501:MPD:HM2	1.94	0.49
1:A:252:ILE:HD12	1:A:299:ILE:HG22	1.94	0.48
1:A:104:ASN:ND2	3:A:606:HOH:O	2.46	0.48
1:A:234:MET:HG2	1:A:315:LEU:HD23	1.98	0.46
1:A:249:ARG:HA	1:A:252:ILE:HG22	1.98	0.46
1:A:186:HIS:HD2	3:A:650:HOH:O	1.99	0.46
1:A:66:ARG:NH1	1:A:66:ARG:HG2	2.31	0.45
1:A:104:ASN:O	1:A:108:GLN:HG3	2.16	0.45
1:A:178:ASP:HB3	2:A:506:MPD:H11	1.98	0.45
1:A:45:PHE:O	1:A:397:ILE:HD13	2.17	0.45
1:A:128:GLY:HA2	1:A:377:PHE:CE2	2.45	0.44
1:A:105:ASP:HA	1:A:108:GLN:HE21	1.82	0.44
1:A:66:ARG:HG2	1:A:66:ARG:HH11	1.83	0.44
1:A:298:LEU:HA	1:A:301:GLU:HB2	1.99	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:182:ASP:OD2	1:A:186:HIS:HE1	2.01	0.42
1:A:299:ILE:HG13	1:A:300:SER:N	2.34	0.42
1:A:404:GLU:O	1:A:408:VAL:HG23	2.19	0.42
1:A:252:ILE:O	1:A:256:LYS:HD2	2.20	0.41
1:A:58:ARG:NH2	2:A:505:MPD:H4	2.34	0.41
1:A:53:LEU:HD13	1:A:103:TRP:CH2	2.56	0.41
1:A:363:MET:HE3	1:A:376:LYS:HD2	2.03	0.41
1:A:223:LYS:HD3	2:A:508:MPD:HM2	2.03	0.41
1:A:275:ILE:O	1:A:275:ILE:HG13	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	363/410 (88%)	358 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	322/357 (90%)	320 (99%)	2 (1%)	86	84

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	158	ASN
1	A	292	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	67	GLN
1	A	102	ASN
1	A	108	GLN
1	A	158	ASN
1	A	186	HIS
1	A	216	ASN
1	A	296	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MPD	A	506	-	7,7,7	0.29	0	9,10,10	0.31	0
2	MPD	A	504	-	7,7,7	0.32	0	9,10,10	0.24	0
2	MPD	A	502	-	7,7,7	0.32	0	9,10,10	0.47	0
2	MPD	A	501	-	7,7,7	0.31	0	9,10,10	0.46	0
2	MPD	A	505	-	7,7,7	0.32	0	9,10,10	0.46	0
2	MPD	A	507	-	7,7,7	0.31	0	9,10,10	0.22	0
2	MPD	A	508	-	7,7,7	0.30	0	9,10,10	0.24	0
2	MPD	A	509	-	7,7,7	0.32	0	9,10,10	0.61	0
2	MPD	A	503	-	7,7,7	0.29	0	9,10,10	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MPD	A	506	-	-	2/5/5/5	-
2	MPD	A	504	-	-	1/5/5/5	-
2	MPD	A	502	-	-	5/5/5/5	-
2	MPD	A	501	-	-	5/5/5/5	-
2	MPD	A	505	-	-	0/5/5/5	-
2	MPD	A	507	-	-	0/5/5/5	-
2	MPD	A	508	-	-	0/5/5/5	-
2	MPD	A	509	-	-	3/5/5/5	-
2	MPD	A	503	-	-	1/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	MPD	C2-C3-C4-O4
2	A	501	MPD	C2-C3-C4-C5
2	A	502	MPD	C2-C3-C4-C5
2	A	503	MPD	O2-C2-C3-C4
2	A	502	MPD	C2-C3-C4-O4
2	A	502	MPD	C1-C2-C3-C4
2	A	509	MPD	CM-C2-C3-C4
2	A	506	MPD	C2-C3-C4-C5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	509	MPD	C2-C3-C4-C5
2	A	501	MPD	O2-C2-C3-C4
2	A	502	MPD	O2-C2-C3-C4
2	A	504	MPD	C2-C3-C4-O4
2	A	506	MPD	C2-C3-C4-O4
2	A	501	MPD	C1-C2-C3-C4
2	A	501	MPD	CM-C2-C3-C4
2	A	502	MPD	CM-C2-C3-C4
2	A	509	MPD	C1-C2-C3-C4

There are no ring outliers.

6 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	506	MPD	2	0
2	A	504	MPD	1	0
2	A	501	MPD	1	0
2	A	505	MPD	2	0
2	A	507	MPD	1	0
2	A	508	MPD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	364/410 (88%)	0.54	54 (14%) 2 1	25, 40, 122, 155	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	275	ILE	8.6
1	A	274	ILE	8.4
1	A	284	PHE	7.4
1	A	286	ALA	6.5
1	A	267	PHE	6.5
1	A	281	GLY	6.5
1	A	271	ILE	5.8
1	A	265	LEU	5.7
1	A	285	GLY	5.6
1	A	262	PHE	5.2
1	A	273	GLY	5.0
1	A	291	ALA	4.8
1	A	276	ALA	4.8
1	A	290	ASN	4.8
1	A	269	GLY	4.7
1	A	268	THR	4.7
1	A	279	ILE	4.7
1	A	282	GLY	4.6
1	A	278	ALA	4.4
1	A	272	GLY	4.4
1	A	283	ILE	4.3
1	A	408	VAL	4.0
1	A	266	SER	4.0
1	A	259	TYR	3.9
1	A	263	VAL	3.8
1	A	280	THR	3.8
1	A	297	ALA	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	293	LYS	3.7
1	A	299	ILE	3.6
1	A	165	ASP	3.5
1	A	405	TYR	3.4
1	A	264	GLY	3.1
1	A	250	GLN	3.0
1	A	160	THR	3.0
1	A	305	LEU	2.9
1	A	407	LYS	2.9
1	A	255	LEU	2.7
1	A	289	GLU	2.7
1	A	44	ILE	2.6
1	A	409	TYR	2.6
1	A	300	SER	2.5
1	A	164	ALA	2.5
1	A	260	ASP	2.5
1	A	406	LYS	2.5
1	A	239	LYS	2.4
1	A	270	ALA	2.4
1	A	158	ASN	2.3
1	A	252	ILE	2.3
1	A	257	LYS	2.2
1	A	246	LYS	2.2
1	A	277	MET	2.2
1	A	157	GLU	2.2
1	A	296	ASN	2.1
1	A	304	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MPD	A	509	8/8	0.79	0.23	116,121,127,128	0
2	MPD	A	508	8/8	0.80	0.20	86,90,101,102	0
2	MPD	A	504	8/8	0.80	0.25	72,79,83,87	0
2	MPD	A	506	8/8	0.82	0.21	66,75,83,84	0
2	MPD	A	507	8/8	0.83	0.22	71,77,85,90	0
2	MPD	A	501	8/8	0.86	0.15	33,51,58,72	0
2	MPD	A	503	8/8	0.88	0.21	84,87,91,94	0
2	MPD	A	505	8/8	0.89	0.15	59,66,69,74	0
2	MPD	A	502	8/8	0.95	0.12	61,63,76,76	0

6.5 Other polymers [i](#)

There are no such residues in this entry.