



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2024 – 03:19 PM EDT

PDB ID : 6EK8  
Title : YaxB from Yersinia enterocolitica  
Authors : Braeuning, B.; Groll, M.  
Deposited on : 2017-09-25  
Resolution : 4.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 4.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1087 (4.30-3.70)
Clashscore	141614	1148 (4.30-3.70)
Ramachandran outliers	138981	1108 (4.30-3.70)
Sidechain outliers	138945	1099 (4.30-3.70)
RSRZ outliers	127900	1028 (4.34-3.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	344	

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2002 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called YaxB.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	249	Total	C	N	O	0	0	0
			2002	1266	341	395			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP A1JM52

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Chain A:

58% 13% 28%

GLY SER ALA GLU ILE THR PHE PRO HIS S10 G11 L12 P15 D16 I17 F22 M28 H31 L32 F35 LYS THR THR GLY VAL GLU VAL GLN E45 K46 V61 R62 E63 S64 L65 L68 L76 Y80 L84 A90 L90 ILE SER GLN ASP ILE THR

GLU SER LYS SER GLU MET ARG LYS GLU ARG I109 I127 E128 K129 I145 I146 E147 E156 R159 E173 I177 E178 R181 A187 ASP ASP VAL ILE ARG GLU HIS ASN LEU VAL ASP ALA PHE LYS ASP LEU ILE THR PRO GLY ASN LEU SER GLU LEU LEU

ALA PRO GLU ILE GLU LEU LEU GLN SER LEU LEU ILE THR LYS LEU LEU GLY GLN PHE SER GLU GLY LEU LYS TYR ILE ASP L246 R249 R250 K251 K252 V281 V284 I285 D288 K291 S292 E297 K300 H306 I307 F308 I309 Q316 L320 L321 L329 L336 L340 L343 ILE

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	111.70Å 111.70Å 169.66Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.82 – 4.00 48.82 – 4.00	Depositor EDS
% Data completeness (in resolution range)	98.7 (48.82-4.00) 98.7 (48.82-4.00)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.25 (at 4.00Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, $R_{free}$	0.326 , 0.339 0.320 , 0.339	Depositor DCC
$R_{free}$ test set	282 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	162.8	Xtriage
Anisotropy	0.375	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 175.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	2002	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	200.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.75% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.28	0/2016	0.47	0/2710

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2002	0	2090	27	1
All	All	2002	0	2090	27	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:65:LEU:HD13	1:A:340:ILE:HG12	1.77	0.66
1:A:249:ALA:HA	1:A:252:LYS:HG2	1.84	0.59
1:A:156:GLU:HA	1:A:159:ARG:HG2	1.88	0.56
1:A:76:LEU:HD21	1:A:329:LEU:HB3	1.88	0.55
1:A:15:PRO:HD3	1:A:307:ILE:HD12	1.88	0.55

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:178:GLU:HA	1:A:181:ARG:HE	1.71	0.54
1:A:146:ILE:HG22	1:A:147:GLU:H	1.73	0.53
1:A:15:PRO:HG2	1:A:17:ILE:HD11	1.90	0.52
1:A:68:LEU:HD21	1:A:129:LYS:HG3	1.92	0.51
1:A:145:ILE:HD12	1:A:284:VAL:HG22	1.93	0.51
1:A:281:VAL:O	1:A:284:VAL:HG12	2.11	0.50
1:A:127:ILE:HG12	1:A:309:ILE:HD11	1.92	0.50
1:A:178:GLU:HA	1:A:181:ARG:HG2	1.95	0.49
1:A:173:GLU:O	1:A:177:ILE:HG12	2.15	0.47
1:A:281:VAL:O	1:A:285:ILE:HG12	2.14	0.47
1:A:320:LEU:HD23	1:A:320:LEU:H	1.81	0.46
1:A:320:LEU:HG	1:A:321:ASN:OD1	2.15	0.45
1:A:28:ASN:O	1:A:32:LEU:HB2	2.16	0.45
1:A:17:ILE:HD12	1:A:300:LYS:HB3	1.99	0.44
1:A:22:PHE:HB2	1:A:297:GLU:HG2	1.99	0.43
1:A:61:VAL:O	1:A:65:LEU:HG	2.19	0.43
1:A:65:LEU:HB2	1:A:336:LEU:HD21	2.00	0.42
1:A:145:ILE:HD11	1:A:288:ASP:HB2	2.01	0.42
1:A:45:GLU:HG2	1:A:46:LYS:N	2.35	0.42
1:A:12:LEU:HD13	1:A:307:ILE:HG23	2.02	0.41
1:A:80:TYR:O	1:A:84:LEU:HB2	2.21	0.40
1:A:306:HIS:HA	1:A:309:ILE:HG22	2.03	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:292:SER:OG	1:A:292:SER:OG[10_445]	1.99	0.21

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	241/344 (70%)	234 (97%)	7 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	229/316 (72%)	219 (96%)	10 (4%)	28	55

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	LEU
1	A	31	HIS
1	A	45	GLU
1	A	62	ARG
1	A	63	GLU
1	A	252	LYS
1	A	288	ASP
1	A	291	LYS
1	A	316	GLN
1	A	343	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	249/344 (72%)	-0.26	2 (0%) 86 79	136, 196, 260, 296	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	246	LEU	3.6
1	A	250	ARG	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.