



## Full wwPDB EM Validation Report ⓘ

Jun 9, 2024 – 08:50 AM EDT

PDB ID : 8ENV  
EMDB ID : EMD-28280  
Title : In situ cryo-EM structure of Pseudomonas phage E217 tail baseplate in C6 map  
Authors : Li, F.; Cingolani, G.; Hou, C.  
Deposited on : 2022-09-30  
Resolution : 3.42 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev92  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*










The reported resolution of this entry is 3.42 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




























Metric	Whole archive (#Entries)	EM structures (#Entries)
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	504	 100%
1	B	504	 100%
1	C	504	 100%
1	D	504	 100%
1	E	504	 100%
1	F	504	 100%
2	G	102	 99%
2	L	102	 99%
2	Q	102	 99%

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	V	102	
2	a	102	
2	f	102	
3	H	108	
3	M	108	
3	R	108	
3	W	108	
3	b	108	
3	g	108	
4	I	152	
4	N	152	
4	S	152	
4	X	152	
4	c	152	
4	h	152	
5	J	417	
5	O	417	
5	T	417	
5	Y	417	
5	d	417	
5	i	417	
6	K	500	
6	P	500	
6	U	500	
6	Z	500	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	e	500	<div><div></div><div>17%</div><div></div><div>100%</div></div>
6	j	500	<div><div></div><div>16%</div><div></div><div>99%</div></div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 80436 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sheath protein gp31.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	504	Total	C	N	O	S	0	0
			3780	2380	639	749	12		
1	B	504	Total	C	N	O	S	0	0
			3780	2380	639	749	12		
1	C	504	Total	C	N	O	S	0	0
			3780	2380	639	749	12		
1	D	504	Total	C	N	O	S	0	0
			3780	2380	639	749	12		
1	E	504	Total	C	N	O	S	0	0
			3780	2380	639	749	12		
1	F	504	Total	C	N	O	S	0	0
			3780	2380	639	749	12		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	17	ALA	GLY	conflict	UNP A0A2K8IA62
B	17	ALA	GLY	conflict	UNP A0A2K8IA62
C	17	ALA	GLY	conflict	UNP A0A2K8IA62
D	17	ALA	GLY	conflict	UNP A0A2K8IA62
E	17	ALA	GLY	conflict	UNP A0A2K8IA62
F	17	ALA	GLY	conflict	UNP A0A2K8IA62

- Molecule 2 is a protein called Structural protein gp33.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	G	102	Total	C	N	O	S	0	0
			816	525	136	150	5		
2	L	102	Total	C	N	O	S	0	0
			816	525	136	150	5		
2	Q	102	Total	C	N	O	S	0	0
			816	525	136	150	5		
2	V	102	Total	C	N	O	S	0	0
			816	525	136	150	5		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					AltConf	Trace
2	a	102	Total	C	N	O	S	0	0
			816	525	136	150	5		
2	f	102	Total	C	N	O	S	0	0
			816	525	136	150	5		

- Molecule 3 is a protein called Sheath initiator gp34.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	H	108	Total	C	N	O	S	0	0
			823	510	137	172	4		
3	M	108	Total	C	N	O	S	0	0
			823	510	137	172	4		
3	R	108	Total	C	N	O	S	0	0
			823	510	137	172	4		
3	W	108	Total	C	N	O	S	0	0
			823	510	137	172	4		
3	b	108	Total	C	N	O	S	0	0
			823	510	137	172	4		
3	g	108	Total	C	N	O	S	0	0
			823	510	137	172	4		

- Molecule 4 is a protein called Ripcord gp36.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	I	152	Total	C	N	O	S	0	0
			1133	726	182	221	4		
4	N	152	Total	C	N	O	S	0	0
			1133	726	182	221	4		
4	S	152	Total	C	N	O	S	0	0
			1133	726	182	221	4		
4	X	152	Total	C	N	O	S	0	0
			1133	726	182	221	4		
4	c	152	Total	C	N	O	S	0	0
			1133	726	182	221	4		
4	h	152	Total	C	N	O	S	0	0
			1133	726	182	221	4		

- Molecule 5 is a protein called Baseplate\_J domain-containing protein gp44.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	J	417	Total	C	N	O	S	0	0
			3036	1901	526	597	12		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					AltConf	Trace
5	O	417	Total	C	N	O	S	0	0
			3036	1901	526	597	12		
5	T	417	Total	C	N	O	S	0	0
			3036	1901	526	597	12		
5	Y	417	Total	C	N	O	S	0	0
			3036	1901	526	597	12		
5	d	417	Total	C	N	O	S	0	0
			3036	1901	526	597	12		
5	i	417	Total	C	N	O	S	0	0
			3036	1901	526	597	12		


- Molecule 6 is a protein called Structural protein gp45.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	K	500	Total	C	N	O	S	0	0
			3818	2435	641	729	13		
6	P	500	Total	C	N	O	S	0	0
			3818	2435	641	729	13		
6	U	500	Total	C	N	O	S	0	0
			3818	2435	641	729	13		
6	Z	500	Total	C	N	O	S	0	0
			3818	2435	641	729	13		
6	e	500	Total	C	N	O	S	0	0
			3818	2435	641	729	13		
6	j	500	Total	C	N	O	S	0	0
			3818	2435	641	729	13		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Sheath protein gp31




- Molecule 1: Sheath protein gp31




- Molecule 1: Sheath protein gp31



- Molecule 1: Sheath protein gp31




- Molecule 1: Sheath protein gp31





- Molecule 1: Sheath protein gp31

Chain F:  100%



- Molecule 2: Structural protein gp33

Chain G:  99%



- Molecule 2: Structural protein gp33

Chain L:  99%



- Molecule 2: Structural protein gp33

Chain Q:  99%



- Molecule 2: Structural protein gp33

Chain V:  97%




- Molecule 2: Structural protein gp33

Chain a:  98%




- Molecule 2: Structural protein gp33

Chain f:  99%




- Molecule 3: Sheath initiator gp34




- Molecule 3: Sheath initiator gp34



- Molecule 3: Sheath initiator gp34



- Molecule 3: Sheath initiator gp34



- Molecule 3: Sheath initiator gp34




- Molecule 3: Sheath initiator gp34




- Molecule 4: Ripcord gp36

Chain I:  99%




- Molecule 4: Ripcord gp36

Chain N:  100%




- Molecule 4: Ripcord gp36

Chain S:  99%




- Molecule 4: Ripcord gp36

Chain X:  100%




- Molecule 4: Ripcord gp36

Chain c:  99%




- Molecule 4: Ripcord gp36

Chain h:  99%




- Molecule 5: Baseplate\_J domain-containing protein gp44


Chain J:  6%  100%




- Molecule 5: Baseplate\_J domain-containing protein gp44




- Molecule 5: Baseplate\_J domain-containing protein gp44




- Molecule 5: Baseplate\_J domain-containing protein gp44




- Molecule 5: Baseplate\_J domain-containing protein gp44




- Molecule 5: Baseplate\_J domain-containing protein gp44



- Molecule 6: Structural protein gp45






- Molecule 6: Structural protein gp45




- Molecule 6: Structural protein gp45




- Molecule 6: Structural protein gp45



- Molecule 6: Structural protein gp45



• Molecule 6: Structural protein gp45



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C6	Depositor
Number of particles used	9300	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	50	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	1500	Depositor
Magnification	81000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.091	Depositor
Minimum map value	-0.063	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.005	Depositor
Recommended contour level	0.01	Depositor
Map size (Å)	448.0, 448.0, 448.0	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.12, 1.12, 1.12	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.31	0/3849	0.51	1/5259 (0.0%)
1	B	0.31	0/3849	0.50	0/5259
1	C	0.31	0/3849	0.50	0/5259
1	D	0.31	0/3849	0.49	0/5259
1	E	0.31	0/3849	0.50	0/5259
1	F	0.31	0/3849	0.50	0/5259
2	G	0.32	0/839	0.55	0/1139
2	L	0.31	0/839	0.50	0/1139
2	Q	0.32	0/839	0.52	0/1139
2	V	0.31	0/839	0.52	0/1139
2	a	0.33	0/839	0.55	1/1139 (0.1%)
2	f	0.32	0/839	0.53	0/1139
3	H	0.30	0/832	0.53	0/1128
3	M	0.31	0/832	0.56	0/1128
3	R	0.31	0/832	0.55	0/1128
3	W	0.31	0/832	0.57	0/1128
3	b	0.31	0/832	0.54	0/1128
3	g	0.31	0/832	0.56	0/1128
4	I	0.34	0/1155	0.55	0/1563
4	N	0.33	0/1155	0.54	0/1563
4	S	0.34	0/1155	0.53	0/1563
4	X	0.33	0/1155	0.53	0/1563
4	c	0.34	0/1155	0.54	0/1563
4	h	0.33	0/1155	0.54	0/1563
5	J	0.27	0/3090	0.52	0/4225
5	O	0.27	0/3090	0.52	0/4225
5	T	0.27	0/3090	0.52	0/4225
5	Y	0.27	0/3090	0.52	0/4225
5	d	0.28	0/3090	0.53	0/4225
5	i	0.27	0/3090	0.52	0/4225
6	K	0.26	0/3922	0.55	1/5363 (0.0%)
6	P	0.32	1/3922 (0.0%)	0.58	4/5363 (0.1%)
6	U	0.27	0/3922	0.53	1/5363 (0.0%)
6	Z	0.27	0/3922	0.55	1/5363 (0.0%)



Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
6	e	0.26	0/3922	0.52	0/5363
6	j	0.27	0/3922	0.55	3/5363 (0.1%)
All	All	0.29	1/82122 (0.0%)	0.53	12/112062 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	G	0	1
2	L	0	1
2	Q	0	1
2	V	0	1
2	a	0	1
2	f	0	1
5	i	0	1
6	j	0	1
All	All	0	8

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	P	452	PRO	CG-CD	-10.79	1.15	1.50

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	P	452	PRO	N-CD-CG	-13.70	82.65	103.20
6	P	452	PRO	CA-CB-CG	-7.10	90.52	104.00
6	P	452	PRO	CA-N-CD	-6.12	102.93	111.50
2	a	67	ASP	CB-CG-OD1	6.00	123.70	118.30
6	K	130	LEU	CA-CB-CG	5.29	127.47	115.30
6	Z	130	LEU	CA-CB-CG	5.27	127.43	115.30
6	j	452	PRO	N-CD-CG	-5.24	95.34	103.20
6	P	130	LEU	CA-CB-CG	5.16	127.18	115.30
1	A	2	ILE	CG1-CB-CG2	-5.05	100.28	111.40
6	U	264	PRO	N-CD-CG	-5.05	95.62	103.20
6	j	16	TRP	CA-CB-CG	-5.02	104.15	113.70
6	j	130	LEU	CA-CB-CG	5.02	126.86	115.30

There are no chirality outliers.

All (8) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	G	65	ARG	Peptide
2	L	65	ARG	Peptide
2	Q	65	ARG	Peptide
2	V	65	ARG	Peptide
2	a	65	ARG	Peptide
2	f	65	ARG	Peptide
5	i	118	GLN	Peptide
6	j	16	TRP	Peptide

## 5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	502/504 (100%)	485 (97%)	16 (3%)	1 (0%)	47	80
1	B	502/504 (100%)	487 (97%)	14 (3%)	1 (0%)	47	80
1	C	502/504 (100%)	479 (95%)	22 (4%)	1 (0%)	47	80
1	D	502/504 (100%)	483 (96%)	18 (4%)	1 (0%)	47	80
1	E	502/504 (100%)	478 (95%)	23 (5%)	1 (0%)	47	80
1	F	502/504 (100%)	477 (95%)	24 (5%)	1 (0%)	47	80
2	G	100/102 (98%)	94 (94%)	6 (6%)	0	100	100
2	L	100/102 (98%)	95 (95%)	5 (5%)	0	100	100
2	Q	100/102 (98%)	94 (94%)	6 (6%)	0	100	100
2	V	100/102 (98%)	93 (93%)	7 (7%)	0	100	100
2	a	100/102 (98%)	94 (94%)	6 (6%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	f	100/102 (98%)	93 (93%)	7 (7%)	0	100	100
3	H	106/108 (98%)	96 (91%)	8 (8%)	2 (2%)	8	37
3	M	106/108 (98%)	97 (92%)	7 (7%)	2 (2%)	8	37
3	R	106/108 (98%)	97 (92%)	7 (7%)	2 (2%)	8	37
3	W	106/108 (98%)	95 (90%)	10 (9%)	1 (1%)	17	53
3	b	106/108 (98%)	96 (91%)	8 (8%)	2 (2%)	8	37
3	g	106/108 (98%)	94 (89%)	11 (10%)	1 (1%)	17	53
4	I	150/152 (99%)	142 (95%)	6 (4%)	2 (1%)	12	45
4	N	150/152 (99%)	141 (94%)	9 (6%)	0	100	100
4	S	150/152 (99%)	139 (93%)	9 (6%)	2 (1%)	12	45
4	X	150/152 (99%)	141 (94%)	9 (6%)	0	100	100
4	c	150/152 (99%)	143 (95%)	6 (4%)	1 (1%)	22	58
4	h	150/152 (99%)	143 (95%)	5 (3%)	2 (1%)	12	45
5	J	415/417 (100%)	373 (90%)	42 (10%)	0	100	100
5	O	415/417 (100%)	361 (87%)	54 (13%)	0	100	100
5	T	415/417 (100%)	368 (89%)	47 (11%)	0	100	100
5	Y	415/417 (100%)	368 (89%)	47 (11%)	0	100	100
5	d	415/417 (100%)	361 (87%)	54 (13%)	0	100	100
5	i	415/417 (100%)	365 (88%)	50 (12%)	0	100	100
6	K	498/500 (100%)	461 (93%)	36 (7%)	1 (0%)	47	80
6	P	498/500 (100%)	458 (92%)	39 (8%)	1 (0%)	47	80
6	U	498/500 (100%)	457 (92%)	41 (8%)	0	100	100
6	Z	498/500 (100%)	458 (92%)	40 (8%)	0	100	100
6	e	498/500 (100%)	460 (92%)	38 (8%)	0	100	100
6	j	498/500 (100%)	461 (93%)	37 (7%)	0	100	100
All	All	10626/10698 (99%)	9827 (92%)	774 (7%)	25 (0%)	50	80

All (25) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2	ILE
1	C	2	ILE
1	D	2	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	F	2	ILE
4	S	89	PHE
4	c	89	PHE
4	h	89	PHE
1	B	2	ILE
1	E	2	ILE
4	I	89	PHE
4	I	96	ILE
4	S	96	ILE
4	h	96	ILE
6	K	454	TYR
6	P	454	TYR
3	M	92	ILE
3	g	92	ILE
3	R	94	ILE
3	H	94	ILE
3	W	92	ILE
3	H	92	ILE
3	M	94	ILE
3	R	92	ILE
3	b	92	ILE
3	b	94	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	400/400 (100%)	400 (100%)	0	100	100
1	B	400/400 (100%)	400 (100%)	0	100	100
1	C	400/400 (100%)	399 (100%)	1 (0%)	92	97
1	D	400/400 (100%)	399 (100%)	1 (0%)	92	97
1	E	400/400 (100%)	399 (100%)	1 (0%)	92	97
1	F	400/400 (100%)	400 (100%)	0	100	100
2	G	84/84 (100%)	83 (99%)	1 (1%)	71	87

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	L	84/84 (100%)	83 (99%)	1 (1%)	71	87
2	Q	84/84 (100%)	84 (100%)	0	100	100
2	V	84/84 (100%)	81 (96%)	3 (4%)	35	66
2	a	84/84 (100%)	83 (99%)	1 (1%)	71	87
2	f	84/84 (100%)	84 (100%)	0	100	100
3	H	92/92 (100%)	92 (100%)	0	100	100
3	M	92/92 (100%)	92 (100%)	0	100	100
3	R	92/92 (100%)	92 (100%)	0	100	100
3	W	92/92 (100%)	92 (100%)	0	100	100
3	b	92/92 (100%)	92 (100%)	0	100	100
3	g	92/92 (100%)	92 (100%)	0	100	100
4	I	124/124 (100%)	124 (100%)	0	100	100
4	N	124/124 (100%)	124 (100%)	0	100	100
4	S	124/124 (100%)	124 (100%)	0	100	100
4	X	124/124 (100%)	124 (100%)	0	100	100
4	c	124/124 (100%)	124 (100%)	0	100	100
4	h	124/124 (100%)	124 (100%)	0	100	100
5	J	320/320 (100%)	320 (100%)	0	100	100
5	O	320/320 (100%)	318 (99%)	2 (1%)	86	94
5	T	320/320 (100%)	318 (99%)	2 (1%)	86	94
5	Y	320/320 (100%)	316 (99%)	4 (1%)	69	86
5	d	320/320 (100%)	320 (100%)	0	100	100
5	i	320/320 (100%)	317 (99%)	3 (1%)	78	90
6	K	404/404 (100%)	404 (100%)	0	100	100
6	P	404/404 (100%)	404 (100%)	0	100	100
6	U	404/404 (100%)	404 (100%)	0	100	100
6	Z	404/404 (100%)	404 (100%)	0	100	100
6	e	404/404 (100%)	403 (100%)	1 (0%)	93	98
6	j	404/404 (100%)	403 (100%)	1 (0%)	93	98
All	All	8544/8544 (100%)	8522 (100%)	22 (0%)	92	97

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	69	LYS
1	D	490	LYS
1	E	256	ASN
2	G	65	ARG
2	L	65	ARG
5	O	106	ARG
5	O	177	ARG
5	T	106	ARG
5	T	256	LYS
2	V	33	MET
2	V	50	ARG
2	V	65	ARG
5	Y	106	ARG
5	Y	203	ARG
5	Y	256	LYS
5	Y	293	ARG
2	a	65	ARG
6	e	296	ARG
5	i	203	ARG
5	i	256	LYS
5	i	411	ASN
6	j	246	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (33) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	351	GLN
1	D	298	GLN
1	E	308	ASN
1	F	191	GLN
1	F	361	GLN
3	H	33	GLN
3	H	89	GLN
3	H	109	HIS
5	J	44	GLN
5	J	236	ASN
5	J	257	GLN
5	J	332	GLN
6	K	312	GLN
3	M	89	GLN
5	O	35	ASN
5	O	44	GLN
5	O	280	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
3	R	64	GLN
3	R	89	GLN
5	T	35	ASN
6	U	288	GLN
5	Y	118	GLN
3	b	33	GLN
5	d	257	GLN
5	d	340	GLN
6	e	312	GLN
6	e	402	ASN
3	g	64	GLN
3	g	89	GLN
3	g	109	HIS
4	h	116	ASN
5	i	280	ASN
6	j	312	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.




## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-28280. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


### 6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 180




Y Index: 220




Z Index: 82


### 6.3.2 Raw map



X Index: 179



Y Index: 220




Z Index: 147

The images above show the largest variance slices of the map in three orthogonal directions.




## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.01. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 1739 nm<sup>3</sup>; this corresponds to an approximate mass of 1571 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ




\*Reported resolution corresponds to spatial frequency of 0.292 Å<sup>-1</sup>



## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.292 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.42	-	-
Author-provided FSC curve	3.51	4.10	3.56
Unmasked-calculated*	3.96	4.75	4.05

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.96 differs from the reported value 3.42 by more than 10 %

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-28280 and PDB model 8ENV. Per-residue inclusion information can be found in section [3](#) on page [8](#).

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.01 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.01).







































































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 91% of all backbone atoms, 76% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary ⓘ




The table lists the average atom inclusion at the recommended contour level (0.01) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7610	 0.4130
A	 0.8390	 0.4620
B	 0.8410	 0.4620
C	 0.8350	 0.4620
D	 0.8340	 0.4610
E	 0.8340	 0.4620
F	 0.8380	 0.4640
G	 0.8330	 0.4750
H	 0.8250	 0.4840
I	 0.8270	 0.5140
J	 0.7360	 0.3740
K	 0.6560	 0.3310
L	 0.8490	 0.4830
M	 0.8270	 0.4850
N	 0.8230	 0.5100
O	 0.7350	 0.3750
P	 0.6560	 0.3340
Q	 0.8310	 0.4860
R	 0.8240	 0.4810
S	 0.8330	 0.5160
T	 0.7370	 0.3770
U	 0.6460	 0.3350
V	 0.8490	 0.4810
W	 0.8310	 0.4850
X	 0.8280	 0.5150
Y	 0.7390	 0.3750
Z	 0.6570	 0.3360
a	 0.8420	 0.4810
b	 0.8270	 0.4810
c	 0.8200	 0.5090
d	 0.7360	 0.3770
e	 0.6550	 0.3290
f	 0.8370	 0.4840
g	 0.8250	 0.4840
h	 0.8320	 0.5150



*Continued on next page...*

*Continued from previous page...*

Chain	Atom inclusion	Q-score
i	 0.7380	 0.3780
j	 0.6590	 0.3350