



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 13, 2024 – 01:04 AM EDT

PDB ID : 3ETE
Title : Crystal structure of bovine glutamate dehydrogenase complexed with hexachlorophene
Authors : Li, M.; Smith, T.J.
Deposited on : 2008-10-07
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.36.2
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2092 (3.00-3.00)
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	501	60% 34% 5% .
1	B	501	61% 34% . .
1	C	501	63% 32% . .
1	D	501	59% 36% . .
1	E	501	61% 34% . .
1	F	501	62% 33% . .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GLU	A	550	-	-	X	-
2	GLU	C	550	-	-	X	-
5	H3P	F	552	-	-	X	-

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 24074 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate dehydrogenase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	495	Total	C	N	O	S	0	0	0
			3866	2443	678	726	19			
1	B	495	Total	C	N	O	S	0	0	0
			3866	2443	678	726	19			
1	C	495	Total	C	N	O	S	0	0	0
			3866	2443	678	726	19			
1	D	495	Total	C	N	O	S	0	0	0
			3866	2443	678	726	19			
1	E	495	Total	C	N	O	S	0	0	0
			3866	2443	678	726	19			
1	F	495	Total	C	N	O	S	0	0	0
			3866	2443	678	726	19			

- Molecule 2 is GLUTAMIC ACID (three-letter code: GLU) (formula: C₅H₉NO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			9	5	1	3		
2	B	1	Total	C	N	O	0	0
			9	5	1	3		
2	C	1	Total	C	N	O	0	0
			9	5	1	3		
2	D	1	Total	C	N	O	0	0
			9	5	1	3		
2	E	1	Total	C	N	O	0	0
			9	5	1	3		
2	F	1	Total	C	N	O	0	0
			9	5	1	3		

- Molecule 3 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: C₂₁H₃₀N₇O₁₇P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 48	C 21	N 7	O 17	P 3	0	0
3	B	1	Total 48	C 21	N 7	O 17	P 3	0	0
3	C	1	Total 48	C 21	N 7	O 17	P 3	0	0
3	D	1	Total 48	C 21	N 7	O 17	P 3	0	0
3	E	1	Total 48	C 21	N 7	O 17	P 3	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	F	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 4 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
4	B	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
4	C	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
4	D	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
4	E	1	Total	C	N	O	P	0	0
			32	10	5	14	3		
4	F	1	Total	C	N	O	P	0	0
			32	10	5	14	3		

- Molecule 5 is 2,2'-methanediylbis(3,4,6-trichlorophenol) (three-letter code: H3P) (formula: $C_{13}H_6Cl_6O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total 21	C 13	Cl 6	O 2	0	0
5	B	1	Total 21	C 13	Cl 6	O 2	0	0
5	C	1	Total 21	C 13	Cl 6	O 2	0	0
5	C	1	Total 21	C 13	Cl 6	O 2	0	0
5	D	1	Total 21	C 13	Cl 6	O 2	0	0
5	F	1	Total 21	C 13	Cl 6	O 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	34	Total O 34 34	0	0
6	B	34	Total O 34 34	0	0
6	C	49	Total O 49 49	0	0
6	D	45	Total O 45 45	0	0
6	E	35	Total O 35 35	0	0
6	F	21	Total O 21 21	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamate dehydrogenase




- Molecule 1: Glutamate dehydrogenase






• Molecule 1: Glutamate dehydrogenase

Chain C: 63% 32% ..



• Molecule 1: Glutamate dehydrogenase


Chain D: 59% 36% ..



• Molecule 1: Glutamate dehydrogenase


Chain E: 61% 34% ..





• Molecule 1: Glutamate dehydrogenase

Chain F: 62% 33%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	123.37Å 101.58Å 166.87Å 90.00° 102.34° 90.00°	Depositor
Resolution (Å)	50.00 – 3.00 48.49 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-3.00) 94.2 (48.49-3.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.37 (at 3.01Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.240 , 0.266 0.328 , 0.346	Depositor DCC
R_{free} test set	6506 reflections (8.07%)	wwPDB-VP
Wilson B-factor (Å ²)	76.5	Xtriage
Anisotropy	0.544	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 37.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.86	EDS
Total number of atoms	24074	wwPDB-VP
Average B, all atoms (Å ²)	91.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: H3P, GTP, NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/3948	0.71	1/5328 (0.0%)
1	B	0.43	0/3948	0.84	4/5328 (0.1%)
1	C	0.44	0/3948	0.71	1/5328 (0.0%)
1	D	0.44	0/3948	0.69	0/5328
1	E	0.46	0/3948	0.70	0/5328
1	F	0.46	0/3948	0.71	0/5328
All	All	0.45	0/23688	0.73	6/31968 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	E	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	419	ARG	NE-CZ-NH1	-23.43	108.59	120.30
1	B	419	ARG	NE-CZ-NH2	23.09	131.85	120.30
1	B	419	ARG	CD-NE-CZ	11.18	139.25	123.60
1	C	40	GLN	N-CA-C	-5.19	96.99	111.00
1	A	40	GLN	N-CA-C	-5.12	97.18	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	493	TYR	Sidechain
1	E	493	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3866	0	3832	139	1
1	B	3866	0	3832	131	0
1	C	3866	0	3832	139	1
1	D	3866	0	3832	150	0
1	E	3866	0	3832	141	0
1	F	3866	0	3832	149	0
2	A	9	0	5	4	0
2	B	9	0	5	3	0
2	C	9	0	5	5	0
2	D	9	0	5	2	0
2	E	9	0	5	1	0
2	F	9	0	5	3	0
3	A	48	0	26	3	0
3	B	48	0	26	4	0
3	C	48	0	26	7	0
3	D	48	0	26	8	0
3	E	48	0	26	5	0
3	F	48	0	26	5	0
4	A	32	0	12	3	0
4	B	32	0	10	1	0
4	C	32	0	12	0	0
4	D	32	0	11	1	0
4	E	32	0	11	2	0
4	F	32	0	11	2	0
5	A	21	0	4	4	0
5	B	21	0	4	0	0
5	C	42	0	9	3	0
5	D	21	0	4	0	0
5	F	21	0	5	6	0
6	A	34	0	0	4	0
6	B	34	0	0	3	0
6	C	49	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	D	45	0	0	5	0
6	E	35	0	0	5	0
6	F	21	0	0	3	0
All	All	24074	0	23271	799	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 799 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:258:HIS:HD2	1:A:261:ARG:HH11	1.28	0.82
1:B:258:HIS:HD2	1:B:261:ARG:HH11	1.29	0.81
1:F:258:HIS:HD2	1:F:261:ARG:HH11	1.28	0.80
1:B:336:ALA:HB3	1:B:337:PRO:HD3	1.64	0.80
1:E:261:ARG:HD2	4:E:553:GTP:O6	1.82	0.79

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:363:ARG:CG	1:C:38:GLU:OE2[2_657]	2.06	0.14

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	493/501 (98%)	444 (90%)	39 (8%)	10 (2%)	7	34
1	B	493/501 (98%)	442 (90%)	43 (9%)	8 (2%)	9	40
1	C	493/501 (98%)	441 (90%)	44 (9%)	8 (2%)	9	40

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	493/501 (98%)	441 (90%)	45 (9%)	7 (1%)	11	43
1	E	493/501 (98%)	435 (88%)	48 (10%)	10 (2%)	7	34
1	F	493/501 (98%)	444 (90%)	42 (8%)	7 (1%)	11	43
All	All	2958/3006 (98%)	2647 (90%)	261 (9%)	50 (2%)	9	39

5 of 50 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	39	GLU
1	A	474	GLY
1	A	496	ALA
1	B	39	GLU
1	B	474	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	412/417 (99%)	380 (92%)	32 (8%)	12	42
1	B	412/417 (99%)	385 (93%)	27 (7%)	16	49
1	C	412/417 (99%)	383 (93%)	29 (7%)	15	47
1	D	412/417 (99%)	383 (93%)	29 (7%)	15	47
1	E	412/417 (99%)	382 (93%)	30 (7%)	14	44
1	F	412/417 (99%)	386 (94%)	26 (6%)	18	51
All	All	2472/2502 (99%)	2299 (93%)	173 (7%)	14	47

5 of 173 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	394	TYR
1	E	410	LEU
1	E	35	ARG
1	E	277	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	F	35	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 89 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	406	ASN
1	E	388	ASN
1	D	484	ASN
1	E	250	GLN
1	F	56	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLU	D	550	-	7,8,9	0.91	0	4,9,11	1.07	0
4	GTP	C	553	-	29,34,34	1.90	7 (24%)	35,54,54	2.14	7 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLU	C	550	-	7,8,9	0.98	0	4,9,11	0.77	0
5	H3P	B	552	-	22,22,22	0.50	0	27,33,33	1.18	2 (7%)
5	H3P	C	554	-	22,22,22	0.38	0	27,33,33	0.79	0
4	GTP	F	553	-	29,34,34	1.76	7 (24%)	35,54,54	5.39	14 (40%)
3	NDP	A	551	-	47,52,52	2.21	9 (19%)	61,80,80	2.48	20 (32%)
4	GTP	E	553	-	29,34,34	2.15	6 (20%)	35,54,54	4.74	13 (37%)
3	NDP	D	551	-	47,52,52	2.00	10 (21%)	61,80,80	2.68	21 (34%)
2	GLU	F	550	-	7,8,9	1.16	0	4,9,11	1.20	0
3	NDP	B	551	-	47,52,52	2.07	8 (17%)	61,80,80	2.60	22 (36%)
4	GTP	D	553	-	29,34,34	2.90	8 (27%)	35,54,54	5.73	17 (48%)
3	NDP	F	551	-	47,52,52	1.76	9 (19%)	61,80,80	2.58	20 (32%)
5	H3P	A	552	-	22,22,22	0.36	0	27,33,33	0.82	0
2	GLU	A	550	-	7,8,9	0.86	0	4,9,11	0.91	0
3	NDP	E	551	-	47,52,52	1.83	9 (19%)	61,80,80	2.52	20 (32%)
5	H3P	F	552	-	22,22,22	0.57	0	27,33,33	0.81	0
4	GTP	B	553	-	29,34,34	2.44	6 (20%)	35,54,54	5.53	18 (51%)
5	H3P	D	552	-	22,22,22	0.69	0	27,33,33	1.21	2 (7%)
2	GLU	B	550	-	7,8,9	1.02	0	4,9,11	1.11	0
2	GLU	E	550	-	7,8,9	0.99	0	4,9,11	0.90	0
3	NDP	C	551	-	47,52,52	1.86	7 (14%)	61,80,80	2.62	26 (42%)
5	H3P	C	552	-	22,22,22	0.38	0	27,33,33	1.31	2 (7%)
4	GTP	A	553	-	29,34,34	1.90	6 (20%)	35,54,54	1.47	7 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLU	D	550	-	-	0/6/7/9	-
4	GTP	C	553	-	-	0/18/38/38	0/3/3/3
2	GLU	C	550	-	-	3/6/7/9	-
5	H3P	B	552	-	-	0/4/4/4	0/2/2/2
5	H3P	C	554	-	-	1/4/4/4	0/2/2/2
4	GTP	F	553	-	-	2/18/38/38	0/3/3/3
3	NDP	A	551	-	-	4/30/77/77	0/5/5/5
4	GTP	E	553	-	-	3/18/38/38	0/3/3/3
3	NDP	D	551	-	-	3/30/77/77	0/5/5/5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLU	F	550	-	-	3/6/7/9	-
3	NDP	B	551	-	-	2/30/77/77	0/5/5/5
4	GTP	D	553	-	-	3/18/38/38	0/3/3/3
3	NDP	F	551	-	-	3/30/77/77	0/5/5/5
5	H3P	A	552	-	-	1/4/4/4	0/2/2/2
2	GLU	A	550	-	-	2/6/7/9	-
3	NDP	E	551	-	-	2/30/77/77	0/5/5/5
5	H3P	F	552	-	-	1/4/4/4	0/2/2/2
4	GTP	B	553	-	-	2/18/38/38	0/3/3/3
5	H3P	D	552	-	-	0/4/4/4	0/2/2/2
2	GLU	B	550	-	-	1/6/7/9	-
2	GLU	E	550	-	-	1/6/7/9	-
3	NDP	C	551	-	-	1/30/77/77	0/5/5/5
5	H3P	C	552	-	-	0/4/4/4	0/2/2/2
4	GTP	A	553	-	-	0/18/38/38	0/3/3/3

The worst 5 of 92 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	553	GTP	PB-O3A	10.15	1.70	1.59
4	B	553	GTP	PB-O3B	9.97	1.70	1.59
3	A	551	NDP	PN-O3	-8.76	1.50	1.59
3	C	551	NDP	PN-O3	-8.35	1.50	1.59
3	D	551	NDP	PN-O3	-7.99	1.50	1.59

The worst 5 of 211 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	553	GTP	C4'-O4'-C1'	-26.68	85.49	109.92
4	F	553	GTP	C4'-O4'-C1'	-25.29	86.76	109.92
4	D	553	GTP	C4'-O4'-C1'	-23.40	88.50	109.92
4	E	553	GTP	C4'-O4'-C1'	-20.04	91.57	109.92
4	D	553	GTP	O4'-C1'-N9	12.17	124.89	108.75

There are no chirality outliers.

5 of 38 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	550	GLU	O-C-CA-CB
2	B	550	GLU	O-C-CA-CB

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
2	E	550	GLU	O-C-CA-CB
2	F	550	GLU	O-C-CA-CB
2	F	550	GLU	N-CA-CB-CG



There are no ring outliers.


21 monomers are involved in 70 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	550	GLU	2	0
2	C	550	GLU	5	0
5	C	554	H3P	2	0
4	F	553	GTP	2	0
3	A	551	NDP	3	0
4	E	553	GTP	2	0
3	D	551	NDP	8	0
2	F	550	GLU	3	0
3	B	551	NDP	4	0
4	D	553	GTP	1	0
3	F	551	NDP	5	0
5	A	552	H3P	4	0
2	A	550	GLU	4	0
3	E	551	NDP	5	0
5	F	552	H3P	6	0
4	B	553	GTP	1	0
2	B	550	GLU	3	0
2	E	550	GLU	1	0
3	C	551	NDP	7	0
5	C	552	H3P	1	0
4	A	553	GTP	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




































5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)




Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around NDP B 551:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around NDP C 551:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NDP D 551:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NDP E 551:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NDP F 551:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GTP A 553:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP B 553:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around GTP C 553:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP D 553:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP E 553:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around GTP F 553:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around H3P A 552:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around H3P B 552:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around H3P C 552:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around H3P C 554:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around H3P D 552:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around H3P F 552:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.