



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 24, 2024 – 01:35 PM EDT

PDB ID : 6EXO
Title : Crystal Structure of Rhodesain in complex with a Macrolactam Inhibitor
Authors : Dietzel, U.; Kisker, C.
Deposited on : 2017-11-08
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	215	 % 95% .
1	B	215	 % 92% 8%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3658 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cysteine protease.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	215	Total	C	N	O	S	0	5	0
			1626	1014	268	330	14			
1	B	215	Total	C	N	O	S	0	11	0
			1657	1033	271	340	13			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	172	ALA	SER	engineered mutation	UNP Q95PM0
B	172	ALA	SER	engineered mutation	UNP Q95PM0

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total 4	C 2	O 2	0	0
2	A	1	Total 4	C 2	O 2	0	0
2	A	1	Total 4	C 2	O 2	0	0
2	A	1	Total 4	C 2	O 2	0	0
2	A	1	Total 4	C 2	O 2	0	0
2	A	1	Total 4	C 2	O 2	0	0
2	A	1	Total 4	C 2	O 2	0	0
2	A	1	Total 4	C 2	O 2	0	0
2	A	1	Total 4	C 2	O 2	0	0
2	A	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is (3 {S},14 {E})-19-chloranyl- {N}-(1-cyanocyclopropyl)-5-oxidanylidene-12,17-dioxa-4-azatricyclo[16.2.2.0^{6,11}]docosa-1(21),6(11),7,9,14,18(22),19-heptaene-3-carboxamide (three-letter code: C3E) (formula: C₂₄H₂₂ClN₃O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	0	0
			32	24	1	3	4		
3	B	1	Total	C	Cl	N	O	0	0
			32	24	1	3	4		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	110	Total	O	0	0
			110	110		
4	B	109	Total	O	0	0
			109	109		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cysteine protease



- Molecule 1: Cysteine protease



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	31.72Å 49.81Å 67.47Å 103.86° 98.98° 99.99°	Depositor
Resolution (Å)	47.18 – 1.90 44.22 – 1.90	Depositor EDS
% Data completeness (in resolution range)	91.4 (47.18-1.90) 91.4 (44.22-1.90)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.195 , 0.251 0.205 , 0.261	Depositor DCC
R_{free} test set	1414 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	28.5	Xtriage
Anisotropy	0.273	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 29.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3658	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: C3E, CSX, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/1670	0.56	0/2277
1	B	0.41	0/1718	0.56	0/2344
All	All	0.40	0/3388	0.56	0/4621

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1626	0	1518	10	0
1	B	1657	0	1548	10	0
2	A	44	0	66	2	0
2	B	48	0	72	1	0
3	A	32	0	0	1	0
3	B	32	0	0	2	0
4	A	110	0	0	1	0
4	B	109	0	0	0	0
All	All	3658	0	3204	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:103[A]:MET:CE	1:A:103[A]:MET:HA	2.12	0.79
1:A:103[A]:MET:HA	1:A:103[A]:MET:HE2	1.77	0.64
1:A:103[A]:MET:HA	1:A:103[A]:MET:HE3	1.78	0.64
1:B:31:ILE:HG23	1:B:48:LEU:HB2	1.84	0.60
1:B:25:CSX:SG	1:B:162:HIS:ND1	2.75	0.59
1:A:73:ASN:HD21	2:A:306:EDO:H22	1.69	0.57
1:B:8:ARG:HD3	1:B:193:TYR:CZ	2.46	0.50
1:B:66:GLY:N	3:B:303:C3E:O	2.39	0.49
1:B:7:TRP:CE2	1:B:133:GLY:HA2	2.48	0.49
1:A:25:CSX:SG	3:A:304:C3E:NAA	2.87	0.47
1:B:129:LEU:HD22	1:B:166[B]:LEU:HD13	1.97	0.47
1:A:103[A]:MET:CE	1:A:103[A]:MET:CA	2.87	0.46
1:A:73:ASN:HD21	2:A:306:EDO:C2	2.29	0.45
1:B:166[A]:LEU:HD11	1:B:178:TRP:CE3	2.52	0.44
1:B:95[A]:ASN:HD21	2:B:312:EDO:C2	2.30	0.44
1:A:8:ARG:NH1	4:A:405:HOH:O	2.50	0.44
3:B:303:C3E:CAM	3:B:303:C3E:CAF	2.98	0.42
1:A:36:GLY:HA3	1:A:134:PRO:HD2	2.01	0.42
1:B:166[B]:LEU:HD21	1:B:178:TRP:CE3	2.54	0.42
1:B:165:LEU:HB3	1:B:181:LYS:HB3	2.01	0.41
1:A:165:LEU:HB3	1:A:181:LYS:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	217/215 (101%)	209 (96%)	8 (4%)	0	100	100
1	B	223/215 (104%)	213 (96%)	10 (4%)	0	100	100
All	All	440/430 (102%)	422 (96%)	18 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	174/169 (103%)	170 (98%)	4 (2%)	50	45
1	B	180/169 (106%)	176 (98%)	4 (2%)	52	47
All	All	354/338 (105%)	346 (98%)	8 (2%)	59	45

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	89	TYR
1	A	103[A]	MET
1	A	103[B]	MET
1	A	104	ASN
1	B	89	TYR
1	B	146	ASP
1	B	161[A]	ASP
1	B	161[B]	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSX	B	25	1	3,6,7	0.73	0	1,6,8	2.51	1 (100%)
1	CSX	A	25	1	3,6,7	0.55	0	1,6,8	1.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSX	B	25	1	-	0/1/5/7	-
1	CSX	A	25	1	-	0/1/5/7	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	25	CSX	CA-CB-SG	2.51	118.83	113.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	25	CSX	1	0
1	A	25	CSX	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

25 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	301	-	3,3,3	0.59	0	2,2,2	0.08	0
3	C3E	B	303	-	34,35,35	1.89	8 (23%)	45,49,49	2.14	6 (13%)
2	EDO	A	309	-	3,3,3	0.41	0	2,2,2	0.35	0
2	EDO	A	307	-	3,3,3	0.45	0	2,2,2	0.42	0
2	EDO	B	304	-	3,3,3	0.45	0	2,2,2	0.25	0
2	EDO	B	311	-	3,3,3	0.47	0	2,2,2	0.21	0
2	EDO	B	301	-	3,3,3	0.52	0	2,2,2	0.22	0
2	EDO	B	307	-	3,3,3	0.48	0	2,2,2	0.29	0
2	EDO	B	313	-	3,3,3	0.46	0	2,2,2	0.26	0
2	EDO	A	305	-	3,3,3	0.54	0	2,2,2	0.08	0
3	C3E	A	304	-	34,35,35	2.00	8 (23%)	45,49,49	0.91	2 (4%)
2	EDO	B	305	-	3,3,3	0.36	0	2,2,2	0.42	0
2	EDO	A	308	-	3,3,3	0.49	0	2,2,2	0.37	0
2	EDO	B	306	-	3,3,3	0.48	0	2,2,2	0.32	0
2	EDO	B	308	-	3,3,3	0.52	0	2,2,2	0.28	0
2	EDO	A	306	-	3,3,3	0.48	0	2,2,2	0.32	0
2	EDO	A	302	-	3,3,3	0.43	0	2,2,2	0.28	0
2	EDO	A	311	-	3,3,3	0.56	0	2,2,2	0.06	0
2	EDO	B	310	-	3,3,3	0.45	0	2,2,2	0.31	0
2	EDO	B	302	-	3,3,3	0.51	0	2,2,2	0.12	0
2	EDO	B	309	-	3,3,3	0.44	0	2,2,2	0.30	0
2	EDO	A	312	-	3,3,3	0.49	0	2,2,2	0.24	0
2	EDO	B	312	-	3,3,3	0.48	0	2,2,2	0.25	0
2	EDO	A	310	-	3,3,3	0.51	0	2,2,2	0.26	0
2	EDO	A	303	-	3,3,3	0.48	0	2,2,2	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	301	-	-	1/1/1/1	-
3	C3E	B	303	-	-	11/30/37/37	0/3/4/4

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	309	-	-	1/1/1/1	-
2	EDO	A	307	-	-	1/1/1/1	-
2	EDO	B	304	-	-	1/1/1/1	-
2	EDO	B	311	-	-	1/1/1/1	-
2	EDO	B	301	-	-	1/1/1/1	-
2	EDO	B	307	-	-	0/1/1/1	-
2	EDO	B	313	-	-	1/1/1/1	-
2	EDO	A	305	-	-	1/1/1/1	-
3	C3E	A	304	-	-	9/30/37/37	0/3/4/4
2	EDO	B	305	-	-	1/1/1/1	-
2	EDO	A	308	-	-	1/1/1/1	-
2	EDO	B	306	-	-	0/1/1/1	-
2	EDO	B	308	-	-	1/1/1/1	-
2	EDO	A	306	-	-	0/1/1/1	-
2	EDO	A	302	-	-	0/1/1/1	-
2	EDO	A	311	-	-	1/1/1/1	-
2	EDO	B	310	-	-	1/1/1/1	-
2	EDO	B	302	-	-	1/1/1/1	-
2	EDO	B	309	-	-	0/1/1/1	-
2	EDO	A	312	-	-	1/1/1/1	-
2	EDO	B	312	-	-	1/1/1/1	-
2	EDO	A	310	-	-	1/1/1/1	-
2	EDO	A	303	-	-	1/1/1/1	-

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	304	C3E	CAE-NAA	5.36	1.22	1.14
3	A	304	C3E	CBD-CAX	-4.89	1.40	1.50
3	B	303	C3E	CAE-NAA	4.70	1.21	1.14
3	A	304	C3E	CA-C	-4.65	1.40	1.52
3	B	303	C3E	CBD-CAX	-4.45	1.41	1.50
3	B	303	C3E	CA-C	-4.34	1.41	1.52
3	A	304	C3E	CB-CG	-4.33	1.40	1.51
3	B	303	C3E	CB-CG	-3.71	1.42	1.51
3	B	303	C3E	CAX-N	3.55	1.41	1.34
3	A	304	C3E	CAX-N	3.35	1.41	1.34
3	A	304	C3E	CAZ-CL1	2.53	1.79	1.73
3	B	303	C3E	CAP-CAG	-2.42	1.39	1.49
3	B	303	C3E	CAQ-CBF	2.10	1.54	1.49
3	B	303	C3E	CAO-CAF	-2.08	1.40	1.49
3	A	304	C3E	CAP-CAG	-2.06	1.40	1.49

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	304	C3E	CAO-CAF	-2.06	1.40	1.49

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	303	C3E	CAQ-CBF-NAU	9.55	127.34	117.25
3	B	303	C3E	CAR-CBF-NAU	-7.55	109.28	117.25
3	B	303	C3E	CA-C-NAU	4.39	126.33	116.77
3	A	304	C3E	CBC-CBD-CAX	-3.15	120.42	126.16
3	B	303	C3E	O-C-NAU	-3.09	116.53	123.14
3	B	303	C3E	C-CA-N	-2.74	103.69	111.16
3	A	304	C3E	CAR-CBF-NAU	2.17	119.55	117.25
3	B	303	C3E	CA-N-CAX	2.08	126.69	121.60

There are no chirality outliers.

All (38) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	303	C3E	CA-C-NAU-CBF
3	B	303	C3E	O-C-NAU-CBF
3	B	303	C3E	CAE-CBF-NAU-C
3	B	303	C3E	CAQ-CBF-NAU-C
3	B	303	C3E	CAR-CBF-NAU-C
3	A	304	C3E	N-CAX-CBD-CBC
3	A	304	C3E	OAB-CAX-CBD-CBC
3	A	304	C3E	CAZ-CBB-OAV-CAO
3	B	303	C3E	OAB-CAX-CBD-CBC
3	B	303	C3E	N-CAX-CBD-CBC
3	B	303	C3E	CAZ-CBB-OAV-CAO
2	A	301	EDO	O1-C1-C2-O2
2	A	308	EDO	O1-C1-C2-O2
2	A	309	EDO	O1-C1-C2-O2
2	A	310	EDO	O1-C1-C2-O2
2	B	301	EDO	O1-C1-C2-O2
2	B	308	EDO	O1-C1-C2-O2
2	B	310	EDO	O1-C1-C2-O2
2	B	311	EDO	O1-C1-C2-O2
2	B	312	EDO	O1-C1-C2-O2
3	B	303	C3E	CAM-CBB-OAV-CAO
3	A	304	C3E	CAF-CAO-OAV-CBB
2	A	307	EDO	O1-C1-C2-O2
3	B	303	C3E	N-CA-CB-CG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	303	EDO	O1-C1-C2-O2
2	B	304	EDO	O1-C1-C2-O2
2	B	313	EDO	O1-C1-C2-O2
2	A	311	EDO	O1-C1-C2-O2
3	A	304	C3E	CAF-CAG-CAP-OAW
2	A	312	EDO	O1-C1-C2-O2
3	B	303	C3E	C-CA-CB-CG
2	A	305	EDO	O1-C1-C2-O2
3	A	304	C3E	CAM-CBB-OAV-CAO
2	B	302	EDO	O1-C1-C2-O2
3	A	304	C3E	N-CAX-CBD-CAK
3	A	304	C3E	CAJ-CBC-OAW-CAP
2	B	305	EDO	O1-C1-C2-O2
3	A	304	C3E	OAB-CAX-CBD-CAK


There are no ring outliers.


4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	303	C3E	2	0
3	A	304	C3E	1	0
2	A	306	EDO	2	0
2	B	312	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand C3E B 303





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	214/215 (99%)	-0.04	3 (1%) 75 77	20, 29, 44, 56	0
1	B	214/215 (99%)	0.00	3 (1%) 75 77	21, 32, 45, 56	0
All	All	428/430 (99%)	-0.02	6 (1%) 75 77	20, 31, 45, 56	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	152	LEU	3.2
1	B	3	ALA	3.2
1	A	174	ASN	3.1
1	A	215	GLY	2.3
1	B	1	ALA	2.2
1	B	131	GLU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSX	A	25	7/8	0.96	0.11	25,26,32,34	0
1	CSX	B	25	7/8	0.96	0.12	24,26,33,34	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	B	311	4/4	0.69	0.30	46,48,49,51	0
2	EDO	A	305	4/4	0.72	0.37	54,55,55,55	0
2	EDO	B	301	4/4	0.73	0.27	62,62,63,63	0
2	EDO	A	303	4/4	0.76	0.19	48,49,49,50	0
2	EDO	B	308	4/4	0.77	0.25	52,53,54,54	0
2	EDO	B	304	4/4	0.78	0.17	53,56,56,56	0
2	EDO	B	312	4/4	0.79	0.18	58,59,59,62	0
3	C3E	B	303	32/32	0.79	0.18	48,53,55,58	0
2	EDO	A	306	4/4	0.81	0.21	52,54,54,55	0
2	EDO	A	308	4/4	0.81	0.20	42,43,44,45	0
2	EDO	A	311	4/4	0.81	0.36	48,52,52,52	0
2	EDO	A	312	4/4	0.83	0.17	55,57,57,58	0
2	EDO	A	302	4/4	0.84	0.22	49,49,50,50	0
3	C3E	A	304	32/32	0.84	0.14	41,47,53,53	0
2	EDO	B	307	4/4	0.84	0.16	54,54,54,55	0
2	EDO	A	310	4/4	0.85	0.23	34,39,39,42	0
2	EDO	B	306	4/4	0.85	0.17	53,54,54,55	0
2	EDO	A	301	4/4	0.86	0.10	43,46,47,47	0
2	EDO	B	313	4/4	0.86	0.21	46,49,49,50	0
2	EDO	B	302	4/4	0.87	0.13	42,43,43,45	0
2	EDO	B	305	4/4	0.87	0.24	44,44,46,46	0
2	EDO	B	309	4/4	0.89	0.15	55,57,58,59	0
2	EDO	A	307	4/4	0.90	0.12	36,40,40,42	0
2	EDO	A	309	4/4	0.90	0.23	52,52,52,53	0
2	EDO	B	310	4/4	0.93	0.10	52,52,53,53	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around C3E B 303:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around C3E A 304:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.