



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 29, 2024 – 12:33 AM EDT

PDB ID : 1EYS  
Title : CRYSTAL STRUCTURE OF PHOTOSYNTHETIC REACTION CENTER FROM A THERMOPHILIC BACTERIUM, THERMOCHROMATIUM TEPIDUM  
Authors : Nogi, T.; Fathir, I.; Kobayashi, M.; Nozawa, T.; Miki, K.  
Deposited on : 2000-05-08  
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	C	382	
2	L	280	
3	M	324	
4	H	259	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
8	BPH	L	606	X	-	-	-

## 2 Entry composition

There are 14 unique types of molecules in this entry. The entry contains 10044 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOTOSYNTHETIC REACTION CENTER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	310	Total	C	N	O	S	0	0	0
			2402	1514	421	451	16			

- Molecule 2 is a protein called PHOTOSYNTHETIC REACTION CENTER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	280	Total	C	N	O	S	0	0	0
			2233	1501	361	361	10			


- Molecule 3 is a protein called PHOTOSYNTHETIC REACTION CENTER.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	M	318	Total	C	N	O	S	0	0	0
			2537	1705	413	409	10			

- Molecule 4 is a protein called PHOTOSYNTHETIC REACTION CENTER.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	238	Total	C	N	O	S	0	0	0
			1837	1187	309	336	5			

- Molecule 5 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
5	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
5	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
5	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0

- Molecule 6 is 2-O-octyl-beta-D-glucopyranose (three-letter code: BGL) (formula: C<sub>14</sub>H<sub>28</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	L	1	Total 20	C 14	O 6	0	0
6	L	1	Total 20	C 14	O 6	0	0
6	L	1	Total 20	C 14	O 6	0	0
6	M	1	Total 20	C 14	O 6	0	0
6	M	1	Total 20	C 14	O 6	0	0
6	M	1	Total 20	C 14	O 6	0	0

- Molecule 7 is BACTERIOCHLOROPHYLL A (three-letter code: BCL) (formula:  $C_{55}H_{74}MgN_4O_6$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	L	1	Total 66	C 55	Mg 1	N 4	O 6	0	0
7	L	1	Total 66	C 55	Mg 1	N 4	O 6	0	0
7	M	1	Total 66	C 55	Mg 1	N 4	O 6	0	0
7	M	1	Total 66	C 55	Mg 1	N 4	O 6	0	0

- Molecule 8 is BACTERIOPHEOPHYTIN A (three-letter code: BPH) (formula:  $\text{C}_{55}\text{H}_{76}\text{N}_4\text{O}_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	L	1	Total	C	N	O	0	0
			65	55	4	6		
8	M	1	Total	C	N	O	0	0
			65	55	4	6		

- Molecule 9 is LAURYL DIMETHYLAMINE-N-OXIDE (three-letter code: LDA) (formula:  $C_{14}H_{31}NO$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	L	1	Total	C	N	O	0	0
			16	14	1	1		

- Molecule 10 is FE (III) ION (three-letter code: FE) (formula: Fe).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	M	1	Total	Fe	0	0
			1	1		

- Molecule 11 is MENAQUINONE 8 (three-letter code: MQ8) (formula: C<sub>51</sub>H<sub>72</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	M	1	Total	C	O	0	0
			53	51	2		

- Molecule 12 is SPIRILLOXANTHIN (three-letter code: CRT) (formula: C<sub>42</sub>H<sub>60</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
12	M	1	Total	C	O	0	0
			44	42	2		

- Molecule 13 is DI-PALMITOYL-3-SN-PHOSPHATIDYLETHANOLAMINE (three-letter code: PEF) (formula:  $C_{37}H_{74}NO_8P$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
13	H	1	Total	C	N	O	P	0	0
			47	37	1	8	1		

- Molecule 14 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
14	C	94	Total	O	0	0
			94	94		
14	L	37	Total	O	0	0
			37	37		
14	M	35	Total	O	0	0
			35	35		
14	H	22	Total	O	0	0
			22	22		

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


#### • Molecule 1: PHOTOSYNTHETIC REACTION CENTER




#### • Molecule 2: PHOTOSYNTHETIC REACTION CENTER




#### • Molecule 3: PHOTOSYNTHETIC REACTION CENTER





● Molecule 4: PHOTOSYNTHETIC REACTION CENTER



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	133.30Å 196.60Å 84.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.20	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-2.20)	Depositor
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, $R_{free}$	0.231 , 0.287	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	10044	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEF, HEM, LDA, MQ8, FE, CRT, BCL, BGL, BPH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	C	0.53	1/2471 (0.0%)	0.72	2/3374 (0.1%)
2	L	0.50	0/2320	0.65	0/3170
3	M	0.49	0/2637	0.67	1/3610 (0.0%)
4	H	0.47	0/1890	0.76	1/2576 (0.0%)
All	All	0.50	1/9318 (0.0%)	0.69	4/12730 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	225	CYS	CB-SG	-5.56	1.72	1.81

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	159	LEU	CA-CB-CG	7.48	132.51	115.30
1	C	230	ASN	N-CA-C	-6.83	92.55	111.00
3	M	47	ILE	N-CA-C	-5.57	95.95	111.00
1	C	3	GLY	N-CA-C	5.11	125.88	113.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2402	0	2323	74	0
2	L	2233	0	2195	85	0
3	M	2537	0	2511	123	0
4	H	1837	0	1831	82	0
5	C	172	0	120	5	0
6	L	60	0	84	10	0
6	M	60	0	84	1	0
7	L	132	0	148	14	0
7	M	132	0	148	16	0
8	L	65	0	75	6	0
8	M	65	0	75	4	0
9	L	16	0	31	2	0
10	M	1	0	0	0	0
11	M	53	0	72	1	0
12	M	44	0	60	2	0
13	H	47	0	73	7	0
14	C	94	0	0	0	0
14	H	22	0	0	1	0
14	L	37	0	0	0	0
14	M	35	0	0	1	0
All	All	10044	0	9830	364	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 364 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:M:69:ILE:HD13	3:M:116:ILE:HG23	1.36	1.06
4:H:7:HIS:HB3	4:H:9:ILE:HG12	1.40	1.00
4:H:151:PRO:HA	4:H:154:MET:SD	2.07	0.94
3:M:33:PRO:HG3	3:M:49:PRO:HD3	1.52	0.92
2:L:86:ALA:H	2:L:95:GLN:HE22	1.06	0.90

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	308/382 (81%)	268 (87%)	31 (10%)	9 (3%)	3	2
2	L	278/280 (99%)	249 (90%)	27 (10%)	2 (1%)	19	19
3	M	316/324 (98%)	279 (88%)	31 (10%)	6 (2%)	6	4
4	H	234/259 (90%)	191 (82%)	33 (14%)	10 (4%)	2	1
All	All	1136/1245 (91%)	987 (87%)	122 (11%)	27 (2%)	5	3

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	2	GLU
1	C	4	PRO
4	H	140	LYS
4	H	142	PHE
4	H	146	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	259/300 (86%)	239 (92%)	20 (8%)	10	11
2	L	228/228 (100%)	212 (93%)	16 (7%)	12	14
3	M	254/258 (98%)	234 (92%)	20 (8%)	10	11
4	H	195/211 (92%)	166 (85%)	29 (15%)	2	2
All	All	936/997 (94%)	851 (91%)	85 (9%)	7	8

5 of 85 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	H	14	ILE
4	H	140	LYS
4	H	61	LEU
4	H	116	SER
4	H	159	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
3	M	198	ASN
4	H	122	HIS
4	H	218	HIS
4	H	189	ASN
1	C	253	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 1 is monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	BGL	L	703	-	20,20,20	0.93	1 (5%)	24,25,25	2.02	7 (29%)
11	MQ8	M	608	-	54,54,54	2.47	17 (31%)	67,69,69	2.79	20 (29%)
7	BCL	L	604	2	64,74,74	2.47	12 (18%)	74,115,115	2.32	23 (31%)
6	BGL	M	702	-	20,20,20	1.09	1 (5%)	24,25,25	2.39	8 (33%)
8	BPH	M	605	-	51,70,70	2.04	10 (19%)	52,101,101	2.37	14 (26%)
8	BPH	L	606	-	51,70,70	1.88	10 (19%)	52,101,101	2.54	16 (30%)
5	HEM	C	611	1	42,50,50	1.69	9 (21%)	46,82,82	1.02	1 (2%)
6	BGL	L	704	-	20,20,20	1.25	2 (10%)	24,25,25	2.39	9 (37%)
13	PEF	H	708	-	46,46,46	2.19	7 (15%)	49,51,51	1.40	6 (12%)
9	LDA	L	707	-	13,15,15	2.46	2 (15%)	14,17,17	0.64	0
5	HEM	C	609	1	42,50,50	1.74	12 (28%)	46,82,82	1.14	2 (4%)
7	BCL	L	602	2	64,74,74	2.31	10 (15%)	74,115,115	2.22	20 (27%)
6	BGL	L	701	-	20,20,20	1.28	2 (10%)	24,25,25	1.95	8 (33%)
6	BGL	M	706	-	20,20,20	0.98	1 (5%)	24,25,25	2.04	6 (25%)
7	BCL	M	603	3	64,74,74	2.38	10 (15%)	74,115,115	2.20	22 (29%)
7	BCL	M	601	3	64,74,74	2.53	11 (17%)	74,115,115	2.33	22 (29%)
6	BGL	M	705	-	20,20,20	1.30	2 (10%)	24,25,25	2.01	6 (25%)
5	HEM	C	610	1	42,50,50	1.76	10 (23%)	46,82,82	1.20	4 (8%)
12	CRT	M	613	-	43,43,43	2.39	13 (30%)	48,54,54	2.61	5 (10%)
5	HEM	C	612	1	42,50,50	1.52	6 (14%)	46,82,82	0.86	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	BGL	L	703	-	-	6/11/31/31	0/1/1/1
11	MQ8	M	608	-	-	12/47/67/67	0/2/2/2
7	BCL	L	604	2	-	8/37/137/137	-
6	BGL	M	702	-	-	7/11/31/31	0/1/1/1
8	BPH	M	605	-	-	11/37/105/105	0/5/6/6
8	BPH	L	606	-	1/1/18/22	15/37/105/105	0/5/6/6
5	HEM	C	611	1	-	4/12/54/54	-
6	BGL	L	704	-	-	5/11/31/31	0/1/1/1
13	PEF	H	708	-	-	34/50/50/50	-

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	LDA	L	707	-	-	4/13/13/13	-
5	HEM	C	609	1	-	4/12/54/54	-
7	BCL	L	602	2	-	10/37/137/137	-
6	BGL	L	701	-	-	7/11/31/31	0/1/1/1
6	BGL	M	706	-	-	5/11/31/31	0/1/1/1
7	BCL	M	603	3	-	8/37/137/137	-
7	BCL	M	601	3	-	21/37/137/137	-
6	BGL	M	705	-	-	4/11/31/31	0/1/1/1
5	HEM	C	610	1	-	6/12/54/54	-
12	CRT	M	613	-	-	5/51/51/51	-
5	HEM	C	612	1	-	4/12/54/54	-

The worst 5 of 148 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	M	601	BCL	C3C-C4C	-12.13	1.36	1.51
7	L	604	BCL	C3C-C4C	-12.06	1.36	1.51
7	M	603	BCL	C3C-C4C	-10.55	1.38	1.51
7	L	602	BCL	C3C-C4C	-10.39	1.38	1.51
7	M	601	BCL	CHC-C1C	10.23	1.42	1.33

The worst 5 of 200 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	M	613	CRT	C37-C36-C35	-11.54	108.70	124.91
8	L	606	BPH	O2D-CGD-CBD	10.96	122.99	110.95
12	M	613	CRT	C4-C5-C6	-9.81	111.13	124.91
7	L	604	BCL	C4A-NA-C1A	8.88	110.73	106.68
7	M	601	BCL	C4A-NA-C1A	8.61	110.61	106.68

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
8	L	606	BPH	C8

5 of 180 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	611	HEM	C2B-C3B-CAB-CBB

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Res	Type	Atoms
6	L	703	BGL	C2'-C1'-O2-C2
6	M	705	BGL	C2'-C1'-O2-C2
7	L	602	BCL	C2C-C3C-CAC-CBC
7	L	602	BCL	C4C-C3C-CAC-CBC



There are no ring outliers.


17 monomers are involved in 65 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	L	703	BGL	4	0
11	M	608	MQ8	1	0
7	L	604	BCL	10	0
6	M	702	BGL	1	0
8	M	605	BPH	4	0
8	L	606	BPH	6	0
6	L	704	BGL	3	0
13	H	708	PEF	7	0
9	L	707	LDA	2	0
5	C	609	HEM	1	0
7	L	602	BCL	7	0
6	L	701	BGL	3	0
7	M	603	BCL	6	0
7	M	601	BCL	10	0
5	C	610	HEM	2	0
12	M	613	CRT	2	0
5	C	612	HEM	2	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



























## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.