Full wwPDB EM Validation Report

Jun 25, 2025 – 05:51 am BST

PDB ID	:	9F10 / pdb_00009f10
EMDB ID	:	EMD-50120
Title	:	CryoEM structure of the F plasmid relaxosome with TraI in its TE mode, without accessory protein TraM. Derived from the ss-27_+8ds+9_+143-R_ deltaTraM Locally-refined 2.94 A Map.
Authors	:	Williams, S.M.; Waksman, G.
Deposited on	:	2024-04-17
Resolution	:	2.94 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

EMDB validation analysis	:	0.0.1.dev118
MolProbity	:	4-5-2 with Phenix2.0rc1
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ	:	1.9.13
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.44

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	170	37% 19% 44%
2	B	135	41% 21% 37%
3	C	99	83% 13%
4	D	94	91% 9%
5	E	131	75% 11% 15%
5	F	131	73% 18% 8%
5	G	131	14% 68% 15% 17%
6	H	1763	27% 5% 68%

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 12286 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called T-strand DNA (96-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	96	Total	C	N	O	P	0	0
			1982	950	367	569	96		

- Molecule 2 is a DNA chain called R-strand DNA (85-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	85	Total	C	N	O	P	0	0
			1739	841	293	520	85		

- Molecule 3 is a protein called Integration host factor subunit alpha.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	95	Total	C	N	O	S	0	0
			764	478	142	143	1		

- Molecule 4 is a protein called Integration host factor subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	94	Total	C	N	O	S	0	0
			749	467	139	140	3		

- Molecule 5 is a protein called Relaxosome protein TraY.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	112	Total	C	N	O	S	0	0
			906	572	169	161	4		
5	F	120	Total	C	N	O	S	0	0
			972	613	180	175	4		
5	G	109	Total	C	N	O	S	0	0
			867	546	160	158	3		

- Molecule 6 is a protein called Multifunctional conjugation protein TraI.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	H	564	Total	C	N	O	S	0	0
			4306	2656	788	842	20		

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	-6	MET	-	initiating methionine	UNP P14565
H	-5	GLU	-	expression tag	UNP P14565
H	-4	PHE	-	expression tag	UNP P14565
H	-3	GLU	-	expression tag	UNP P14565
H	-2	LEU	-	expression tag	UNP P14565
H	-1	GLY	-	expression tag	UNP P14565
H	0	THR	-	expression tag	UNP P14565
H	16	PHE	TYR	engineered mutation	UNP P14565

- Molecule 7 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
7	H	1	Total	Mg	0
			1	1	

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: T-strand DNA (96-MER)

Chain A:





• Molecule 2: R-strand DNA (85-MER)

Chain B:





• Molecule 3: Integration host factor subunit alpha

Chain C:

• Molecule 4: Integration host factor subunit beta

Chain D:





• Molecule 5: Relaxosome protein TraY

Chain E:



• Molecule 5: Relaxosome protein TraY

Chain F:



• Molecule 5: Relaxosome protein TraY

Chain G:



• Molecule 6: Multifunctional conjugation protein TraI

Chain H:




R566 I570 D575 R576 R579 A584 K592 E595
Q600 E606 I609 I614 K619 V623 L626 P627 R642 D648 R651 M654 Y669 R673 L683 L704 M710 A713 E716 R719 L727 R728 V729 L735 V749 P750 G751 E771 E776 V792 A796 L827
T834

4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	330708	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/Å^2$)	50.09	Depositor
Minimum defocus (nm)	900	Depositor
Maximum defocus (nm)	2700	Depositor
Magnification	105000	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	5.031	Depositor
Minimum map value	-3.401	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.056	Depositor
Recommended contour level	0.4	Depositor
Map size (Å)	462.0, 462.0, 462.0	wwPDB
Map dimensions	560, 560, 560	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.825, 0.825, 0.825	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| > 5$	RMSZ	$\#\|Z\| > 5$
1	A	0.33	0/2228	0.49	0/3439
2	B	0.33	0/1946	0.51	0/3002
3	C	0.23	0/775	0.29	0/1036
4	D	0.20	0/763	0.23	0/1021
5	E	0.20	0/921	0.28	0/1232
5	F	0.20	0/989	0.29	0/1326
5	G	0.16	0/882	0.28	0/1186
6	H	0.18	0/4376	0.32	0/5920
All	All	0.24	0/12880	0.38	0/18162

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1982	0	1088	35	0
2	B	1739	0	974	22	0
3	C	764	0	779	11	0
4	D	749	0	756	9	0
5	E	906	0	928	9	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	F	972	0	988	17	0
5	G	867	0	853	13	0
6	H	4306	0	4209	61	0
7	H	1	0	0	0	0
All	All	12286	0	10575	151	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (151) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:E:22:VAL:HG11	5:E:102:ILE:HG12	1.63	0.80
6:H:651:ARG:H	6:H:654:MET:HE2	1.52	0.74
6:H:60:LEU:HD21	6:H:165:VAL:HG11	1.69	0.74
2:B:33:DT:H2'	2:B:34:DT:H71	1.72	0.71
5:G:47:ASP:OD2	5:G:51:ARG:NH1	2.23	0.70
6:H:710:MET:HE1	6:H:771:GLU:HA	1.75	0.69
6:H:122:SER:OG	6:H:176:SER:OG	2.10	0.67
5:G:57:ARG:HG2	5:G:59:GLY:H	1.61	0.65
6:H:713:ALA:H	6:H:716:GLU:HG3	1.62	0.64
2:B:38:DT:O2	4:D:62:ARG:NH1	2.30	0.64
1:A:45:DA:N3	3:C:63:ARG:NH1	2.46	0.64
5:F:108:PHE:HA	5:F:118:ARG:HG2	1.79	0.64
5:F:13:MET:HE3	5:F:71:THR:HG23	1.79	0.64
2:B:46:DA:OP2	3:C:76:ARG:NH1	2.24	0.63
2:B:11:DC:H2'	2:B:12:DA:C8	2.33	0.63
3:C:35:ARG:HG2	4:D:86:LEU:HD13	1.81	0.63
6:H:112:PHE:O	6:H:116:GLN:NE2	2.32	0.62
5:F:57:ARG:NH2	5:F:96:GLU:OE1	2.33	0.61
4:D:42:ARG:NH1	4:D:44:GLU:OE1	2.23	0.61
1:A:50:DA:OP1	4:D:42:ARG:NH2	2.35	0.60
6:H:575:ASP:O	6:H:579:ARG:NH1	2.35	0.60
6:H:719:ARG:NH1	6:H:771:GLU:OE2	2.36	0.58
5:G:48:HIS:NE2	5:G:103:ASP:OD2	2.33	0.57
6:H:729:VAL:HG11	6:H:735:LEU:HD21	1.86	0.57
6:H:68:ARG:NH1	6:H:70:GLN:OE1	2.38	0.56
6:H:584:ALA:HA	6:H:614:ILE:HD13	1.86	0.56
1:A:85:DC:H2"	1:A:84:DA:C8	2.41	0.55
3:C:22:ASP:OD1	6:H:673:ARG:NH1	2.40	0.55
2:B:81:DT:H2"	2:B:82:DT:H71	1.89	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:74:DT:H2"	1:A:73:DA:C8	2.42	0.54
1:A:5:DG:H2"	1:A:4:DG:O5'	2.07	0.53
5:F:57:ARG:NH1	5:F:99:ASP:OD2	2.34	0.53
5:F:39:PHE:HA	5:F:42:VAL:HG22	1.89	0.53
2:B:85:DG:O6	5:G:73:ARG:NH2	2.42	0.53
6:H:792:VAL:HG13	6:H:796:ALA:HB3	1.90	0.53
5:G:18:LEU:O	5:G:68:LYS:N	2.42	0.53
5:G:48:HIS:CE1	5:G:52:TYR:HD2	2.27	0.52
5:F:57:ARG:HH22	5:F:96:GLU:HA	1.74	0.52
1:A:88:DT:OP1	5:G:11:GLY:N	2.42	0.52
1:A:84:DA:C8	1:A:84:DA:H5'	2.45	0.52
1:A:58:DT:H6	1:A:58:DT:H5'	1.75	0.52
6:H:184:GLY:O	6:H:188:ASN:ND2	2.32	0.51
6:H:576:ARG:HH12	6:H:609:ILE:HD12	1.76	0.51
6:H:626:LEU:HB2	6:H:627:PRO:HD3	1.93	0.51
1:A:48:DT:H2'	1:A:47:DC:C6	2.46	0.51
2:B:86:DA:H2'	2:B:87:DT:C6	2.47	0.51
5:E:72:MET:HE3	5:E:74:LEU:HD11	1.94	0.50
1:A:2:DG:OP1	6:H:88:LYS:NZ	2.34	0.50
6:H:88:LYS:HB3	6:H:231:VAL:HG12	1.94	0.49
1:A:2:DG:H2"	1:A:1:DT:OP2	2.12	0.49
5:F:109:PRO:HD2	5:F:118:ARG:HB3	1.93	0.49
2:B:45:DT:H4'	2:B:46:DA:OP1	2.13	0.49
2:B:10:DG:H2"	2:B:11:DC:OP2	2.13	0.49
5:F:112:TYR:HB3	5:F:116:ILE:HB	1.95	0.48
6:H:600:GLN:HG2	6:H:776:GLU:HG3	1.94	0.48
6:H:619:LYS:HE3	6:H:626:LEU:HA	1.94	0.48
1:A:4:DG:C4	6:H:1:MET:HE2	2.48	0.48
6:H:2:MET:HE2	6:H:4:ILE:HG12	1.96	0.48
1:A:2:DG:N2	6:H:255:ASP:OD1	2.47	0.47
3:C:62:GLY:HA3	3:C:73:ILE:HD11	1.96	0.47
5:F:19:PRO:HB2	5:F:22:VAL:HG23	1.97	0.47
6:H:168:HIS:CG	6:H:169:ASN:H	2.32	0.47
5:F:107:LYS:O	5:F:118:ARG:NH1	2.47	0.47
6:H:669:TYR:HB3	6:H:683:LEU:HB3	1.97	0.47
1:A:41:DA:H8	1:A:41:DA:H5'	1.80	0.47
5:G:75:ASP:OD1	5:G:78:THR:OG1	2.22	0.46
6:H:108:GLN:HG3	6:H:207:GLN:HE22	1.80	0.46
6:H:136:THR:O	6:H:138:ASN:ND2	2.46	0.46
6:H:592:LYS:HE3	6:H:623:VAL:HG22	1.97	0.46
5:F:82:LEU:HD21	5:F:97:ALA:HB2	1.97	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:43:GLN:O	4:D:1:MET:N	2.48	0.46
6:H:2:MET:HE3	6:H:82:LEU:HD13	1.98	0.46
1:A:30:DT:H1'	1:A:29:DC:H5'	1.98	0.46
6:H:126:MET:HB2	6:H:131:SER:HB3	1.98	0.46
1:A:43:DT:H2"	1:A:42:DT:H5"	1.97	0.46
1:A:6:DG:O4'	6:H:193:GLN:NE2	2.49	0.46
5:G:82:LEU:HD21	5:G:97:ALA:HB2	1.98	0.46
5:F:14:VAL:HG22	5:F:93:LYS:HE3	1.98	0.45
2:B:61:DA:H5'	2:B:61:DA:C8	2.51	0.45
5:E:27:ILE:HG12	5:E:37:ARG:HD3	1.98	0.45
3:C:3:LEU:HB2	4:D:36:THR:HG21	1.99	0.45
5:G:48:HIS:O	5:G:48:HIS:ND1	2.49	0.45
1:A:46:DT:H5'	1:A:46:DT:C6	2.52	0.45
2:B:53:DT:H2"	2:B:54:DA:N7	2.32	0.45
1:A:41:DA:H5'	1:A:41:DA:C8	2.51	0.45
2:B:41:DT:H2"	2:B:42:DA:C8	2.52	0.45
6:H:113:ALA:HB2	6:H:200:TYR:HB2	1.98	0.45
2:B:93:DA:H2"	2:B:94:DA:C8	2.52	0.44
6:H:104:ASP:HA	6:H:107:ASN:ND2	2.32	0.44
6:H:178:ASP:N	6:H:178:ASP:OD1	2.50	0.44
3:C:5:LYS:NZ	3:C:28:GLU:OE1	2.43	0.44
5:E:30:SER:OG	5:E:35:ARG:O	2.30	0.44
1:A:8:DT:H2'	6:H:190:TYR:CD2	2.53	0.44
1:A:2:DG:H3'	6:H:237:ARG:HB3	2.00	0.44
6:H:654:MET:HE3	6:H:704:LEU:HD11	2.00	0.44
1:A:29:DC:H2"	1:A:28:DA:C8	2.53	0.44
1:A:46:DT:H6	1:A:46:DT:H2'	1.66	0.44
6:H:141:MET:HG2	6:H:162:VAL:HG22	1.99	0.44
5:E:21:ASP:O	5:E:24:SER:OG	2.31	0.43
5:E:80:GLN:HG3	6:H:642:ARG:NH2	2.33	0.43
2:B:73:DT:H4'	5:F:3:ARG:HD2	1.99	0.43
6:H:116:GLN:HG2	6:H:199:LEU:HD13	1.99	0.43
5:G:92:CYS:SG	5:G:93:LYS:N	2.92	0.43
5:G:74:LEU:HB3	5:G:78:THR:OG1	2.18	0.43
6:H:92:MET:HE3	6:H:92:MET:HB2	1.95	0.43
5:G:19:PRO:HB2	5:G:22:VAL:HG23	2.01	0.43
6:H:33:TRP:CZ2	6:H:47:VAL:HG13	2.53	0.43
2:B:28:DT:H2"	2:B:29:DG:C8	2.54	0.43
5:E:14:VAL:HG22	5:E:93:LYS:HE3	2.00	0.43
5:F:22:VAL:HG11	5:F:102:ILE:HG12	2.01	0.43
1:A:90:DA:H2"	1:A:89:DA:C8	2.54	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:H:111:ASP:O	6:H:115:ARG:HG3	2.19	0.42
6:H:229:VAL:HG23	6:H:291:TYR:CZ	2.54	0.42
2:B:30:DA:C8	2:B:30:DA:H5'	2.55	0.42
6:H:178:ASP:CG	6:H:183:THR:H	2.27	0.42
5:E:8:SER:HB2	6:H:648:ASP:HB3	2.00	0.42
6:H:154:PRO:HD3	6:H:277:TRP:CE2	2.54	0.42
2:B:91:DT:H2"	2:B:92:DA:H8	1.85	0.42
6:H:576:ARG:NH1	6:H:606:GLU:OE1	2.52	0.42
1:A:5:DG:N7	6:H:201:ARG:NH2	2.58	0.42
2:B:12:DA:H2"	2:B:13:DA:C8	2.55	0.42
5:F:118:ARG:H	5:F:118:ARG:HG3	1.35	0.42
2:B:91:DT:H2"	2:B:92:DA:C8	2.55	0.41
6:H:113:ALA:O	6:H:117:VAL:HG23	2.20	0.41
1:A:87:DA:C8	1:A:86:DT:H72	2.56	0.41
1:A:44:DT:H2"	1:A:43:DT:C6	2.55	0.41
1:A:26:DC:H2"	1:A:25:DA:C8	2.55	0.41
3:C:86:LYS:HB3	3:C:86:LYS:HE2	1.86	0.41
6:H:120:LEU:HD13	6:H:188:ASN:HB3	2.03	0.41
2:B:47:DG:H2"	2:B:48:DA:H5"	2.03	0.41
1:A:78:DA:H5'	1:A:78:DA:H8	1.86	0.41
1:A:-1:DG:O6	6:H:3:SER:HB2	2.21	0.41
2:B:47:DG:H4'	2:B:48:DA:OP1	2.20	0.41
5:F:48:HIS:NE2	5:F:103:ASP:OD2	2.34	0.41
1:A:41:DA:C2	2:B:42:DA:C2	3.09	0.41
1:A:2:DG:O6	6:H:254:ARG:NH2	2.51	0.41
6:H:89:SER:HA	6:H:92:MET:HG2	2.02	0.41
5:E:21:ASP:OD1	5:E:21:ASP:N	2.53	0.41
6:H:70:GLN:HB3	6:H:75:ARG:HG3	2.02	0.41
6:H:92:MET:SD	6:H:229:VAL:HG12	2.61	0.40
6:H:203:LYS:O	6:H:207:GLN:HG3	2.21	0.40
5:F:32:ARG:NE	5:F:110:ASP:OD1	2.36	0.40
6:H:136:THR:HB	6:H:138:ASN:HD21	1.86	0.40
1:A:30:DT:H2"	1:A:29:DC:C6	2.57	0.40
3:C:74:THR:HG22	3:C:76:ARG:HH21	1.85	0.40
3:C:81:PHE:HB3	4:D:80:PHE:HB3	2.04	0.40
6:H:570:ILE:HD11	6:H:827:LEU:HD12	2.03	0.40
1:A:37:DA:C2	4:D:64:PRO:HG2	2.57	0.40
6:H:727:LEU:HD22	6:H:749:VAL:HG21	2.03	0.40
1:A:37:DA:N1	4:D:64:PRO:HG2	2.36	0.40
6:H:203:LYS:HG3	6:H:207:GLN:HE21	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	C	93/99 (94%)	92 (99%)	1 (1%)	0	100	100
4	D	92/94 (98%)	92 (100%)	0	0	100	100
5	E	108/131 (82%)	106 (98%)	2 (2%)	0	100	100
5	F	118/131 (90%)	115 (98%)	3 (2%)	0	100	100
5	G	107/131 (82%)	104 (97%)	3 (3%)	0	100	100
6	H	556/1763 (32%)	536 (96%)	20 (4%)	0	100	100
All	All	1074/2349 (46%)	1045 (97%)	29 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	82/86 (95%)	82 (100%)	0	100	100
4	D	79/79 (100%)	79 (100%)	0	100	100
5	E	97/116 (84%)	97 (100%)	0	100	100
5	F	104/116 (90%)	103 (99%)	1 (1%)	73	84
5	G	90/116 (78%)	90 (100%)	0	100	100
6	H	454/1428 (32%)	454 (100%)	0	100	100
All	All	906/1941 (47%)	905 (100%)	1 (0%)	92	96

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	F	118	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
6	H	116	GLN
6	H	679	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Map visualisation

This section contains visualisations of the EMDB entry EMD-50120. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections

6.1.1 Primary map



X



Y



Z

6.1.2 Raw map



X



Y



Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map

X Index: 280



Y Index: 280

Z Index: 280

6.2.2 Raw map

X Index: 280



Y Index: 280

Z Index: 280

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map

X Index: 281



Y Index: 296

Z Index: 308

6.3.2 Raw map



X Index: 282



Y Index: 296

Z Index: 308

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



X



Y



Z

6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views (i)

6.5.1 Primary map

X Y Z

The images above show the 3D surface view of the map at the recommended contour level 0.4. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map

X Y Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation (i)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency

A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

6.6.1 emd_50120_msk_1.map (i)



X



Y



Z

7 Map analysis

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate



The volume at the recommended contour level is 114 nm^3; this corresponds to an approximate mass of 103 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.340 Å^{-1}

8 Fourier-Shell correlation

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC



*Reported resolution corresponds to spatial frequency of 0.340 Å^{-1}

8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.94	-	-
Author-provided FSC curve	2.94	3.15	2.98
Unmasked-calculated*	4.13	7.70	4.32

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.13 differs from the reported value 2.94 by more than 10 %

9 Map-model fit

This section contains information regarding the fit between EMDB map EMD-50120 and PDB model 9F10. Per-residue inclusion information can be found in section 3 on page 5.

9.1 Map-model overlay



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.4 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model (i)



X Y Z

The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.4).

9.4 Atom inclusion (i)



At the recommended contour level, 91% of all backbone atoms, 87% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.4) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8720	0.5140
A	0.9400	0.5760
B	0.9470	0.5780
C	0.9340	0.5710
D	0.9410	0.5880
E	0.9100	0.5550
F	0.8730	0.5220
G	0.6870	0.3530
H	0.8150	0.4600