



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 29, 2024 – 02:27 PM EDT

PDB ID : 1FBI  
Title : CRYSTAL STRUCTURE OF A CROSS-REACTION COMPLEX BETWEEN FAB F9.13.7 AND GUINEA-FOWL LYSOZYME  
Authors : Lescar, J.; Alzari, P.M.  
Deposited on : 1995-01-19  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : NOT EXECUTED  
EDS : NOT EXECUTED  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	L	214	46% 47% 7%
1	P	214	47% 46% 6%
2	H	221	44% 48% 6% •
2	Q	221	55% 40% 5%
3	X	129	49% 44% 6% •
3	Y	129	47% 46% 5% •

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8521 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called IGG1 F9.13.7 FAB (LIGHT CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	L	214	Total	C	N	O	S	0	0	0
			1641	1023	273	338	7			
1	P	214	Total	C	N	O	S	0	0	0
			1644	1022	271	344	7			

- Molecule 2 is a protein called IGG1 F9.13.7 FAB (HEAVY CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	221	Total	C	N	O	S	0	0	0
			1624	1034	257	326	7			
2	Q	221	Total	C	N	O	S	0	0	0
			1614	1035	253	319	7			

- Molecule 3 is a protein called GUINEA FOWL LYSOZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	X	129	Total	C	N	O	S	0	0	0
			1001	614	193	184	10			
3	Y	129	Total	C	N	O	S	0	0	0
			997	611	192	184	10			


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


#### • Molecule 1: IGG1 F9.13.7 FAB (LIGHT CHAIN)

Chain L: 




#### • Molecule 1: IGG1 F9.13.7 FAB (LIGHT CHAIN)


Chain P: 




#### • Molecule 2: IGG1 F9.13.7 FAB (HEAVY CHAIN)

Chain H: 






• Molecule 2: IGG1 F9.13.7 FAB (HEAVY CHAIN)



• Molecule 3: GUINEA FOWL LYSOZYME



• Molecule 3: GUINEA FOWL LYSOZYME



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.70Å 195.50Å 50.20Å 90.00° 108.50° 90.00°	Depositor
Resolution (Å)	7.00 – 3.00	Depositor
% Data completeness (in resolution range)	(Not available) (7.00-3.00)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.190 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	8521	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	L	0.50	0/1676	0.76	0/2277
1	P	0.44	0/1679	0.72	1/2283 (0.0%)
2	H	0.54	0/1671	0.82	1/2294 (0.0%)
2	Q	0.43	0/1661	0.66	0/2283
3	X	0.55	0/1022	0.86	1/1379 (0.1%)
3	Y	0.46	0/1018	0.72	0/1375
All	All	0.49	0/8727	0.75	3/11891 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	P	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	X	63	TRP	N-CA-C	5.84	126.77	111.00
1	P	78	LEU	CA-CB-CG	5.67	128.35	115.30
2	H	99	LEU	N-CA-C	-5.62	95.83	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	P	50	TYR	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	L	1641	0	1555	98	0
1	P	1644	0	1543	92	0
2	H	1624	0	1541	114	0
2	Q	1614	0	1530	82	0
3	X	1001	0	960	63	0
3	Y	997	0	949	62	0
All	All	8521	0	8078	481	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 29.

The worst 5 of 481 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:P:159:VAL:HG22	1:P:179:LEU:HD23	1.47	0.96
1:L:49:TYR:HB2	2:H:108:VAL:HG21	1.49	0.94
3:Y:78:ILE:HD12	3:Y:79:PRO:HD2	1.58	0.85
2:Q:14:PRO:HG3	2:Q:120:VAL:HG12	1.59	0.84
2:Q:219:ILE:O	2:Q:221:PRO:HD3	1.81	0.81

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	L	212/214 (99%)	167 (79%)	31 (15%)	14 (7%)	<b>1</b> <b>6</b>

*Continued on next page...*



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	P	212/214 (99%)	168 (79%)	35 (16%)	9 (4%)	3	16
2	H	219/221 (99%)	186 (85%)	20 (9%)	13 (6%)	1	9
2	Q	219/221 (99%)	179 (82%)	30 (14%)	10 (5%)	2	14
3	X	127/129 (98%)	97 (76%)	23 (18%)	7 (6%)	2	10
3	Y	127/129 (98%)	93 (73%)	26 (20%)	8 (6%)	1	7
All	All	1116/1128 (99%)	890 (80%)	165 (15%)	61 (6%)	2	10

5 of 61 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	L	15	LEU
1	L	80	GLN
1	L	110	ASP
1	L	171	SER
1	L	172	THR

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	L	185/192 (96%)	162 (88%)	23 (12%)	4	20
1	P	186/192 (97%)	171 (92%)	15 (8%)	11	40
2	H	179/190 (94%)	162 (90%)	17 (10%)	8	32
2	Q	174/190 (92%)	155 (89%)	19 (11%)	6	25
3	X	105/105 (100%)	94 (90%)	11 (10%)	7	27
3	Y	104/105 (99%)	93 (89%)	11 (11%)	6	26
All	All	933/974 (96%)	837 (90%)	96 (10%)	7	28

5 of 96 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	P	93	THR

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
2	Q	96	CYS
1	P	98	PHE
1	P	213	GLU
2	Q	130	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	P	37	GLN
2	Q	164	ASN
1	P	55	HIS
3	Y	27	ASN
2	Q	82	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.