



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 9, 2024 – 08:35 AM EST

PDB ID : 5FBH
Title : Crystal structure of the extracellular domain of human calcium sensing receptor with bound Gd3+
Authors : Zhang, T.; Zhang, C.; Miller, C.L.; Zou, J.; Moremen, K.W.; Brown, E.M.; Yang, J.J.; Hu, J.
Deposited on : 2015-12-14
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	3333 (2.70-2.70)
Clashscore	180529	3684 (2.70-2.70)
Ramachandran outliers	177936	3633 (2.70-2.70)
Sidechain outliers	177891	3633 (2.70-2.70)
RSRZ outliers	164620	3333 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	568	
1	B	568	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 7394 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Extracellular calcium-sensing receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	470	Total	C	N	O	S	0	0	0
			3628	2328	615	673	12			
1	A	472	Total	C	N	O	S	0	0	0
			3604	2312	607	673	12			

There are 92 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-26	MET	-	initiating methionine	UNP P41180
B	-25	ARG	-	expression tag	UNP P41180
B	-24	LEU	-	expression tag	UNP P41180
B	-23	LEU	-	expression tag	UNP P41180
B	-22	THR	-	expression tag	UNP P41180
B	-21	ALA	-	expression tag	UNP P41180
B	-20	LEU	-	expression tag	UNP P41180
B	-19	PHE	-	expression tag	UNP P41180
B	-18	ALA	-	expression tag	UNP P41180
B	-17	TYR	-	expression tag	UNP P41180
B	-16	PHE	-	expression tag	UNP P41180
B	-15	ILE	-	expression tag	UNP P41180
B	-14	VAL	-	expression tag	UNP P41180
B	-13	ALA	-	expression tag	UNP P41180
B	-12	LEU	-	expression tag	UNP P41180
B	-11	ILE	-	expression tag	UNP P41180
B	-10	LEU	-	expression tag	UNP P41180
B	-9	ALA	-	expression tag	UNP P41180
B	-8	PHE	-	expression tag	UNP P41180
B	-7	SER	-	expression tag	UNP P41180
B	-6	VAL	-	expression tag	UNP P41180
B	-5	SER	-	expression tag	UNP P41180
B	-4	ALA	-	expression tag	UNP P41180
B	-3	LYS	-	expression tag	UNP P41180
B	-2	SER	-	expression tag	UNP P41180

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	MET	-	expression tag	UNP P41180
B	0	HIS	-	expression tag	UNP P41180
B	1	HIS	-	expression tag	UNP P41180
B	2	HIS	-	expression tag	UNP P41180
B	3	HIS	-	expression tag	UNP P41180
B	4	HIS	-	expression tag	UNP P41180
B	5	HIS	-	expression tag	UNP P41180
B	6	HIS	-	expression tag	UNP P41180
B	7	HIS	-	expression tag	UNP P41180
B	8	SER	-	expression tag	UNP P41180
B	9	ALA	-	expression tag	UNP P41180
B	10	TRP	-	expression tag	UNP P41180
B	11	SER	-	expression tag	UNP P41180
B	12	HIS	-	expression tag	UNP P41180
B	13	PRO	-	expression tag	UNP P41180
B	14	GLN	-	expression tag	UNP P41180
B	15	PHE	-	expression tag	UNP P41180
B	16	GLU	-	expression tag	UNP P41180
B	17	LYS	-	expression tag	UNP P41180
B	18	GLU	-	expression tag	UNP P41180
B	19	PHE	-	expression tag	UNP P41180
A	-26	MET	-	initiating methionine	UNP P41180
A	-25	ARG	-	expression tag	UNP P41180
A	-24	LEU	-	expression tag	UNP P41180
A	-23	LEU	-	expression tag	UNP P41180
A	-22	THR	-	expression tag	UNP P41180
A	-21	ALA	-	expression tag	UNP P41180
A	-20	LEU	-	expression tag	UNP P41180
A	-19	PHE	-	expression tag	UNP P41180
A	-18	ALA	-	expression tag	UNP P41180
A	-17	TYR	-	expression tag	UNP P41180
A	-16	PHE	-	expression tag	UNP P41180
A	-15	ILE	-	expression tag	UNP P41180
A	-14	VAL	-	expression tag	UNP P41180
A	-13	ALA	-	expression tag	UNP P41180
A	-12	LEU	-	expression tag	UNP P41180
A	-11	ILE	-	expression tag	UNP P41180
A	-10	LEU	-	expression tag	UNP P41180
A	-9	ALA	-	expression tag	UNP P41180
A	-8	PHE	-	expression tag	UNP P41180
A	-7	SER	-	expression tag	UNP P41180
A	-6	VAL	-	expression tag	UNP P41180

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	SER	-	expression tag	UNP P41180
A	-4	ALA	-	expression tag	UNP P41180
A	-3	LYS	-	expression tag	UNP P41180
A	-2	SER	-	expression tag	UNP P41180
A	-1	MET	-	expression tag	UNP P41180
A	0	HIS	-	expression tag	UNP P41180
A	1	HIS	-	expression tag	UNP P41180
A	2	HIS	-	expression tag	UNP P41180
A	3	HIS	-	expression tag	UNP P41180
A	4	HIS	-	expression tag	UNP P41180
A	5	HIS	-	expression tag	UNP P41180
A	6	HIS	-	expression tag	UNP P41180
A	7	HIS	-	expression tag	UNP P41180
A	8	SER	-	expression tag	UNP P41180
A	9	ALA	-	expression tag	UNP P41180
A	10	TRP	-	expression tag	UNP P41180
A	11	SER	-	expression tag	UNP P41180
A	12	HIS	-	expression tag	UNP P41180
A	13	PRO	-	expression tag	UNP P41180
A	14	GLN	-	expression tag	UNP P41180
A	15	PHE	-	expression tag	UNP P41180
A	16	GLU	-	expression tag	UNP P41180
A	17	LYS	-	expression tag	UNP P41180
A	18	GLU	-	expression tag	UNP P41180
A	19	PHE	-	expression tag	UNP P41180


- Molecule 2 is GADOLINIUM ION (three-letter code: GD3) (formula: Gd).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Gd 1 1	0	0
2	A	1	Total Gd 1 1	0	0

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	2	Total Mg 2 2	0	0
3	A	1	Total Mg 1 1	0	0

- Molecule 4 is BICARBONATE ION (three-letter code: BCT) (formula: CHO_3).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	1	3		
4	A	1	Total	C	O	0	0
			4	1	3		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Cl	0	0
			1	1		
5	A	1	Total	Cl	0	0
			1	1		

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $\text{C}_8\text{H}_{15}\text{NO}_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		
6	B	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		
6	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 7 is CYCLOMETHYLTRYPTOPHAN (three-letter code: TCR) (formula: $C_{12}H_{12}N_2O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	B	1	Total	C	N	O	0	0
			16	12	2	2		
7	A	1	Total	C	N	O	0	0
			16	12	2	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	11	Total	O	0	0
			11	11		
8	A	6	Total	O	0	0
			6	6		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Extracellular calcium-sensing receptor



- Molecule 1: Extracellular calcium-sensing receptor



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	172.11Å 83.11Å 94.47Å 90.00° 105.15° 90.00°	Depositor
Resolution (Å)	46.08 – 2.70 46.08 – 2.70	Depositor EDS
% Data completeness (in resolution range)	97.4 (46.08-2.70) 97.7 (46.08-2.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.91 (at 2.69Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.182 , 0.236 0.188 , 0.236	Depositor DCC
R_{free} test set	1731 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	68.7	Xtriage
Anisotropy	0.038	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 44.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7394	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TCR, NAG, BCT, CL, MG, GD3, CSO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	0/3676	0.73	1/5007 (0.0%)
1	B	0.65	0/3701	0.78	2/5031 (0.0%)
All	All	0.62	0/7377	0.76	3/10038 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	537	VAL	C-N-CA	6.91	151.00	122.00
1	B	455	VAL	CB-CA-C	-5.57	100.82	111.40
1	B	331	ARG	NE-CZ-NH1	5.01	122.80	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3604	0	3360	32	0
1	B	3628	0	3437	28	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	4	0	0	1	0
4	B	4	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	56	0	52	0	0
6	B	42	0	39	0	0
7	A	16	0	11	0	0
7	B	16	0	11	1	0
8	A	6	0	0	0	0
8	B	11	0	0	0	0
All	All	7394	0	6910	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 58 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:314:VAL:O	1:A:317:THR:HG23	1.80	0.81
1:B:314:VAL:O	1:B:317:THR:HG23	1.86	0.76
1:A:71:LEU:C	1:A:71:LEU:HD23	2.15	0.67
1:A:193:GLN:HE22	1:A:297:GLU:H	1.48	0.62
1:B:468:ASN:HA	1:B:478:THR:HG22	1.84	0.59

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	464/568 (82%)	436 (94%)	26 (6%)	2 (0%)	30	55
1	B	462/568 (81%)	440 (95%)	21 (4%)	1 (0%)	44	68

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	926/1136 (82%)	876 (95%)	47 (5%)	3 (0%)	37 61

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	118	ASN
1	A	62	ARG
1	A	89	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	362/487 (74%)	339 (94%)	23 (6%)	14 34
1	B	370/487 (76%)	346 (94%)	24 (6%)	14 34
All	All	732/974 (75%)	685 (94%)	47 (6%)	14 34

5 of 47 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	179	GLN
1	A	314	VAL
1	A	184	LEU
1	A	227	ARG
1	A	334	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	176	ASN
1	A	413	HIS
1	A	193	GLN
1	A	463	HIS
1	B	357	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CSO	A	236	1	3,6,7	0.70	0	1,6,8	0.94	0
1	CSO	A	482	1	3,6,7	0.64	0	1,6,8	0.75	0
1	CSO	B	236	1	3,6,7	0.56	0	1,6,8	0.80	0
1	CSO	B	482	1	3,6,7	0.40	0	1,6,8	0.04	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	236	1	-	0/1/5/7	-
1	CSO	A	482	1	-	0/1/5/7	-
1	CSO	B	236	1	-	0/1/5/7	-
1	CSO	B	482	1	-	0/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 7 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
6	NAG	A	606	1	14,14,15	0.60	0	17,19,21	1.60	3 (17%)
7	TCR	A	609	-	17,18,18	2.51	7 (41%)	16,26,26	2.39	3 (18%)
6	NAG	A	608	1	14,14,15	0.56	0	17,19,21	2.16	6 (35%)
6	NAG	A	605	1	14,14,15	0.58	0	17,19,21	1.25	1 (5%)
7	TCR	B	609	-	17,18,18	1.79	3 (17%)	16,26,26	1.96	4 (25%)
6	NAG	B	606	1	14,14,15	0.53	0	17,19,21	2.24	4 (23%)
4	BCT	A	603	-	3,3,3	0.58	0	2,3,3	3.02	2 (100%)
6	NAG	A	607	1	14,14,15	0.55	0	17,19,21	1.49	4 (23%)
4	BCT	B	604	-	3,3,3	0.63	0	2,3,3	0.65	0
6	NAG	B	608	1	14,14,15	0.52	0	17,19,21	2.08	7 (41%)
6	NAG	B	607	1	14,14,15	0.57	0	17,19,21	0.87	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	A	606	1	-	1/6/23/26	0/1/1/1
7	TCR	A	609	-	-	0/4/13/13	0/3/3/3
6	NAG	A	608	1	-	4/6/23/26	0/1/1/1
6	NAG	A	605	1	-	2/6/23/26	0/1/1/1
7	TCR	B	609	-	-	0/4/13/13	0/3/3/3
6	NAG	B	606	1	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	A	607	1	-	2/6/23/26	0/1/1/1
6	NAG	B	608	1	-	2/6/23/26	0/1/1/1
6	NAG	B	607	1	-	2/6/23/26	0/1/1/1

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	609	TCR	CG-CD1	5.45	1.49	1.39
7	A	609	TCR	CB-CG	-4.58	1.46	1.51
7	B	609	TCR	CG-CD1	3.28	1.45	1.39
7	A	609	TCR	CZ2-CE2	-2.95	1.37	1.41
7	A	609	TCR	CG-CD2	2.92	1.46	1.41

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	609	TCR	CB-CA-N	7.53	128.06	112.50
6	B	606	NAG	C1-O5-C5	7.50	122.24	112.19
6	A	608	NAG	C1-O5-C5	5.23	119.19	112.19
6	A	608	NAG	C8-C7-N2	4.00	122.75	116.12
6	A	605	NAG	C1-O5-C5	3.99	117.53	112.19

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	607	NAG	O5-C5-C6-O6
6	B	606	NAG	O5-C5-C6-O6
6	A	608	NAG	O5-C5-C6-O6
6	B	607	NAG	C4-C5-C6-O6
6	A	608	NAG	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	609	TCR	1	0
4	A	603	BCT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	470/568 (82%)	-0.11	9 (1%) 66 65	40, 67, 110, 144	0
1	B	468/568 (82%)	-0.26	7 (1%) 71 71	38, 60, 98, 126	0
All	All	938/1136 (82%)	-0.18	16 (1%) 69 68	38, 63, 105, 144	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	361	GLN	4.2
1	A	22	PRO	3.3
1	A	207	ASN	3.0
1	A	499	GLU	2.9
1	B	534	SER	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSO	B	236	7/8	0.88	0.14	53,57,78,78	0
1	CSO	B	482	7/8	0.94	0.07	54,65,80,80	0
1	CSO	A	236	7/8	0.96	0.06	48,52,65,77	0
1	CSO	A	482	7/8	0.96	0.06	58,60,67,82	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	NAG	A	606	14/15	0.64	0.13	93,124,141,144	0
6	NAG	A	608	14/15	0.74	0.13	96,106,121,134	0
3	MG	A	602	1/1	0.79	0.09	95,95,95,95	0
6	NAG	B	607	14/15	0.81	0.10	109,121,131,133	0
6	NAG	B	606	14/15	0.85	0.11	88,112,125,126	0
3	MG	B	603	1/1	0.85	0.10	77,77,77,77	0
6	NAG	A	607	14/15	0.86	0.09	89,109,114,114	0
5	CL	B	605	1/1	0.86	0.10	71,71,71,71	0
6	NAG	A	605	14/15	0.89	0.10	95,105,120,121	0
3	MG	B	602	1/1	0.89	0.15	62,62,62,62	0
6	NAG	B	608	14/15	0.93	0.08	59,62,79,81	0
5	CL	A	604	1/1	0.94	0.12	65,65,65,65	0
7	TCR	A	609	16/16	0.94	0.08	32,42,50,55	0
7	TCR	B	609	16/16	0.95	0.10	31,47,58,63	0
4	BCT	A	603	4/4	0.95	0.07	61,61,67,70	0
4	BCT	B	604	4/4	0.97	0.08	45,56,57,64	0
2	GD3	A	601	1/1	0.98	0.07	120,120,120,120	0
2	GD3	B	601	1/1	0.99	0.07	107,107,107,107	0

6.5 Other polymers [i](#)

There are no such residues in this entry.