



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 12, 2024 – 08:05 AM EDT

PDB ID : 5FFG
Title : Crystal structure of integrin alpha V beta 6 head
Authors : Dong, X.; Springer, T.A.
Deposited on : 2015-12-18
Resolution : 2.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	1763 (2.26-2.26)
Clashscore	180529	1919 (2.26-2.26)
Ramachandran outliers	177936	1884 (2.26-2.26)
Sidechain outliers	177891	1885 (2.26-2.26)
RSRZ outliers	164620	1763 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	601	<div> <div>3%</div> <div> <div></div> <div>85%</div> <div>13%</div> <div></div> </div> </div>
2	B	257	<div> <div>2%</div> <div> <div></div> <div>84%</div> <div>12%</div> <div></div> </div> </div>
3	C	2	<div> <div></div> <div>100%</div> </div>
4	D	3	<div> <div></div> <div>100%</div> </div>
4	F	3	<div> <div></div> <div> <div>67%</div> <div>33%</div> </div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	6	 17% 83%
6	G	8	 38% 62%

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 7099 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Integrin alpha-V.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	591	Total	C	N	O	S	0	0	0
			4580	2904	778	877	21			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	400	GLY	-	insertion	UNP P06756
A	401	CYS	MET	conflict	UNP P06756
A	597	THR	CYS	conflict	UNP P06756
A	599	GLY	-	expression tag	UNP P06756
A	600	LEU	-	expression tag	UNP P06756
A	601	GLU	-	expression tag	UNP P06756

- Molecule 2 is a protein called Integrin beta-6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	248	Total	C	N	O	S	0	0	0
			1941	1240	311	379	11			

There are 7 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	270	CYS	ILE	conflict	UNP P18564
B	362	HIS	-	expression tag	UNP P18564
B	363	HIS	-	expression tag	UNP P18564
B	364	HIS	-	expression tag	UNP P18564
B	365	HIS	-	expression tag	UNP P18564
B	366	HIS	-	expression tag	UNP P18564
B	367	HIS	-	expression tag	UNP P18564

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	3	Total	C	N	O	0	0	0
			39	22	2	15			
4	F	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
5	E	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 6 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
6	G	8	Total	C	N	O	0	0	0
			94	52	2	40			

- Molecule 7 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	4	Total	Ca	0	0
			4	4		
7	B	2	Total	Ca	0	0
			2	2		

- Molecule 8 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	O	0	0
			14	8	1	5		
8	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 9 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total	O	S		0	0
			5	4	1			

- Molecule 10 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
10	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 11 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	A	1	Total	C	O	0	0
			7	4	3		
11	B	1	Total	C	O	0	0
			7	4	3		


- Molecule 12 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	168	Total	O	0	0
			168	168		
12	B	73	Total	O	0	0
			73	73		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Integrin alpha-V



- Molecule 2: Integrin beta-6



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose





- Molecule 4: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:




- Molecule 5: alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain E:



- Molecule 6: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	215.59Å 77.56Å 59.07Å 90.00° 101.46° 90.00°	Depositor
Resolution (Å)	46.99 – 2.25 46.99 – 2.25	Depositor EDS
% Data completeness (in resolution range)	98.5 (46.99-2.25) 98.5 (46.99-2.25)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.71 (at 2.24Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, R_{free}	0.208 , 0.246 0.209 , 0.248	Depositor DCC
R_{free} test set	905 reflections (2.02%)	wwPDB-VP
Wilson B-factor (Å ²)	38.6	Xtriage
Anisotropy	0.463	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 55.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7099	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MAN, SO4, PEG, MES, CA, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.27	1/4683 (0.0%)	0.46	0/6338
2	B	0.26	0/1980	0.45	0/2685
All	All	0.27	1/6663 (0.0%)	0.46	0/9023

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	123	GLU	C-N	5.22	1.44	1.34

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4580	0	4428	48	0
2	B	1941	0	1917	20	0
3	C	28	0	25	0	0
4	D	39	0	34	0	0
4	F	39	0	34	1	0
5	E	72	0	61	1	0
6	G	94	0	79	0	0
7	A	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	2	0	0	0	0
8	A	14	0	13	1	0
8	B	14	0	13	0	0
9	A	5	0	0	1	0
10	A	12	0	12	0	0
11	A	7	0	10	2	0
11	B	7	0	10	0	0
12	A	168	0	0	8	0
12	B	73	0	0	2	0
All	All	7099	0	6636	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 67 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:141:ARG:NH1	12:B:2101:HOH:O	2.23	0.72
1:A:440:PRO:HB2	1:A:488:LEU:HD12	1.71	0.71
2:B:113:ASP:N	12:B:2102:HOH:O	2.27	0.68
1:A:397:ALA:HB2	1:A:403:PRO:HD3	1.76	0.67
1:A:488:LEU:HD21	1:A:492:LEU:HD21	1.78	0.66

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	587/601 (98%)	565 (96%)	21 (4%)	1 (0%)	44	51
2	B	246/257 (96%)	234 (95%)	12 (5%)	0	100	100
All	All	833/858 (97%)	799 (96%)	33 (4%)	1 (0%)	48	57

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	65	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	484/491 (99%)	476 (98%)	8 (2%)	56	66
2	B	221/230 (96%)	220 (100%)	1 (0%)	86	91
All	All	705/721 (98%)	696 (99%)	9 (1%)	65	74

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	488	LEU
2	B	243	ASN
1	A	155	CYS
1	A	178	TYR
1	A	275	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

22 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	C	1	1,3	14,14,15	0.52	0	17,19,21	0.52	0
3	NAG	C	2	3	14,14,15	0.26	0	17,19,21	0.56	0
4	NAG	D	1	1,4	14,14,15	0.23	0	17,19,21	0.51	0
4	NAG	D	2	4	14,14,15	0.22	0	17,19,21	0.39	0
4	BMA	D	3	4	11,11,12	0.60	0	15,15,17	0.75	0
5	NAG	E	1	1,5	14,14,15	0.29	0	17,19,21	0.39	0
5	NAG	E	2	5	14,14,15	0.34	0	17,19,21	0.40	0
5	BMA	E	3	5	11,11,12	0.77	1 (9%)	15,15,17	0.81	0
5	MAN	E	4	5	11,11,12	0.97	1 (9%)	15,15,17	0.84	0
5	MAN	E	5	5	11,11,12	0.74	0	15,15,17	0.93	1 (6%)
5	MAN	E	6	5	11,11,12	0.65	0	15,15,17	1.01	2 (13%)
4	NAG	F	1	1,4	14,14,15	0.32	0	17,19,21	0.45	0
4	NAG	F	2	4	14,14,15	0.27	0	17,19,21	0.44	0
4	BMA	F	3	4	11,11,12	0.61	0	15,15,17	0.80	0
6	NAG	G	1	1,6	14,14,15	0.28	0	17,19,21	0.57	0
6	NAG	G	2	6	14,14,15	0.34	0	17,19,21	0.51	0
6	BMA	G	3	6	11,11,12	0.81	0	15,15,17	0.67	0
6	MAN	G	4	6	11,11,12	2.06	4 (36%)	15,15,17	1.92	5 (33%)
6	MAN	G	5	6	11,11,12	0.88	1 (9%)	15,15,17	1.42	3 (20%)
6	MAN	G	6	6	11,11,12	0.84	0	15,15,17	1.03	1 (6%)
6	MAN	G	7	6	11,11,12	0.75	0	15,15,17	1.02	2 (13%)
6	MAN	G	8	6	11,11,12	0.63	0	15,15,17	1.05	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1
4	NAG	D	1	1,4	-	1/6/23/26	0/1/1/1
4	NAG	D	2	4	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	BMA	D	3	4	-	0/2/19/22	0/1/1/1
5	NAG	E	1	1,5	-	0/6/23/26	0/1/1/1
5	NAG	E	2	5	-	2/6/23/26	0/1/1/1
5	BMA	E	3	5	-	2/2/19/22	0/1/1/1
5	MAN	E	4	5	-	0/2/19/22	0/1/1/1
5	MAN	E	5	5	-	0/2/19/22	0/1/1/1
5	MAN	E	6	5	-	0/2/19/22	0/1/1/1
4	NAG	F	1	1,4	-	4/6/23/26	0/1/1/1
4	NAG	F	2	4	-	2/6/23/26	0/1/1/1
4	BMA	F	3	4	-	0/2/19/22	0/1/1/1
6	NAG	G	1	1,6	-	2/6/23/26	0/1/1/1
6	NAG	G	2	6	-	1/6/23/26	0/1/1/1
6	BMA	G	3	6	-	2/2/19/22	0/1/1/1
6	MAN	G	4	6	-	0/2/19/22	0/1/1/1
6	MAN	G	5	6	-	2/2/19/22	0/1/1/1
6	MAN	G	6	6	-	2/2/19/22	0/1/1/1
6	MAN	G	7	6	-	0/2/19/22	0/1/1/1
6	MAN	G	8	6	-	0/2/19/22	0/1/1/1

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	G	4	MAN	C2-C3	4.19	1.58	1.52
6	G	4	MAN	O2-C2	3.81	1.51	1.43
6	G	4	MAN	C1-C2	2.62	1.58	1.52
5	E	4	MAN	O5-C1	-2.26	1.39	1.43
6	G	5	MAN	C1-C2	2.24	1.57	1.52

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	G	4	MAN	C1-O5-C5	5.51	119.58	112.19
6	G	5	MAN	C1-O5-C5	3.05	116.27	112.19
6	G	8	MAN	C1-O5-C5	2.64	115.72	112.19
5	E	6	MAN	C1-O5-C5	2.59	115.65	112.19
6	G	5	MAN	C1-C2-C3	2.47	113.24	109.64

There are no chirality outliers.

5 of 24 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	G	3	BMA	O5-C5-C6-O6
3	C	2	NAG	O5-C5-C6-O6
4	F	2	NAG	O5-C5-C6-O6
3	C	2	NAG	C4-C5-C6-O6
4	F	2	NAG	C4-C5-C6-O6



There are no ring outliers.


2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	E	2	NAG	1	0
4	F	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	NAG	A	2027	1	14,14,15	0.32	0	17,19,21	0.39	0
10	MES	A	2029	-	12,12,12	2.30	1 (8%)	15,16,16	1.91	4 (26%)
9	SO4	A	2028	-	4,4,4	0.24	0	6,6,6	0.05	0
8	NAG	B	2003	2	14,14,15	0.24	0	17,19,21	0.44	0
11	PEG	B	2004	-	6,6,6	0.65	0	5,5,5	0.67	0
11	PEG	A	2030	-	6,6,6	0.64	0	5,5,5	0.67	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	NAG	A	2027	1	-	2/6/23/26	0/1/1/1
10	MES	A	2029	-	-	4/6/14/14	0/1/1/1
8	NAG	B	2003	2	-	0/6/23/26	0/1/1/1
11	PEG	B	2004	-	-	0/4/4/4	-
11	PEG	A	2030	-	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	A	2029	MES	C8-S	-7.69	1.66	1.77

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
10	A	2029	MES	C5-N4-C3	4.41	118.34	108.84
10	A	2029	MES	C6-C5-N4	-2.77	105.92	110.12
10	A	2029	MES	C7-N4-C5	2.22	117.15	111.24
10	A	2029	MES	O2S-S-C8	2.12	109.93	106.73

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
10	A	2029	MES	C7-C8-S-O3S
8	A	2027	NAG	C4-C5-C6-O6
8	A	2027	NAG	O5-C5-C6-O6
11	A	2030	PEG	O1-C1-C2-O2
11	A	2030	PEG	O2-C3-C4-O4

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	2027	NAG	1	0
9	A	2028	SO4	1	0
11	A	2030	PEG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	591/601 (98%)	0.30	16 (2%) 56 56	32, 68, 100, 158	0
2	B	248/257 (96%)	0.08	6 (2%) 59 60	29, 51, 89, 138	0
All	All	839/858 (97%)	0.24	22 (2%) 57 58	29, 63, 100, 158	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	488	LEU	6.1
1	A	566	TYR	3.9
1	A	469	LEU	3.8
1	A	582	PHE	3.6
2	B	280	SER	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	MAN	G	5	11/12	0.24	0.16	116,127,139,142	0
6	MAN	G	4	11/12	0.28	0.16	140,145,150,152	0
4	BMA	D	3	11/12	0.44	0.14	99,115,119,121	0
5	MAN	E	5	11/12	0.59	0.14	108,112,122,123	0
4	BMA	F	3	11/12	0.60	0.14	96,105,109,110	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	MAN	E	4	11/12	0.66	0.14	91,98,106,107	0
6	MAN	G	7	11/12	0.69	0.12	113,117,125,127	0
4	NAG	D	2	14/15	0.77	0.11	84,98,109,115	0
3	NAG	C	2	14/15	0.77	0.11	86,98,101,106	0
6	MAN	G	8	11/12	0.77	0.10	74,83,96,101	0
6	BMA	G	3	11/12	0.78	0.11	100,107,118,130	0
6	MAN	G	6	11/12	0.78	0.11	80,92,96,104	0
4	NAG	F	2	14/15	0.79	0.13	76,87,100,102	0
6	NAG	G	2	14/15	0.82	0.11	88,96,99,100	0
5	MAN	E	6	11/12	0.86	0.10	67,75,85,86	0
6	NAG	G	1	14/15	0.88	0.10	57,75,90,96	0
4	NAG	D	1	14/15	0.90	0.10	62,80,91,92	0
4	NAG	F	1	14/15	0.90	0.12	51,67,97,101	0
3	NAG	C	1	14/15	0.90	0.12	39,61,73,75	0
5	BMA	E	3	11/12	0.91	0.09	47,57,76,90	0
5	NAG	E	1	14/15	0.92	0.09	35,48,59,62	0
5	NAG	E	2	14/15	0.93	0.08	30,43,48,54	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain F:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
8	NAG	B	2003	14/15	0.59	0.14	78,94,99,103	0
8	NAG	A	2027	14/15	0.74	0.14	73,92,96,100	0
11	PEG	A	2030	7/7	0.81	0.14	63,71,80,83	0
11	PEG	B	2004	7/7	0.83	0.17	82,84,88,89	0
10	MES	A	2029	12/12	0.85	0.16	51,79,98,98	0
9	SO4	A	2028	5/5	0.89	0.12	61,65,68,68	5
7	CA	A	2003	1/1	0.96	0.07	55,55,55,55	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	CA	A	2001	1/1	0.97	0.05	49,49,49,49	0
7	CA	A	2004	1/1	0.97	0.06	61,61,61,61	0
7	CA	B	2002	1/1	0.97	0.08	53,53,53,53	1
7	CA	A	2002	1/1	0.98	0.04	58,58,58,58	0
7	CA	B	2001	1/1	0.99	0.02	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.