



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 12, 2024 – 12:15 AM EDT

PDB ID : 1FO6
Title : CRYSTAL STRUCTURE ANALYSIS OF N-CARBAMOYL-D-AMINO-ACID AMIDOHYDROLASE
Authors : Wang, W.-C.; Hsu, W.-H.; Chien, F.-T.; Chen, C.-Y.
Deposited on : 2000-08-25
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	304	 75% 18% 6% ..
1	B	304	 75% 21% . .
1	C	304	 73% 24% . .
1	D	304	 67% 27% 5% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 10288 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-CARBAMoyL-D-AMINO-ACID AMIDOHYDROLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	302	Total	C	N	O	S	0	0	0
			2391	1527	419	431	14			
1	B	302	Total	C	N	O	S	0	0	0
			2391	1527	419	431	14			
1	C	302	Total	C	N	O	S	0	0	0
			2391	1527	419	431	14			
1	D	302	Total	C	N	O	S	0	0	0
			2391	1527	419	431	14			

- Molecule 2 is XENON (three-letter code: XE) (formula: Xe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Xe	0	0
			1	1		
2	C	1	Total	Xe	0	0
			1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	200	Total	O	0	0
			200	200		
3	B	202	Total	O	0	0
			202	202		
3	C	169	Total	O	0	0
			169	169		
3	D	151	Total	O	0	0
			151	151		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: N-CARBAMoyL-D-AMINO-ACID AMIDOHYDROLASE

Chain A: 




• Molecule 1: N-CARBAMoyL-D-AMINO-ACID AMIDOHYDROLASE


Chain B: 




• Molecule 1: N-CARBAMoyL-D-AMINO-ACID AMIDOHYDROLASE

Chain C: 





● Molecule 1: N-CARBAMOYL-D-AMINO-ACID AMIDOHYDROLASE



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	70.23Å 67.53Å 137.48Å 90.00° 96.12° 90.00°	Depositor
Resolution (Å)	30.00 – 1.95	Depositor
% Data completeness (in resolution range)	94.9 (30.00-1.95)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.187 , 0.239	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	10288	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: XE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	0/2455	1.84	48/3327 (1.4%)
1	B	0.95	0/2455	1.71	33/3327 (1.0%)
1	C	0.87	0/2455	1.81	48/3327 (1.4%)
1	D	0.89	0/2455	2.15	54/3327 (1.6%)
All	All	0.92	0/9820	1.89	183/13308 (1.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	9
1	B	0	3
1	C	0	6
1	D	0	4
All	All	0	22

There are no bond length outliers.

The worst 5 of 183 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	38	ARG	CD-NE-CZ	39.53	178.94	123.60
1	D	56	ARG	CD-NE-CZ	36.00	174.00	123.60
1	C	38	ARG	NE-CZ-NH2	-28.27	106.17	120.30
1	D	107	ARG	NE-CZ-NH2	27.48	134.04	120.30
1	D	107	ARG	NE-CZ-NH1	-21.34	109.63	120.30

There are no chirality outliers.

5 of 22 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	117	LYS	Mainchain
1	A	118	SER	Mainchain
1	A	126	ARG	Mainchain
1	A	54	PHE	Mainchain
1	A	78	VAL	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2391	0	2335	47	0
1	B	2391	0	2335	55	0
1	C	2391	0	2335	41	0
1	D	2391	0	2335	75	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
3	A	200	0	0	2	0
3	B	202	0	0	2	0
3	C	169	0	0	4	0
3	D	151	0	0	8	0
All	All	10288	0	9340	205	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 205 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:7:LEU:HB2	1:C:274:LEU:HD11	1.37	1.05
1:D:20:THR:HG22	1:D:23:GLN:H	1.20	1.03
1:B:31:MET:HE3	1:B:269:VAL:HG22	1.43	0.98
1:A:7:LEU:HB2	1:A:274:LEU:HD11	1.47	0.94
1:A:21:ARG:HG2	1:A:21:ARG:HH11	1.33	0.92

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	300/304 (99%)	290 (97%)	10 (3%)	0	100	100
1	B	300/304 (99%)	291 (97%)	9 (3%)	0	100	100
1	C	300/304 (99%)	294 (98%)	6 (2%)	0	100	100
1	D	300/304 (99%)	288 (96%)	12 (4%)	0	100	100
All	All	1200/1216 (99%)	1163 (97%)	37 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	249/251 (99%)	242 (97%)	7 (3%)	43	33
1	B	249/251 (99%)	245 (98%)	4 (2%)	62	58
1	C	249/251 (99%)	241 (97%)	8 (3%)	39	27
1	D	249/251 (99%)	241 (97%)	8 (3%)	39	27
All	All	996/1004 (99%)	969 (97%)	27 (3%)	44	34

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	79	ARG
1	C	290	LYS
1	D	139	TYR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	224	SER
1	C	293	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	292	HIS
1	D	292	HIS
1	C	202	ASN
1	D	216	HIS
1	C	201	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.