



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 17, 2025 – 12:22 pm GMT

PDB ID : 9FVQ  
Title : Ferric-mycobactin receptor (FemA) in complex with pyochelin  
Authors : Moynie, L.  
Deposited on : 2024-06-27  
Resolution : 2.03 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	3.0
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.003 (Gargrove)
Density-Fitness	:	1.0.11
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	2096 (2.04-2.04)
Clashscore	180529	2229 (2.04-2.04)
Ramachandran outliers	177936	2217 (2.04-2.04)
Sidechain outliers	177891	2217 (2.04-2.04)
RSRZ outliers	164620	2096 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	670	<div> <div>3%</div> <div> <div></div> <div>86%</div> <div>11%</div> <div>.</div> </div> </div>
1	B	670	<div> <div>2%</div> <div> <div></div> <div>86%</div> <div>12%</div> <div>.</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	A	812	-	-	X	-
4	EDO	B	820	-	-	X	-

## 2 Entry composition [i](#)


There are 9 unique types of molecules in this entry. The entry contains 21968 atoms, of which 10522 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ferric-mycobactin receptor, FemA.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	670	Total	C	H	N	O	S	133	6	0
			10200	3218	5034	935	1004	9			
1	B	670	Total	C	H	N	O	S	133	16	0
			10285	3242	5080	941	1013	9			

- Molecule 2 is PYOCHELIN FE(III) (three-letter code: 188) (formula:  $C_{14}H_{14}FeN_2O_3S_2$ ).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	Fe	H	N	O	0	1
			36	14	1	14	2	3		
2	B	1	Total	C	Fe	H	N	O	0	1
			36	14	1	14	2	3		

- Molecule 3 is Pyochelin Fe(III) isomer (three-letter code: A1IG8) (formula:  $C_{14}H_{14}FeN_2O_3S_2$ ).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
3	A	1	Total	C	Fe	H	N	O	S	0	1
			35	14	1	13	2	3	2		
3	B	1	Total	C	Fe	H	N	O	S	0	1
			35	14	1	13	2	3	2		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	2	0
			10	2	6	2		
4	A	1	Total	C	H	O	2	0
			10	2	6	2		

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	2	0
			10	2	6	2		
4	A	1	Total	C	H	O	2	0
			10	2	6	2		
4	A	1	Total	C	H	O	1	0
			9	2	5	2		
4	A	1	Total	C	H	O	2	0
			10	2	6	2		
4	A	1	Total	C	H	O	1	0
			9	2	5	2		
4	A	1	Total	C	H	O	2	0
			10	2	6	2		
4	A	1	Total	C	H	O	1	0
			9	2	5	2		
4	A	1	Total	C	H	O	2	0
			10	2	6	2		
4	A	1	Total	C	H	O	2	1
			10	2	6	2		
4	A	1	Total	C	H	O	2	1
			10	2	6	2		
4	B	1	Total	C	H	O	2	0
			10	2	6	2		
4	B	1	Total	C	H	O	2	0
			10	2	6	2		
4	B	1	Total	C	H	O	2	0
			10	2	6	2		
4	B	1	Total	C	H	O	2	0
			10	2	6	2		
4	B	1	Total	C	H	O	2	0
			10	2	6	2		
4	B	1	Total	C	H	O	2	0
			10	2	6	2		
4	B	1	Total	C	H	O	2	0
			10	2	6	2		
4	B	1	Total	C	H	O	2	0
			10	2	6	2		
4	B	1	Total	C	H	O	2	0
			10	2	6	2		
4	B	1	Total	C	H	O	2	1
			10	2	6	2		

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	H	O	2	0
			10	2	6	2		

- Molecule 5 is (HYDROXYETHYLOXY)TRI(ETHYLOXY)OCTANE (three-letter code: C8E) (formula:  $C_{16}H_{34}O_5$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	H	O	2	0
			39	12	25	2		
5	A	1	Total	C	H	O	0	0
			40	12	25	3		
5	A	1	Total	C	H	O	0	0
			40	12	25	3		
5	A	1	Total	C	H	O	0	1
			40	12	25	3		
5	A	1	Total	C	H	O	0	1
			15	4	8	3		
5	B	1	Total	C	H	O	0	1
			15	4	8	3		
5	B	1	Total	C	H	O	2	0
			39	12	25	2		
5	B	1	Total	C	H	O	0	0
			40	12	25	3		
5	B	1	Total	C	H	O	0	1
			26	8	17	1		
5	B	1	Total	C	H	O	0	1
			15	4	8	3		

- Molecule 6 is pentane-2,4-dione (three-letter code: P2D) (formula:  $C_5H_8O_2$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	0	0
			15	5	8	2		
6	B	1	Total	C	H	O	0	0
			15	5	8	2		

- Molecule 7 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	9	Total	K	0	0
			9	9		
7	B	6	Total	K	0	0
			6	6		

- Molecule 8 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula:  $C_2H_6OS$ ).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
8	B	1	Total	C	H	O	S	0	0
			10	2	6	1	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	375	Total	O	0	0
			375	375		
9	B	365	Total	O	0	0
			365	365		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: Ferric-mycobactin receptor, FemA



#### • Molecule 1: Ferric-mycobactin receptor, FemA



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	83.84Å 84.66Å 86.97Å 89.96° 118.67° 113.58°	Depositor
Resolution (Å)	66.72 – 2.03 66.72 – 2.03	Depositor EDS
% Data completeness (in resolution range)	96.6 (66.72-2.03) 96.6 (66.72-2.03)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.42 (at 2.03Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, $R_{free}$	0.182 , 0.223 0.183 , 0.224	Depositor DCC
$R_{free}$ test set	2007 reflections (1.72%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	19.9	Xtriage
Anisotropy	0.280	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.42 , 39.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.000 for -h-l,-k,l	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	21968	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, A1IG8, K, P2D, DMS, 188, C8E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.72	3/5293 (0.1%)	1.24	44/7189 (0.6%)
1	B	0.70	0/5367	1.22	34/7291 (0.5%)
All	All	0.71	3/10660 (0.0%)	1.23	78/14480 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	7
1	B	0	7
All	All	0	14

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	151	GLU	CD-OE2	5.44	1.31	1.25
1	A	712	SER	CA-CB	-5.16	1.45	1.52
1	A	211	GLU	CD-OE1	5.07	1.31	1.25

The worst 5 of 78 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	767	ARG	NE-CZ-NH1	16.49	128.55	120.30
1	B	767	ARG	NE-CZ-NH2	-15.31	112.64	120.30
1	A	767	ARG	NE-CZ-NH2	-13.86	113.37	120.30
1	A	767	ARG	NE-CZ-NH1	12.55	126.58	120.30
1	A	613	ARG	NE-CZ-NH2	10.12	125.36	120.30

There are no chirality outliers.

5 of 14 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	123[A]	ARG	Sidechain
1	A	156	ARG	Sidechain
1	A	453	ARG	Sidechain
1	A	534	ARG	Sidechain
1	A	613	ARG	Sidechain

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5166	5034	5005	55	0
1	B	5205	5080	5013	50	0
2	A	22	14	14	0	0
2	B	22	14	14	0	0
3	A	22	13	0	1	0
3	B	22	13	0	0	0
4	A	48	69	58	7	0
4	B	48	72	65	9	0
5	A	66	108	73	3	0
5	B	52	83	48	3	0
6	A	7	8	8	0	0
6	B	7	8	8	0	0
7	A	9	0	0	0	0
7	B	6	0	0	0	0
8	B	4	6	6	0	0
9	A	375	0	0	11	0
9	B	365	0	0	11	0
All	All	11446	10522	10312	106	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 106 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:451:HIS:ND1	1:B:494:ASP:OD2	1.88	1.06
1:B:152[A]:GLN:OE1	9:B:901:HOH:O	1.71	1.05
1:A:163:GLN:HG2	4:A:812:EDO:H11	1.57	0.85
1:B:609:LEU:HD22	1:B:636[B]:VAL:HG12	1.59	0.83
1:A:220:LYS:HE3	9:A:1035:HOH:O	1.79	0.82

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	674/670 (101%)	656 (97%)	17 (2%)	1 (0%)	48	44
1	B	684/670 (102%)	666 (97%)	16 (2%)	2 (0%)	37	30
All	All	1358/1340 (101%)	1322 (97%)	33 (2%)	3 (0%)	44	38

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	622	SER
1	B	659	PRO
1	A	659	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	539/535 (101%)	528 (98%)	11 (2%)	50	47
1	B	548/535 (102%)	532 (97%)	16 (3%)	37	32
All	All	1087/1070 (102%)	1060 (98%)	27 (2%)	45	38

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	379	ASP
1	B	438	PRO
1	B	600	LEU
1	B	424	TYR
1	B	494	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	344	HIS
1	A	491	ASN
1	A	526	ASN
1	B	120	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 56 ligands modelled in this entry, 15 are monoatomic - leaving 41 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	EDO	A	814	-	3,3,3	0.69	0	2,2,2	0.93	0
5	C8E	A	809	-	14,14,20	0.27	0	13,13,19	0.28	0
4	EDO	A	807	-	3,3,3	0.86	0	2,2,2	0.13	0
4	EDO	A	812	-	3,3,3	0.35	0	2,2,2	0.92	0
4	EDO	A	816	-	3,3,3	0.34	0	2,2,2	0.40	0
4	EDO	B	820	-	3,3,3	0.61	0	2,2,2	0.50	0
5	C8E	B	809	-	13,13,20	0.44	0	12,12,19	0.57	0
4	EDO	A	815	7	3,3,3	0.11	0	2,2,2	0.63	0
4	EDO	B	804	-	3,3,3	1.17	0	2,2,2	0.31	0
4	EDO	B	807	-	3,3,3	0.81	0	2,2,2	0.79	0
4	EDO	B	808	-	3,3,3	0.57	0	2,2,2	0.30	0
8	DMS	B	821	-	3,3,3	0.51	0	3,3,3	0.26	0
4	EDO	B	811	-	3,3,3	0.97	0	2,2,2	0.89	0
4	EDO	A	805	-	3,3,3	0.06	0	2,2,2	0.16	0
4	EDO	A	813	7	3,3,3	0.56	0	2,2,2	0.57	0
4	EDO	A	806	-	3,3,3	0.36	0	2,2,2	0.25	0
5	C8E	B	810	-	14,14,20	0.43	0	13,13,19	0.38	0
4	EDO	A	803	-	3,3,3	0.47	0	2,2,2	0.25	0
4	EDO	B	816	-	3,3,3	0.24	0	2,2,2	0.64	0
4	EDO	B	814	-	3,3,3	0.53	0	2,2,2	0.30	0
5	C8E	A	804	-	13,13,20	0.40	0	12,12,19	0.41	0
4	EDO	B	803	-	3,3,3	0.48	0	2,2,2	0.54	0
4	EDO	B	815	-	3,3,3	0.23	0	2,2,2	0.16	0
4	EDO	A	808	7	3,3,3	0.51	0	2,2,2	0.15	0
6	P2D	B	806	3,2	6,6,6	0.81	0	6,7,7	1.65	2 (33%)
4	EDO	B	805	-	3,3,3	0.37	0	2,2,2	0.46	0
6	P2D	A	811	3,2	6,6,6	0.86	0	6,7,7	1.08	0
5	C8E	A	810	-	14,14,20	0.64	0	13,13,19	0.43	0
4	EDO	B	802	-	3,3,3	0.42	0	2,2,2	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	814	-	-	1/1/1/1	-

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	C8E	A	809	-	-	5/12/12/18	-
4	EDO	A	807	-	-	0/1/1/1	-
4	EDO	A	812	-	-	0/1/1/1	-
4	EDO	A	816	-	-	1/1/1/1	-
4	EDO	B	820	-	-	1/1/1/1	-
5	C8E	B	809	-	-	6/11/11/18	-
4	EDO	A	815	7	-	0/1/1/1	-
4	EDO	B	804	-	-	0/1/1/1	-
4	EDO	B	807	-	-	0/1/1/1	-
4	EDO	B	808	-	-	1/1/1/1	-
4	EDO	B	811	-	-	0/1/1/1	-
4	EDO	A	805	-	-	0/1/1/1	-
4	EDO	A	813	7	-	0/1/1/1	-
4	EDO	A	806	-	-	0/1/1/1	-
5	C8E	B	810	-	-	10/12/12/18	-
4	EDO	A	803	-	-	1/1/1/1	-
4	EDO	B	816	-	-	1/1/1/1	-
4	EDO	B	814	-	-	0/1/1/1	-
5	C8E	A	804	-	-	5/11/11/18	-
4	EDO	B	803	-	-	0/1/1/1	-
4	EDO	B	815	-	-	0/1/1/1	-
4	EDO	A	808	7	-	0/1/1/1	-
6	P2D	B	806	3,2	-	1/4/4/4	-
4	EDO	B	805	-	-	1/1/1/1	-
6	P2D	A	811	3,2	-	2/4/4/4	-
5	C8E	A	810	-	-	3/12/12/18	-
4	EDO	B	802	-	-	0/1/1/1	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	806	P2D	O2-C2-C1	2.55	127.89	121.40
6	B	806	P2D	C1-C2-C3	-2.30	109.94	117.89

There are no chirality outliers.

5 of 39 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	809	C8E	C6-C7-C8-O9

*Continued on next page...*

*Continued from previous page...*





Mol	Chain	Res	Type	Atoms
5	A	810	C8E	O9-C10-C11-O12
5	B	809	C8E	C14-C13-O12-C11
5	A	809	C8E	C3-C4-C5-C6
5	B	810	C8E	C6-C7-C8-O9


There are no ring outliers.

11 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	814	EDO	1	0
4	A	812	EDO	4	0
4	B	820	EDO	5	0
5	B	809	C8E	2	0
4	A	815	EDO	1	0
4	B	804	EDO	3	0
4	B	811	EDO	1	0
4	A	813	EDO	1	0
5	B	810	C8E	1	0
5	A	804	C8E	1	0
5	A	810	C8E	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	670/670 (100%)	-0.45	21 (3%)	51	53	9, 19, 46, 65	3 (0%)
1	B	670/670 (100%)	-0.49	14 (2%)	63	66	7, 19, 43, 67	9 (1%)
All	All	1340/1340 (100%)	-0.47	35 (2%)	57	58	7, 19, 43, 67	12 (0%)

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	340	PRO	4.3
1	A	578	ALA	4.3
1	B	437	GLY	3.9
1	A	340	PRO	3.8
1	B	692	PHE	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
8	DMS	B	821	4/4	0.82	0.20	47,52,59,73	0
4	EDO	A	819[B]	4/4	0.83	0.17	48,50,51,52	6
4	EDO	A	813	4/4	0.84	0.18	40,41,48,48	1
5	C8E	B	809	14/21	0.85	0.16	27,36,45,50	2
4	EDO	B	817[B]	4/4	0.85	0.17	36,45,48,49	2
5	C8E	A	810	15/21	0.86	0.16	37,44,53,56	0
5	C8E	A	817[B]	15/21	0.86	0.20	38,52,63,67	8
7	K	B	827	1/1	0.88	0.37	55,55,55,55	0
5	C8E	B	818[B]	9/21	0.89	0.15	28,35,39,46	17
5	C8E	B	801[B]	7/21	0.89	0.14	35,39,48,50	8
4	EDO	A	815	4/4	0.89	0.18	41,42,48,49	1
7	K	A	829	1/1	0.90	0.35	65,65,65,65	0
5	C8E	B	810	15/21	0.90	0.14	37,43,45,49	0
4	EDO	B	816	4/4	0.90	0.21	56,63,65,66	2
5	C8E	A	820[B]	7/21	0.91	0.14	36,40,49,50	4
5	C8E	A	809	15/21	0.91	0.13	34,39,46,52	0
7	K	A	825	1/1	0.92	0.21	58,58,58,58	0
7	K	A	823	1/1	0.92	0.41	65,65,65,65	0
4	EDO	A	808	4/4	0.93	0.14	27,29,31,32	1
4	EDO	B	820	4/4	0.93	0.16	32,33,33,34	2
4	EDO	B	804	4/4	0.93	0.13	23,30,40,40	2
7	K	B	826	1/1	0.93	0.20	64,64,64,64	0
4	EDO	B	807	4/4	0.93	0.09	27,29,31,32	2
4	EDO	A	818[B]	4/4	0.93	0.16	39,40,54,55	6
4	EDO	A	816	4/4	0.94	0.11	29,33,39,40	2
5	C8E	B	819[B]	7/21	0.94	0.12	31,37,50,57	8
4	EDO	B	808	4/4	0.94	0.10	26,32,41,42	2
4	EDO	A	807	4/4	0.94	0.10	21,27,35,35	2
6	P2D	A	811	7/7	0.95	0.10	23,28,31,32	0
4	EDO	B	805	4/4	0.95	0.09	29,31,43,43	2
4	EDO	A	814	4/4	0.95	0.13	21,35,42,43	2
5	C8E	A	804	14/21	0.95	0.09	19,26,39,41	2
7	K	B	823	1/1	0.95	0.27	49,49,49,49	0
4	EDO	B	802	4/4	0.95	0.12	25,26,29,29	2
4	EDO	B	814	4/4	0.95	0.11	19,23,33,34	2
4	EDO	A	812	4/4	0.95	0.14	22,26,30,31	2
4	EDO	B	811	4/4	0.96	0.08	29,34,35,37	2
7	K	A	824	1/1	0.96	0.16	44,44,44,44	0
7	K	B	824	1/1	0.96	0.23	42,42,42,42	0
4	EDO	A	805	4/4	0.96	0.07	17,24,27,27	2
7	K	A	827	1/1	0.96	0.14	48,48,48,48	0
7	K	A	828	1/1	0.96	0.32	45,45,45,45	0
4	EDO	B	815	4/4	0.97	0.07	17,26,35,35	2

Continued on next page...




*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	A1IG8	B	813[B]	22/22	0.97	0.08	17,21,28,32	35
7	K	B	825	1/1	0.97	0.18	54,54,54,54	0
2	188	B	812[A]	22/22	0.97	0.07	20,22,26,29	36
6	P2D	B	806	7/7	0.97	0.07	21,26,29,32	0
3	A1IG8	A	802[B]	22/22	0.97	0.08	20,24,29,35	35
7	K	A	822	1/1	0.98	0.20	33,33,33,33	0
4	EDO	A	803	4/4	0.98	0.04	13,15,19,19	2
4	EDO	B	803	4/4	0.98	0.07	24,27,28,28	2
2	188	A	801[A]	22/22	0.98	0.07	18,22,24,26	36
7	K	A	826	1/1	0.98	0.21	50,50,50,50	0
4	EDO	A	806	4/4	0.98	0.06	21,22,23,24	2
7	K	A	821	1/1	0.98	0.14	39,39,39,39	0
7	K	B	822	1/1	0.99	0.15	40,40,40,40	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around C8E B 809:**

2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray  
mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around C8E A 810:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around C8E B 810:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)








**Electron density around C8E A 809:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around C8E A 804:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.