



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 1, 2025 – 01:54 pm BST

PDB ID : 9FWY / pdb_00009fwy
Title : Structure of the Nothoceros aenigmaticus LFY DNA-binding domain bound to DNA
Authors : Verhage, L.; Zubieta, C.; Nanao, M.H.; Parcy, F.
Deposited on : 2024-07-01
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0rc1
Xtriage (Phenix) : 2.0rc1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.44


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2335 (2.90-2.90)
Clashscore	180529	2564 (2.90-2.90)
Ramachandran outliers	177936	2514 (2.90-2.90)
Sidechain outliers	177891	2516 (2.90-2.90)
RSRZ outliers	164620	2337 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	169	<div> <div>3%</div> <div>79%</div> <div>16%</div> <div>5%</div> </div>
1	B	169	<div> <div>6%</div> <div>69%</div> <div>23%</div> <div>6%</div> </div>
2	F	29	<div> <div>7%</div> <div>55%</div> <div>45%</div> </div>
3	G	29	<div> <div>28%</div> <div>55%</div> <div>45%</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3812 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Floricaula/leafy-like transcription factor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	160	Total	C	N	O	S	0	0	0
			1309	830	238	231	10			
1	B	159	Total	C	N	O	S	0	0	0
			1304	828	239	227	10			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	178	GLY	-	expression tag	UNP W8EDT4
A	179	ALA	-	expression tag	UNP W8EDT4
A	180	MET	-	expression tag	UNP W8EDT4
A	181	GLU	-	expression tag	UNP W8EDT4
A	346	HIS	-	expression tag	UNP W8EDT4
B	178	GLY	-	expression tag	UNP W8EDT4
B	179	ALA	-	expression tag	UNP W8EDT4
B	180	MET	-	expression tag	UNP W8EDT4
B	181	GLU	-	expression tag	UNP W8EDT4
B	346	HIS	-	expression tag	UNP W8EDT4

- Molecule 2 is a DNA chain called DNA (29-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	F	29	Total	C	N	O	P	0	0	0
			597	280	122	167	28			

- Molecule 3 is a DNA chain called DNA (29-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	29	Total	C	N	O	P	0	0	0
			586	278	103	177	28			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	3	Total 3	O 3	0	0
4	B	3	Total 3	O 3	0	0
4	F	2	Total 2	O 2	0	0
4	G	8	Total 8	O 8	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Floricaula/leafy-like transcription factor



- Molecule 1: Floricaula/leafy-like transcription factor



- Molecule 2: DNA (29-MER)



- Molecule 3: DNA (29-MER)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	43.22Å 90.90Å 152.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	78.00 – 2.90 78.00 – 2.94	Depositor EDS
% Data completeness (in resolution range)	95.0 (78.00-2.90) 95.1 (78.00-2.94)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 2.96Å)	Xtriage
Refinement program	REFMAC v5.8	Depositor
R, R_{free}	0.257 , 0.297 0.274 , 0.313	Depositor DCC
R_{free} test set	642 reflections (4.78%)	wwPDB-VP
Wilson B-factor (Å ²)	61.3	Xtriage
Anisotropy	0.585	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 47.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3812	wwPDB-VP
Average B, all atoms (Å ²)	81.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 47.07 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0466e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.45	0/1340	0.87	0/1798
1	B	0.46	0/1336	0.90	0/1791
2	F	0.27	0/672	0.81	1/1036 (0.1%)
3	G	0.27	0/654	0.81	0/1007
All	All	0.40	0/4002	0.86	1/5632 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	F	12	DC	O3'-P-O5'	-5.04	96.44	104.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	290	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1309	0	1309	17	0
1	B	1304	0	1308	29	0
2	F	597	0	322	7	0
3	G	586	0	327	13	0
4	A	3	0	0	0	0
4	B	3	0	0	0	0
4	F	2	0	0	0	0
4	G	8	0	0	0	0
All	All	3812	0	3266	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:280:SER:O	1:B:284:ARG:HG3	1.74	0.87
1:B:188:ARG:HH21	1:B:257:TYR:HB3	1.43	0.83
1:B:334:LYS:O	1:B:338:LEU:HD23	1.81	0.80
3:G:15:DC:H2'	3:G:16:DT:H72	1.63	0.79
1:B:315:ASP:O	1:B:319:LEU:HD12	1.91	0.70
3:G:15:DC:H2'	3:G:16:DT:C7	2.23	0.68
1:B:281:ASP:HA	1:B:284:ARG:HD3	1.77	0.66
2:F:16:DC:H2''	2:F:17:DG:C8	2.31	0.65
3:G:22:DC:H2'	3:G:23:DC:C6	2.32	0.65
2:F:27:DC:H2''	2:F:28:DA:C8	2.33	0.64
1:A:218:GLU:OE2	1:A:325:LYS:HD3	1.99	0.62
1:B:188:ARG:NH2	1:B:257:TYR:HB3	2.16	0.60
3:G:16:DT:H2''	3:G:17:DG:C8	2.36	0.60
1:B:194:PHE:CE2	1:B:213:LEU:HB2	2.38	0.59
3:G:22:DC:H2''	3:G:23:DC:O5'	2.03	0.57
1:B:278:ASP:O	1:B:282:GLU:HG2	2.05	0.56
2:F:7:DG:H2''	2:F:8:DG:N7	2.25	0.52
1:B:303:LEU:N	1:B:304:PRO:HD2	2.25	0.52
1:A:197:THR:HG23	1:A:201:GLU:HB3	1.93	0.51
1:B:281:ASP:HA	1:B:284:ARG:CD	2.41	0.51
2:F:3:DC:H2''	2:F:4:DA:C8	2.46	0.51
3:G:13:DC:H2''	3:G:14:DG:C8	2.46	0.51
1:A:316:ILE:HD12	1:A:319:LEU:HD12	1.93	0.51
1:B:316:ILE:HA	1:B:319:LEU:HD13	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:G:22:DC:H2'	3:G:23:DC:C5	2.47	0.50
1:B:283:LEU:HD11	1:B:300:ALA:HB1	1.93	0.50
1:B:198:GLU:HG2	1:B:199:PRO:HD2	1.92	0.49
1:A:270:TYR:OH	1:A:325:LYS:CD	2.60	0.49
3:G:22:DC:C2'	3:G:23:DC:C6	2.96	0.49
1:B:260:LYS:HB3	1:B:261:PRO:HD3	1.96	0.47
1:A:260:LYS:HB3	1:A:261:PRO:HD3	1.96	0.47
1:A:270:TYR:OH	1:A:325:LYS:HD3	2.14	0.47
1:B:293:ASN:HB2	2:F:7:DG:OP1	2.15	0.46
3:G:15:DC:H2'	3:G:16:DT:C5	2.49	0.46
1:A:259:ASN:O	1:A:263:MET:HG2	2.16	0.46
2:F:13:DC:H2''	2:F:14:DA:C8	2.51	0.46
1:A:263:MET:O	1:A:267:VAL:HG22	2.16	0.46
1:A:194:PHE:CE1	1:A:213:LEU:HB2	2.51	0.46
1:A:271:ALA:HA	1:A:326:LEU:HD21	1.97	0.45
3:G:1:DG:H2'	3:G:2:DT:H72	1.98	0.45
1:A:279:LYS:HA	1:A:282:GLU:HG2	1.97	0.45
1:B:259:ASN:O	1:B:263:MET:HG2	2.16	0.45
1:B:271:ALA:HA	1:B:326:LEU:HD21	1.98	0.45
1:A:303:LEU:HD12	1:A:303:LEU:H	1.81	0.44
3:G:27:DG:H2''	3:G:28:DG:C8	2.52	0.44
1:B:210:LEU:O	1:B:213:LEU:HB3	2.18	0.44
1:B:263:MET:O	1:B:267:VAL:HG22	2.17	0.44
1:B:214:PHE:CE2	1:B:273:HIS:CG	3.06	0.44
1:B:210:LEU:HB2	1:B:284:ARG:NH2	2.34	0.43
1:A:255:ALA:HA	1:A:257:TYR:CE2	2.54	0.43
1:A:210:LEU:O	1:A:213:LEU:HB3	2.18	0.42
1:B:197:THR:HG22	1:B:198:GLU:N	2.33	0.42
1:B:303:LEU:H	1:B:303:LEU:HD12	1.85	0.42
3:G:22:DC:H4'	3:G:23:DC:OP1	2.18	0.42
1:B:265:GLN:HE21	1:B:266:TYR:HE1	1.68	0.42
1:B:255:ALA:HA	1:B:257:TYR:CE2	2.55	0.42
1:A:303:LEU:HB2	1:A:304:PRO:HD3	2.02	0.41
1:B:213:LEU:HD23	1:B:214:PHE:CE1	2.56	0.41
1:A:270:TYR:OH	1:A:325:LYS:HG2	2.21	0.41
1:B:323:HIS:CD2	1:B:325:LYS:HB2	2.55	0.41
1:A:336:ILE:CD1	1:B:237:LYS:HG2	2.51	0.41
3:G:24:DC:H2''	3:G:25:DG:C8	2.56	0.41
1:B:325:LYS:N	1:B:325:LYS:CD	2.84	0.40
2:F:22:DG:H1'	2:F:23:DT:C5'	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	158/169 (94%)	151 (96%)	6 (4%)	1 (1%)	22	52
1	B	157/169 (93%)	149 (95%)	8 (5%)	0	100	100
All	All	315/338 (93%)	300 (95%)	14 (4%)	1 (0%)	37	66

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	199	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	140/145 (97%)	138 (99%)	2 (1%)	62	86
1	B	139/145 (96%)	133 (96%)	6 (4%)	25	57
All	All	279/290 (96%)	271 (97%)	8 (3%)	37	72

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	188	ARG
1	A	294	VAL
1	B	248	ARG
1	B	252	VAL
1	B	283	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	303	LEU
1	B	331	VAL
1	B	338	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	208	ASN
1	A	253	GLN
1	A	265	GLN
1	B	253	GLN
1	B	340	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	160/169 (94%)	0.45	5 (3%) 51 46	36, 57, 86, 123	0
1	B	159/169 (94%)	0.66	10 (6%) 27 23	40, 79, 124, 136	0
2	F	29/29 (100%)	1.11	2 (6%) 24 20	55, 97, 157, 174	0
3	G	29/29 (100%)	1.27	8 (27%) 2 2	53, 91, 166, 181	0
All	All	377/396 (95%)	0.65	25 (6%) 26 22	36, 67, 129, 181	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	G	29	DC	4.1
3	G	28	DG	3.4
3	G	4	DC	3.3
1	B	202	LEU	3.0
2	F	1	DG	3.0
1	A	345	LYS	2.9
1	A	292	GLU	2.7
1	A	202	LEU	2.6
1	B	344	SER	2.5
1	B	296	ALA	2.5
3	G	3	DG	2.3
3	G	27	DG	2.3
3	G	1	DG	2.3
3	G	5	DT	2.3
1	B	301	CYS	2.3
1	B	197	THR	2.2
1	A	286	CYS	2.2
1	B	203	ALA	2.2
1	B	286	CYS	2.2
3	G	20	DC	2.1
1	A	339	CYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	205	GLY	2.1
1	B	204	ARG	2.1
2	F	21	DG	2.0
1	B	314	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.