



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 13, 2024 – 08:26 AM EDT

PDB ID : 4G44
Title : Structure of P450 CYP121 in complex with lead compound MB286, 3-((1H-1,2,4-triazol-1-yl)methyl)aniline
Authors : Hudson, S.A.; McLean, K.J.; Surade, S.; Yang, Y.-Q.; Leys, D.; Ciulli, A.; Munro, A.W.; Abell, C.
Deposited on : 2012-07-16
Resolution : 1.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.36.2
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2024 (1.28-1.20)
Clashscore	141614	1007 (1.26-1.22)
Ramachandran outliers	138981	2053 (1.28-1.20)
Sidechain outliers	138945	2051 (1.28-1.20)
RSRZ outliers	127900	1987 (1.28-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	396	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3990 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome P450 121.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	395	Total	C	N	O	S	0	30	1
			3074	1970	530	562	12			

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is 3-(1H-1,2,4-triazol-1-ylmethyl)aniline (three-letter code: TZM) (formula: $C_9H_{10}N_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			13	9	4		
4	A	1	Total	C	N	0	0
			13	9	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	832	Total	O	0	0
			832	832		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cytochrome P450 121



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	77.64Å 77.64Å 264.10Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.11 – 1.24 47.11 – 1.24	Depositor EDS
% Data completeness (in resolution range)	96.6 (47.11-1.24) 96.6 (47.11-1.24)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.19 (at 1.24Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.161 , 0.179 0.160 , 0.178	Depositor DCC
R_{free} test set	6402 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	8.4	Xtriage
Anisotropy	0.029	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 58.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3990	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, SO4, TZM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	2.78	31/3290 (0.9%)	1.82	63/4475 (1.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (31) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	64[A]	GLU	CG-CD	64.03	2.48	1.51
1	A	64[B]	GLU	CG-CD	64.03	2.48	1.51
1	A	194[A]	GLU	CG-CD	63.12	2.46	1.51
1	A	194[B]	GLU	CG-CD	63.12	2.46	1.51
1	A	33[A]	LYS	CD-CE	22.63	2.07	1.51
1	A	33[B]	LYS	CD-CE	22.63	2.07	1.51
1	A	81	GLU	CD-OE1	9.39	1.35	1.25
1	A	56	GLU	CD-OE2	7.96	1.34	1.25
1	A	272	GLU	CD-OE2	7.87	1.34	1.25
1	A	396	TRP	CG-CD1	7.63	1.47	1.36
1	A	326	GLU	CD-OE1	7.40	1.33	1.25
1	A	330	PRO	N-CD	-7.15	1.37	1.47
1	A	32	ARG	CZ-NH2	-6.86	1.24	1.33
1	A	62	MET	CB-CG	6.61	1.72	1.51
1	A	26	ARG	CZ-NH1	-6.52	1.24	1.33
1	A	56	GLU	CG-CD	-6.48	1.42	1.51
1	A	35	ARG	CZ-NH2	6.39	1.41	1.33
1	A	279[A]	SER	CA-CB	6.07	1.62	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	279[B]	SER	CA-CB	6.07	1.62	1.52
1	A	254	GLN	CD-OE1	6.01	1.37	1.24
1	A	303	GLU	CG-CD	-6.00	1.43	1.51
1	A	326	GLU	CD-OE2	-5.92	1.19	1.25
1	A	310	GLU	CD-OE1	5.74	1.31	1.25
1	A	205	GLU	CD-OE1	-5.55	1.19	1.25
1	A	32	ARG	NE-CZ	5.44	1.40	1.33
1	A	81	GLU	CD-OE2	5.37	1.31	1.25
1	A	58	ARG	CZ-NH1	5.15	1.39	1.33
1	A	216[A]	SER	CB-OG	-5.09	1.35	1.42
1	A	216[B]	SER	CB-OG	-5.09	1.35	1.42
1	A	208[A]	ARG	CZ-NH2	-5.04	1.26	1.33
1	A	208[B]	ARG	CZ-NH2	-5.04	1.26	1.33

All (63) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	64[A]	GLU	CG-CD-OE2	-26.19	65.93	118.30
1	A	64[B]	GLU	CG-CD-OE2	-26.19	65.93	118.30
1	A	64[A]	GLU	CG-CD-OE1	25.11	168.53	118.30
1	A	64[B]	GLU	CG-CD-OE1	25.11	168.53	118.30
1	A	33[A]	LYS	CD-CE-NZ	25.11	169.44	111.70
1	A	33[B]	LYS	CD-CE-NZ	25.11	169.44	111.70
1	A	35	ARG	NE-CZ-NH1	24.30	132.45	120.30
1	A	252	ARG	NE-CZ-NH2	13.93	127.27	120.30
1	A	352	ARG	NE-CZ-NH1	-13.21	113.70	120.30
1	A	56	GLU	OE1-CD-OE2	11.51	137.12	123.30
1	A	370	ASP	CB-CG-OD2	-11.36	108.08	118.30
1	A	329	ARG	NE-CZ-NH2	11.12	125.86	120.30
1	A	64[A]	GLU	CB-CG-CD	-11.04	84.40	114.20
1	A	64[B]	GLU	CB-CG-CD	-11.04	84.40	114.20
1	A	286	ARG	NE-CZ-NH1	10.78	125.69	120.30
1	A	275	ARG	NE-CZ-NH2	10.30	125.45	120.30
1	A	35	ARG	NE-CZ-NH2	-10.13	115.23	120.30
1	A	33[A]	LYS	CG-CD-CE	-9.61	83.07	111.90
1	A	33[B]	LYS	CG-CD-CE	-9.61	83.07	111.90
1	A	122	ASP	CB-CG-OD1	9.45	126.81	118.30
1	A	291	ASP	CB-CG-OD1	9.27	126.65	118.30
1	A	194[A]	GLU	CB-CG-CD	-8.88	90.22	114.20
1	A	194[B]	GLU	CB-CG-CD	-8.88	90.22	114.20
1	A	286	ARG	NE-CZ-NH2	-8.40	116.10	120.30
1	A	194[A]	GLU	CG-CD-OE2	8.39	135.08	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	194[B]	GLU	CG-CD-OE2	8.39	135.08	118.30
1	A	383	ARG	NE-CZ-NH2	8.39	124.49	120.30
1	A	194[A]	GLU	CG-CD-OE1	-8.24	101.83	118.30
1	A	194[B]	GLU	CG-CD-OE1	-8.24	101.83	118.30
1	A	352	ARG	NH1-CZ-NH2	8.12	128.33	119.40
1	A	303	GLU	OE1-CD-OE2	-7.87	113.86	123.30
1	A	383	ARG	NE-CZ-NH1	-7.74	116.43	120.30
1	A	370	ASP	CB-CG-OD1	7.73	125.26	118.30
1	A	396	TRP	CD1-NE1-CE2	7.61	115.85	109.00
1	A	328	ASP	CB-CG-OD2	7.25	124.83	118.30
1	A	32	ARG	NE-CZ-NH1	7.17	123.89	120.30
1	A	208[A]	ARG	CD-NE-CZ	7.08	133.51	123.60
1	A	208[B]	ARG	CD-NE-CZ	7.08	133.51	123.60
1	A	183	ASP	CB-CG-OD1	6.80	124.42	118.30
1	A	387	ARG	NE-CZ-NH1	6.71	123.65	120.30
1	A	99[A]	MET	CB-CA-C	-6.66	97.08	110.40
1	A	99[B]	MET	CB-CA-C	-6.66	97.08	110.40
1	A	113	LEU	CA-CB-CG	6.61	130.50	115.30
1	A	139	ASP	CB-CG-OD2	-6.60	112.36	118.30
1	A	35	ARG	NH1-CZ-NH2	-6.50	112.25	119.40
1	A	139	ASP	CB-CG-OD1	6.47	124.12	118.30
1	A	208[A]	ARG	CA-CB-CG	6.43	127.54	113.40
1	A	208[B]	ARG	CA-CB-CG	6.43	127.54	113.40
1	A	56	GLU	CG-CD-OE2	-6.41	105.48	118.30
1	A	376	ASP	CB-CG-OD1	6.26	123.94	118.30
1	A	252	ARG	NE-CZ-NH1	-6.18	117.21	120.30
1	A	58	ARG	NE-CZ-NH2	6.09	123.35	120.30
1	A	291	ASP	CB-CG-OD2	-6.05	112.86	118.30
1	A	137	PHE	CB-CG-CD2	-6.04	116.58	120.80
1	A	134[A]	ARG	NE-CZ-NH2	5.88	123.24	120.30
1	A	134[B]	ARG	NE-CZ-NH2	5.88	123.24	120.30
1	A	362[A]	LEU	CB-CA-C	5.86	121.32	110.20
1	A	362[B]	LEU	CB-CA-C	5.86	121.32	110.20
1	A	183	ASP	CB-CG-OD2	-5.75	113.13	118.30
1	A	371	LEU	CB-CG-CD2	5.71	120.71	111.00
1	A	300[A]	ARG	CG-CD-NE	5.37	123.08	111.80
1	A	300[B]	ARG	CG-CD-NE	5.37	123.08	111.80
1	A	173	ASP	CB-CG-OD1	-5.07	113.73	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	72	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3074	0	3088	40	1
2	A	43	0	30	4	0
3	A	15	0	0	1	0
4	A	26	0	20	5	0
5	A	832	0	0	28	3
All	All	3990	0	3138	47	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:37[A]:ILE:HD11	5:A:1236:HOH:O	1.58	1.02
1:A:383:ARG:NE	5:A:1236:HOH:O	1.96	0.97
1:A:317:GLU:CG	5:A:1250:HOH:O	2.21	0.88
4:A:406:TZM:CAE	5:A:839:HOH:O	2.22	0.87
1:A:390:GLU:HB2	5:A:1312:HOH:O	1.75	0.86
4:A:406:TZM:NAI	5:A:839:HOH:O	2.12	0.82
1:A:111:GLN:NE2	1:A:114[B]:ARG:HH21	1.78	0.81
1:A:37[A]:ILE:CD1	5:A:1236:HOH:O	2.25	0.76
1:A:99[A]:MET:SD	5:A:707:HOH:O	2.44	0.76
1:A:383:ARG:CD	5:A:1236:HOH:O	2.29	0.75
3:A:404:SO4:O3	5:A:1137:HOH:O	2.05	0.74
1:A:391:ARG:NH2	5:A:1312:HOH:O	2.23	0.72
1:A:86[B]:MET:SD	5:A:1190:HOH:O	2.47	0.71
1:A:17:ARG:NE	5:A:1055:HOH:O	2.23	0.71
1:A:340[A]:ARG:NH2	5:A:1023:HOH:O	2.21	0.70
1:A:29:GLU:HG2	1:A:32:ARG:HB3	1.75	0.68
1:A:58:ARG:HG2	5:A:1137:HOH:O	1.95	0.67
1:A:353:ARG:CG	5:A:1018:HOH:O	2.43	0.66

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41[B]:GLU:OE2	5:A:789:HOH:O	2.14	0.66
1:A:277:ASN:HB2	5:A:711:HOH:O	1.96	0.64
1:A:86[A]:MET:HG3	2:A:401:HEM:HBD2	1.82	0.60
1:A:277:ASN:ND2	5:A:1132:HOH:O	2.34	0.60
1:A:29:GLU:OE1	5:A:1032:HOH:O	2.18	0.57
1:A:383:ARG:HG2	5:A:1236:HOH:O	2.06	0.56
4:A:406:TZM:H9	5:A:839:HOH:O	1.94	0.54
1:A:29:GLU:OE1	5:A:1139:HOH:O	2.18	0.54
1:A:331:ASN:H	1:A:331:ASN:HD22	1.55	0.52
1:A:144:ALA:HA	1:A:154:GLN:HE22	1.74	0.51
1:A:205:GLU:OE1	1:A:208[A]:ARG:HD2	2.11	0.51
1:A:86[A]:MET:CE	5:A:809:HOH:O	2.58	0.50
1:A:56:GLU:OE2	1:A:340[B]:ARG:HD2	2.11	0.50
1:A:74:ASN:C	1:A:74:ASN:HD22	2.14	0.49
1:A:246:LEU:HD22	1:A:362[B]:LEU:CD2	2.43	0.49
4:A:406:TZM:NAA	5:A:594:HOH:O	2.11	0.48
1:A:271[A]:GLU:OE2	1:A:352:ARG:NE	2.35	0.47
1:A:37[B]:ILE:HG22	5:A:1030:HOH:O	2.13	0.47
1:A:390:GLU:CB	5:A:1312:HOH:O	2.48	0.47
1:A:86[A]:MET:HG3	2:A:401:HEM:CBD	2.45	0.46
1:A:205:GLU:O	1:A:208[A]:ARG:HG2	2.17	0.44
1:A:352:ARG:HH11	1:A:352:ARG:HD2	1.52	0.43
2:A:401:HEM:HMC1	2:A:401:HEM:HBC2	2.00	0.43
1:A:104:PRO:HA	1:A:109[B]:LEU:HD23	2.01	0.43
2:A:401:HEM:C4D	4:A:405:TZM:CAG	3.02	0.43
1:A:84:ASN:OD1	5:A:1147:HOH:O	2.22	0.41
1:A:331:ASN:HD22	1:A:331:ASN:N	2.17	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:1267:HOH:O	5:A:1290:HOH:O[5_555]	2.05	0.15
5:A:936:HOH:O	5:A:936:HOH:O[10_555]	2.13	0.07
1:A:254:GLN:CG	5:A:1120:HOH:O[5_555]	2.19	0.01

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	423/396 (107%)	417 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	337/326 (103%)	331 (98%)	6 (2%)	59	21

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	ARG
1	A	74	ASN
1	A	277	ASN
1	A	300[A]	ARG
1	A	300[B]	ARG
1	A	331	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	ASN
1	A	111	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	154	GLN
1	A	251	GLN
1	A	331	ASN
1	A	342	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no monosaccharides in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	401	1,4	42,50,50	1.80	10 (23%)	46,82,82	1.56	11 (23%)
4	TZM	A	405	2	12,14,14	2.30	5 (41%)	15,18,18	2.40	8 (53%)
3	SO4	A	402	-	4,4,4	1.98	1 (25%)	6,6,6	0.92	0
3	SO4	A	404	-	4,4,4	1.61	1 (25%)	6,6,6	1.66	2 (33%)
3	SO4	A	403	-	4,4,4	0.55	0	6,6,6	0.48	0
4	TZM	A	406	-	12,14,14	1.95	4 (33%)	15,18,18	2.17	7 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TZM	A	405	2	-	1/4/4/4	0/2/2/2
4	TZM	A	406	-	-	0/4/4/4	0/2/2/2
2	HEM	A	401	1,4	-	0/12/54/54	-

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	405	TZM	NAJ-NAM	-5.06	1.28	1.35
2	A	401	HEM	O2D-CGD	-4.70	1.15	1.30
4	A	405	TZM	CAF-NAM	4.36	1.38	1.33
4	A	406	TZM	NAJ-NAM	-4.10	1.29	1.35
2	A	401	HEM	CBD-CGD	3.87	1.59	1.50
2	A	401	HEM	CHB-C1B	3.56	1.43	1.34
2	A	401	HEM	CHA-C4D	3.48	1.43	1.34
3	A	402	SO4	O2-S	-3.37	1.24	1.44
3	A	404	SO4	O4-S	-3.02	1.23	1.48
4	A	406	TZM	CAH-CAL	-2.96	1.43	1.51
2	A	401	HEM	C3B-C4B	-2.77	1.39	1.44
4	A	405	TZM	CAG-CAL	2.42	1.43	1.39
4	A	406	TZM	CAG-CAL	2.37	1.43	1.39
2	A	401	HEM	C3D-C2D	2.32	1.41	1.36
4	A	406	TZM	CAG-CAK	2.30	1.43	1.39
2	A	401	HEM	C1B-NB	-2.29	1.36	1.40
2	A	401	HEM	C3B-C2B	2.26	1.41	1.37
4	A	405	TZM	CAK-NAA	2.16	1.45	1.38
2	A	401	HEM	CBD-CAD	-2.12	1.44	1.51
4	A	405	TZM	CAH-NAM	-2.10	1.43	1.47
2	A	401	HEM	CMB-C2B	-2.03	1.46	1.50

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	405	TZM	CAL-CAH-NAM	4.06	117.79	111.94
4	A	405	TZM	CAH-CAL-CAD	-4.01	112.56	120.39
4	A	405	TZM	NAI-CAF-NAM	-3.99	107.41	112.24
2	A	401	HEM	C4A-C3A-C2A	-3.82	104.34	107.00
4	A	406	TZM	NAI-CAF-NAM	-3.50	108.01	112.24
4	A	406	TZM	CAC-CAK-CAG	3.34	122.91	118.62
4	A	406	TZM	CAH-NAM-NAJ	3.30	126.05	120.87

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	HEM	O2D-CGD-O1D	3.22	131.60	123.33
2	A	401	HEM	C3B-C2B-C1B	-3.20	104.01	106.41
3	A	404	SO4	O4-S-O3	2.94	124.76	108.54
4	A	406	TZM	CAG-CAK-NAA	-2.94	115.60	120.56
4	A	406	TZM	CAF-NAI-CAE	2.82	107.07	102.62
4	A	405	TZM	CAB-CAD-CAL	-2.78	116.71	120.61
4	A	406	TZM	CAK-CAG-CAL	-2.72	118.37	121.01
4	A	405	TZM	CAH-CAL-CAG	2.67	125.44	120.23
2	A	401	HEM	O1D-CGD-CBD	-2.62	114.79	123.09
2	A	401	HEM	C3D-C4D-ND	2.54	112.96	110.17
4	A	406	TZM	CAL-CAH-NAM	-2.50	108.34	111.94
4	A	405	TZM	CAD-CAL-CAG	2.46	121.94	118.55
3	A	404	SO4	O3-S-O2	-2.39	97.04	109.56
2	A	401	HEM	CHA-C4D-C3D	-2.39	120.82	125.23
2	A	401	HEM	C3B-C4B-NB	2.33	111.14	109.47
4	A	405	TZM	CAK-CAG-CAL	-2.31	118.77	121.01
2	A	401	HEM	CMA-C3A-C2A	2.31	129.29	124.94
2	A	401	HEM	C2B-C1B-NB	2.23	112.41	109.84
2	A	401	HEM	C4C-CHD-C1D	2.16	125.41	122.56
2	A	401	HEM	C4B-CHC-C1C	2.06	125.28	122.56
4	A	405	TZM	CAD-CAB-CAC	2.01	122.83	120.24

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	405	TZM	CAL-CAH-NAM-CAF


There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	HEM	4	0
4	A	405	TZM	1	0
3	A	404	SO4	1	0
4	A	406	TZM	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	395/396 (99%)	-0.20	4 (1%) 82 78	5, 9, 21, 44	4 (1%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	THR	6.3
1	A	126	THR	3.0
1	A	107	PRO	2.9
1	A	3	ALA	2.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	TZM	A	406	13/13	0.88	0.12	17,21,24,29	0
4	TZM	A	405	13/13	0.95	0.08	9,12,19,20	0
3	SO4	A	403	5/5	0.96	0.11	15,15,18,20	5

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	SO4	A	402	5/5	0.97	0.14	14,21,23,28	0
3	SO4	A	404	5/5	0.98	0.08	11,14,17,18	0
2	HEM	A	401	43/43	0.99	0.06	5,7,11,13	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.